

Closing Tues: HW 9.5

Closing Thurs: HW 9.6, HW 9.7(1)

Monday is a holiday (no class, no MSC)!

9.5/9.6 Product, Quotient, Chain rules

Consider the three functions:

$$y = (x^5 + 4x + 7)(x^4 + 2x) \quad \left. \vphantom{y} \right\} \text{PRODUCT}$$
$$= f(x)g(x)$$

$$y = \frac{x^4 + 5x}{x^7 - x^2} = \frac{f(x)}{g(x)} \quad \left. \vphantom{y} \right\} \text{QUOTIENT}$$

$$y = (4x^2 - 3x)^{10} = f(g(x)) \quad \left. \vphantom{y} \right\} \text{COMPOSITION}$$

$$f(x) = x^5 + 4x + 7 = \text{First}$$
$$g(x) = x^4 + 2x = \text{Second}$$

$$f'(x) = 5x^4 + 4$$

$$g'(x) = 4x^3 + 2$$

$$f(x) = x^4 + 5x = \text{NUMERATOR}$$

$$g(x) = x^7 - x^2 = \text{DENOMINATOR}$$

$$f'(x) = 4x^3 + 5$$

$$g'(x) = 7x^6 - 2x$$

$$f(u) = u^{10} \Rightarrow f'(u) = 10u^9$$

$$u = g(x) = 4x^2 - 3x \Rightarrow g'(x) = 8x - 3$$

THE PRODUCT, QUOTIENT, AND
CHAIN RULES TELL US WHAT TO
DO NEXT.

$$\text{PRODUCT RULE: } \frac{d}{dx}(f(x)g(x)) = f(x)g'(x) + f'(x)g(x).$$

$$F \quad S \qquad F \quad S' + F' \quad S$$

Ex $y = \underbrace{(x^5 + 4x + 7)}_F \underbrace{(x^4 + 2x)}_S \Rightarrow y' = \underbrace{(x^5 + 4x + 7)}_F \underbrace{(4x^3 + 2)}_{S'} + \underbrace{(5x^4 + 4)}_{F'} \underbrace{(x^4 + 2x)}_S$

Ex $y = \underbrace{x^3}_F \underbrace{(x^2 + x^{10})}_S \Rightarrow y' = \underbrace{x^3}_F \underbrace{(2x + 10x^9)}_{S'} + \underbrace{3x^2}_{F'} \underbrace{(x^2 + x^{10})}_S$
 $= 2x^4 + 10x^{12} + 3x^4 + 3x^{12} = 5x^4 + 13x^{12}$ ✓

OR, EXPAND FIRST

$y = x^3 x^2 + x^3 x^{10} = x^5 + x^{13} \Rightarrow y' = 5x^4 + 13x^{12}$ ✓

proof of product rule

(just for your own interest)

We are trying to find a pattern for

$$\frac{f(x+h)g(x+h) - f(x)g(x)}{h}$$

Adding and subtracting

$f(x+h)g(x)$ in the numerator gives

$$\frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h}$$

Then rearranging gives

$$\begin{aligned} &= \frac{f(x+h)g(x+h) - f(x+h)g(x) + f(x+h)g(x) - f(x)g(x)}{h} \\ &= f(x+h) \frac{g(x+h) - g(x)}{h} + \frac{f(x+h) - f(x)}{h} g(x) \end{aligned}$$

As $h \rightarrow 0$, we see the expression above is approaching

$$f(x)g'(x) + f'(x)g(x)$$

QUOTIENT RULE:

$$\frac{d}{dx} \left(\frac{f(x)}{g(x)} \right) = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}$$

$$\left(\frac{N}{D} \right)' = \frac{D \cdot N' - N \cdot D'}{D^2}$$

$$\text{EX)} \quad y = \frac{x^4 + 5x}{x^7 - x^2} \Rightarrow y' = \frac{(x^7 - x^2)(4x^3 + 5) - (x^4 + 5x)(7x^6 - 2x)}{(x^7 - x^2)^2}$$

$$\begin{aligned} \text{EX)} \quad y = \frac{4x + 2}{x^3} &\Rightarrow y' = \frac{x^3(4) - (4x + 2)3x^2}{x^6} = \frac{4x^3 - 12x^3 - 6x^2}{x^6} \\ &= \frac{-8x^3 - 6x^2}{x^6} = -8x^{-3} - 6x^{-4} \end{aligned}$$

OR, EXPAND FIRST

$$\begin{aligned} y = \frac{1}{x^3}(4x + 2) &= \frac{4x}{x^3} + \frac{2}{x^3} = 4x^{-2} + 2x^{-3} \Rightarrow y' = -8x^{-3} - 6x^{-4} \\ &= -\frac{8}{x^3} - \frac{6}{x^4} \end{aligned} \quad \leftarrow \text{ONLY POSITIVE EXPONENTS}$$

You try: Differentiate

1. $y = x^2(x^3 + 1)$

EXPANDING FIRST $\Rightarrow y = x^5 + x^2$
 $y' = 5x^4 + 2x$

PRODUCT RULE $\Rightarrow y' = x^2 \cdot 3x^2 + 2x(x^3 + 1)$
 $= 3x^4 + 2x^4 + 2x$
 $= 5x^4 + 2x$

2. $y = \frac{5}{x^3}$

REWRITE FIRST, $y = 5x^{-3}$
 $y' = -15x^{-4} = -\frac{15}{x^4}$

QUOTIENT RULE $y' = \frac{x^3(0) - 5 \cdot 3x^2}{x^6}$
 $= \frac{-15x^2}{x^6} = -15x^{-4}$

HW NOTE

ANSWER WRITTEN
USING ONLY POSITIVE
EXPONENTS

$$3. y = (x^2 + 3x)(\sqrt{x} - 5x^3)$$

PRODUCT RULE

$$y' = (x^2 + 3x) \left(\frac{1}{2}x^{-\frac{1}{2}} - 15x^2 \right) + (2x + 3)(x^{\frac{1}{2}} - 5x^3)$$

or

$$y = x^{2.5} - 5x^5 + 3x^{1.5} - 15x^4$$

$$y' = 2.5x^{1.5} - 25x^4 + 4.5x^{0.5} - 60x^3$$

$$4. y = \frac{x^5}{3x^3 - x^5}$$

QUOTIENT RULE (ONLY OPTION)

$$y' = \frac{(3x^3 - x^5)5x^4 - x^5(9x^2 - 5x^4)}{(3x^3 - x^5)^2}$$

Equations for Tangent lines

(HW 9.5: Problems 8 and 9)

Recall: All the points (x, y) on a given line can be described by an equation of the form

$$y = m(x - x_0) + y_0$$

where

m = slope of the line


(x_0, y_0) = any point on the line

Review Question:

Find the equation of the line that has slope 8 and goes through $(3, 7)$.

$$y = 8(x - 3) + 7$$

$$y = m(x - x_0) + y_0$$



means

$$\frac{y - 7}{x - 3} = 8$$

For ALL (x, y) ON THE LINE.

CAN ALSO BE EXPANDED TO GET

$$y = 8x - 24 + 7$$

$$y = 8x - 17$$

$$y = mx + b$$

Since $f'(x)$ is the slope of the tangent line, we can use it to get the equation for the tangent line.

Example: Let

$$f(x) = \frac{x^3 + 3}{2x - 1}$$


Find the equation for the tangent line at $x = 1$.

$$y = f(x) = \frac{x^3 + 3}{2x - 1} = \text{HEIGHT ON GRAPH}$$

$$y' = f'(x) = \frac{(2x-1)(3x^2) - (x^3+3) \cdot 2}{(2x-1)^2} = \text{TANGENT SLOPE ON GRAPH}$$

$$f(1) = \frac{1^3 + 3}{2(1) - 1} = 4$$

$$f'(1) = \frac{(2(1)-1)(3(1)^2) - ((1^3+3) \cdot 2)}{(2(1)-1)^2} = \frac{1 \cdot 3 - 4 \cdot 2}{1^2} = \frac{3 - 8}{1} = -5$$



$$y = -5(x-1) + 4$$

$$y = -5x + 5 + 4$$

$$y = -5x + 9$$

Section 9.6: The chain rule

CHAIN RULE:

$$\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$$

EX) $y = (4x^2 - 3x)^{10} = f(g(x))$

OUTSIDE = $f(u) = u^{10} \Rightarrow 10u^9$

INSIDE = $g(x) = 4x^2 - 3x \Rightarrow 8x - 3$

$$y' = 10(4x^2 - 3x)^9 \cdot (8x - 3)$$

EX) $y = (6x + x^{14})^{100}$

$$y' = 100(6x + x^{14})^{99} \cdot (6 + 14x^{13})$$

EX) $y = \sqrt{x^2 + 3x} = (x^2 + 3x)^{1/2}$

$$y' = \frac{1}{2}(x^2 + 3x)^{-1/2} (2x + 3) = \frac{1}{2} \frac{1}{(x^2 + 3x)^{1/2}} (2x + 3) = \frac{2x + 3}{2(x^2 + 3x)^{1/2}}$$

proof of chain rule

(just for your own interest)

We are trying to find a pattern for

$$\frac{f(g(x+h)) - f(g(x))}{h}$$

Multiplying top and bottom by

$g(x+h) - g(x)$ gives

$$\left(\frac{f(g(x+h)) - f(g(x))}{h} \right) \left(\frac{g(x+h) - g(x)}{g(x+h) - g(x)} \right)$$

Rearranging gives

$$\left(\frac{f(g(x+h)) - f(g(x))}{g(x+h) - g(x)} \right) \left(\frac{g(x+h) - g(x)}{h} \right)$$

As $h \rightarrow 0$, we see the expression

above is approaching

$$f'(g(x))g'(x)$$

All Rules:

$$\frac{d}{dx}(f(x) + g(x)) = f'(x) + g'(x).$$

$$\frac{d}{dx}(cf(x)) = cf'(x).$$

$$\frac{d}{dx}(x^n) = nx^{n-1}.$$

$$\frac{d}{dx}(f(x)g(x)) = f(x)g'(x) + f'(x)g(x).$$

$$\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}$$

$$\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$$