

MATH 112  
Final Exam  
March 10, 2007

Name SOLUTION OUTLINES

Student ID # \_\_\_\_\_

Section \_\_\_\_\_

HONOR STATEMENT

"I affirm that my work upholds the highest standards of honesty and academic integrity at the University of Washington, and that I have neither given nor received any unauthorized assistance on this exam."

SIGNATURE: \_\_\_\_\_

1	14		6	10	
2	12		7	10	
3	10		8	12	
4	10		9	10	
5	12		Total	100	

- Your exam should consist of 9 problems. Check that you have a complete exam.
- Turn your cell phone OFF and put it away for the duration of the exam.
- Unless otherwise indicated, you must show your work. The correct answer with no supporting work may result in no credit.
- Unless otherwise indicated, you may round your FINAL ANSWER to two digits after the decimal.
- You may use a calculator for arithmetic purposes only (such as plugging into the quadratic formula or plugging into a function). ALL other work must be written and demonstrated on your exam. No credit will be given for guess-and-check or calculator methods, even if they give the correct answer.
- There are multiple versions of the exam. Any student found engaging in academic misconduct will receive a score of 0 on this exam.

GOOD LUCK!

1. (14 points) The total revenue and total cost for selling Frumtops are:

$$TR(q) = -5q^2 + 80q \text{ and } TC(q) = q^3 - 12q^2 + 60q + 60,$$

where  $q$  is in thousands of Frumtops and  $TR$  and  $TC$  are both in thousands of dollars.

(a) Use the derivative rules to find formulas for marginal revenue and marginal cost.

DERIVATIVES

ANSWER:  $MR(q) =$  \_\_\_\_\_  
 $MC(q) =$  \_\_\_\_\_

(b) What does it cost to produce the 6,001<sup>st</sup> Frumtop?

ASKING ABOUT MC AT 6000  
WHICH IS  $q = 6$  thousand Frumtops

COMPUTE  $MC(6)$

ANSWER: \_\_\_\_\_ dollars

(c) What quantity maximizes profit?

SOLVE  $MR(q) = MC(q)$

ANSWER:  $q =$  \_\_\_\_\_ thousand Frumtops

(d) Find all quantities at which the slope of the tangent line to the marginal revenue graph is the same as the slope of the tangent line to the marginal cost graph.

SAME AS  $MC'(x)$

SAME AS  $MR'(x)$

FIND  $MR'(x)$  AND  $MC'(x)$   
SOLVE  $MR'(x) = MC'(x)$

ANSWER:  $q =$  \_\_\_\_\_ thousand Frumtops

(e) Is the graph of total revenue concave up, concave down, or neither at  $q = 5$ ? (As always, you must show your work.)

NEED TO FIND  $TR''(x)$   
THEN COMPUTE  $TR''(5)$  IS IT POSITIVE OR NEGATIVE

ANSWER: (circle one) concave up concave down neither

2. (12 points) Let  $y = f(x) = x^2 - 5x + 2\ln(x) + 6$

(a) Find all the critical numbers of  $f(x)$ .

FIND  $f'(x)$   
SOLVE  $f'(x) = 0$

NOTE:  $f'(x) = 2x - 5 + 2\frac{1}{x} \stackrel{?}{=} 0$   
SO MULTIPLY BOTH SIDES BY  $x$   
TO GET  $2x^2 - 5x + 2 = 0$

ANSWER:  $x = \underline{\hspace{2cm}}$

(b) Use the Second Derivative Test to determine whether each critical number you found in part (a) gives a local minimum or a local maximum, or state that the test is inconclusive.

FIND  $f''(x)$

PLUG EACH CRITICAL NUMBER FROM (a)  
INTO  $f''(x)$ .

2<sup>nd</sup> Deriv. TEST → IF  $f''$  IS POSITIVE, THEN THE CRITICAL PT GIVES A LOCAL MIN.  
IF  $f''$  IS NEGATIVE, THEN THE CRITICAL PT GIVES A LOCAL MAX.

(c) Find the the global maximum and global minimum values of  $f(x)$  on the interval from  $x = 1$  to  $x = 10$ .

COMPUTE  $f(1) = ?$ ,  $f(10) = ?$

AND COMPUTE THE VALUE OF  $f(x)$  AT  
EACH CRITICAL NUMBER BETWEEN 1 AND 10 FROM  
PART (a). BIGGEST OUTPUT = GLOBAL MAX  
SMALLEST OUTPUT = GLOBAL MIN

ANSWERS: GLOBAL MINIMUM:  $\underline{\hspace{2cm}}$

GLOBAL MAXIMUM:  $\underline{\hspace{2cm}}$

3. (10 points) This week it is your job to buy muffins and bagels for a meeting at your office. Each muffin will feed only 1 person, while each bagel will be cut in half to feed 2 people.

A muffin costs \$1.50 and a bagel costs \$2.00. Your boss gives you \$30.00 to spend. When you get to the bakery they only have 12 bagels left.

Let  $x$  be the number of muffins that you buy and  $y$  be the number of bagels you buy. You want to maximize the number of people that you can feed.

- (a) Give the constraints and the objective function for this problem.

ANSWER: objective function: \_\_\_\_\_

constraints: \_\_\_\_\_

- (b) Sketch the feasible region, clearly label all vertices with their exact values.


SKIP (THIS IS A MATH III PROBLEM)  
WE WON'T ASK THIS ON OUR MATH 112 FINAL.

- (c) How many muffins,  $x$ , and bagels,  $y$ , should you buy to maximize the number of people you can feed? Justify your answer.

ANSWER:  $x =$  \_\_\_\_\_ muffins

$y =$  \_\_\_\_\_ bagels

4. (10 points) The graph below is of the function  $y = f(x)$ .



(a) Give two values of  $x$  at which  $f'(x) = 2$ .  
 WANT TO FIND WHEN THE SLOPE OF THE TANGENT LINE IS 2, DRAW A LINE OF SLOPE 2. THEN FIND A TANGENT LINE WITH THAT SLOPE

ANSWER:  $x =$  \_\_\_\_\_ and  $x =$  \_\_\_\_\_

(b) Approximate the value of  $\frac{f(4+h) - f(4)}{h}$  if  $h = 0.001$ .

$\frac{f(4.001) - f(4)}{0.001} \approx$  slope of the tangent at 4. DRAW IT. FIND THE SLOPE

ANSWER:  $\frac{f(4+h) - f(4)}{h} =$  \_\_\_\_\_

(c) Give an interval of length 1 over which the graph of  $f'(x)$  is positive and decreasing.

$f'(x)$  positive is the same as  $f(x)$  increasing (0.5 TO 3.5 and 6.5 TO 10)  
 $f'(x)$  decreasing is the same as  $f''(x)$  negative so  $f(x)$  concave down (2 TO 5 AND 8 TO 10)

ANSWER: from  $x =$  \_\_\_\_\_ to  $x =$  \_\_\_\_\_

(d) Give an interval over which the graph of  $f'(x)$  is shaped like this:

$f'(x) = 0 \Leftrightarrow f(x)$  horizontal tangent  
 $f'(x)$  negative  $\Leftrightarrow f(x)$  decreasing  
 WE NEED HORIZ. TANGENT, DEC., HORIZ. TANGENT

GIVE ANY ONE MINUTE INTERVAL WHERE BOTH ARE HAPPENING

ANSWER: from  $x =$  \_\_\_\_\_ to  $x =$  \_\_\_\_\_

5. (12 points) Two balloons are next to each other at time  $t = 0$ . You are given the following information about them:

Altitude vs. time:  $A(t) = \frac{1}{3}t^3 - \frac{7}{2}t^2 + 10t + 50$   
 for balloon A  
 Rate of ascent:  $A'(t) =$   
 for balloon A  
 Altitude vs. time  $B(t) =$   
 for balloon B:  
 Rate of ascent  $B'(t) = 20 - 10t$   
 for balloon B:

DERIVATIVE

ANTIDERIVATIVE

NOTE THAT  
 $B(0) = A(0) = 50$   
 SO YOU CAN FIND  
 C

where  $t$  is in minutes and Altitude is in feet.

(a) Find the highest altitude Balloon A reaches in the first 3 minutes.

GLOBAL MAX OF A FROM 0 TO 3  
 COMPUTE  $A(0) = ?$  AND  $A(3) = ?$   
 FIND CRITICAL NUMBERS  $A'(t) = 0$  ← SOLVE  
 IF ANY CRITICAL # IS BETWEEN 0 AND 3  
 THEN COMPUTE THE VALUE OF  $A(t)$  AT THAT VALUE  
 LARGEST OUTPUT

ANSWER: highest altitude = \_\_\_\_\_ feet

(b) Use the fact that  $B(0) = A(0)$  to determine the formula for  $B(t)$  without any undetermined constants.

$B(t) = \int 20 - 10t \, dt = 20t - 5t^2 + C$   
 USE THE FACT THAT  $B(0) = 50$  TO FIND C

ANSWER:  $B(t) =$  \_\_\_\_\_


(c) Find all times,  $t$ , at which balloon B is ascending 3 ft per minute faster than balloon A.

SOLVE  $B'(t) = A'(t) + 3$

ANSWER:  $t =$  \_\_\_\_\_ minutes

6. (10 points)

The graph below is of the function  $y = f(t)$ .



Using the graph above, we define a new function

$$A(m) = \int_0^m f(t) dt =$$

SIGNED AREA UNDER  $f(t)$  FROM 0 TO  $m$

(a) Find the value of  $t$  between 0 and 17 at which  $f(t)$  has a global maximum.

HEIGHT ON GRAPH  
JUST READ OFF LOCATION CORRESPONDING TO HIGHEST POINT

ANSWER:  $t =$  \_\_\_\_\_

(b) Find all values of  $m$  between 0 and 17 at which  $A(m)$  has a local minimum.

$A(m)$  LOCAL MIN  
 $\Rightarrow A'(x) = f(x) = 0$   
AND  $A'(x) = f(x)$  CHANGED FROM POSITIVE TO NEGATIVE

AREA

ANSWER:  $m =$  \_\_\_\_\_

(c) Determine the value of  $f'(4)$ .

SLOPE ON  $f(t)$   
FIND TWO POINTS  
COMPUTE SLOPE

ANSWER:  $f'(4) =$  \_\_\_\_\_

(d) Determine the value of  $A'(4)$ .

SAME AS  $f(4) =$  HEIGHT


ANSWER:  $A'(4) =$  \_\_\_\_\_

(e) Compute  $\int_6^{10} f(t) dt$ .

AREA FROM 6 TO 10

ANSWER:  $\int_6^{10} f(t) dt =$  \_\_\_\_\_

7. (10 points) The graphs below are of marginal revenue ( $MR$ ) and marginal cost ( $MC$ ) (both in dollars) for producing  $q$  thousand Chubkins.



- (a) Determine the increase in total revenue, if  $q$  increases from 12 to 20 thousand Chubkins.

AREA UNDER MR FROM 12 TO 20

ANSWER: \_\_\_\_\_ thousand dollars

- (b) If  $q$  changes from 6 to 8 thousand Chubkins, does profit increase or decrease? By how much?

MR > MC SO PROFIT INCREASES!  
AREA BETWEEN MR AND MC FROM 6 TO 8 IS HOW MUCH PROFIT GOES UP

ANSWER: Profit (circle one) increases decreases by \_\_\_\_\_ thousand dollars.

- (c) What is the largest value of total revenue?

OCCURS WHEN  $TR'(x) = MR(x) = 0$  SO AT  $x = 20$ .  
FIND AREA UNDER MR FROM 0 TO 20 TO FIND  $TR(20)$

ANSWER: \_\_\_\_\_ thousand dollars

- (d) If fixed costs are \$11.2 thousand dollars, what is the profit earned for selling 2 thousand Chubkins?

$$P(2) = TR(2) - TC(2)$$

$$= TR(2) - VC(2) - FC$$

$$= (\text{AREA BETWEEN MR AND MC FROM 0 TO 2}) - FC$$

ANSWER: \_\_\_\_\_ thousand dollars

↑ COMPUTE



8. (12 points)

(a) Evaluate:  $\int_1^2 x^3 + 4x - 1 \, dx$ 

$$= \left. \frac{1}{4} x^4 + 2x^2 - x \right|_1^2$$

$$= \left( \frac{1}{4} (2)^4 + 2(2)^2 - (2) \right) - \left( \frac{1}{4} (1)^4 + 2(1)^2 - (1) \right)$$

$$= ? \quad \leftarrow \text{FINISH}$$

(b) Evaluate:  $\int 3x + \frac{4}{x} - \sqrt{x} \, dx$ 

$$= \int 3x + 4 \cdot \frac{1}{x} - x^{1/2} \, dx$$

$$= \frac{3}{2} x^2 + 4 \ln(x) - \frac{2}{3} x^{3/2} + C$$

FINAL ANSWER

(c) Let  $z = \frac{t^2 + e^{5t+1}}{t^3 - 4}$ . Find  $\frac{dz}{dt}$ . Do not simplify.

QUOTIENT RULE!

DON'T FORGET

$$\frac{dz}{dt} = \frac{(t^3 - 4)(2t + e^{5t+1} \cdot 5) - (t^2 + e^{5t+1})(3t^2)}{(t^3 - 4)^2}$$

FINAL ANSWER

(d) Let  $f(x, y) = 3x^3y + x^2 \ln(x) + y^2$ . Find  $f_x(x, y)$ .

↑ ↑  
BOTH X'S, PRODUCT RULE NEEDED!

$$f_x(x, y) = 9x^2y + x^2 \cdot \frac{1}{x} + 2x \ln(x)$$

$$= 9x^2y + x + 2x \ln(x)$$

FINAL ANSWER

9. (10 points) Let  $z = f(x, y) = -3x^2 + 2y^2 + 36x + 12xy$ .

(a) Write out the formulas for  $f_x(x, y)$  and  $f_y(x, y)$ .

$$f_x(x, y) = \frac{-6x + 36 + 12y}{1}$$

$$f_y(x, y) = \frac{4y + 12x}{1}$$

(b) Use partial derivatives to tell which of the following numbers is bigger:

$$A = \frac{f(1.0001, 1) - f(1, 1)}{0.0001} \quad \text{or} \quad B = \frac{f(1, 1.0001) - f(1, 1)}{0.0001}$$

Justify your answer. (That is, you must show work to get credit.)

$$A \approx f_x(1, 1) \quad (x \text{ IS CHANGING!})$$

$$B \approx f_y(1, 1) \quad (y \text{ IS CHANGING!})$$

COMPUTE! WHICH IS BIGGER

ANSWER: (circle one)

A

B

(c) Find all points  $(x, y)$  which are candidates for local maxima or local minima of  $f(x, y)$  or show that there are no such candidates.

SOLVE SYSTEM

$$\begin{aligned} \textcircled{i} \quad -6x + 36 + 12y &= 0 \\ \textcircled{ii} \quad 4y + 12x &= 0 \Rightarrow 4y = -12x \\ \text{From } \textcircled{ii} \quad y &= -3x \end{aligned}$$

SUBSTITUTING  $\textcircled{ii}$  INTO  $\textcircled{i}$  GIVES

$$-6x + 36 + 12(-3x) = 0$$

$$-6x + 36 - 36x = 0$$

$$36 = 42x$$

$$x = \frac{36}{42} = \frac{6}{7}$$

$$\text{AND FROM } \textcircled{ii} \quad y = -3\left(\frac{6}{7}\right) = -\frac{18}{7}$$

ANSWER: (Check one)  The candidate(s) is/are:  $(x, y) =$

There are no such candidates.

$$\left(\frac{6}{7}, -\frac{18}{7}\right)$$

CHECK!