

Closing Tue: Taylor Notes 1, 2
Closing Thu: Taylor Notes 3
Final: Sat, June 2, 5:00pm, KANE 130
Eight pages, covers everything.

TN 2 & 3: Higher order approx.

Recall: *1st Taylor polynomial*

$$T_1(x) = f(b) + f'(b)(x - b)$$

Error Bound

On interval $[a,b]$, if $|f''(x)| \leq M$,
then $|f(x) - T_1(x)| \leq \frac{M}{2}|x - b|^2$.

Entry Task: Let $f(x) = x^{1/3}$.

- (a) Find the 1st Taylor Polynomial based at $b = 8$.
- (b) Give a bound on the error over the interval $[7,9]$.

2nd Taylor Polynomial is given by

$$T_2(x) = f(b) + f'(b)(x - b) + \frac{1}{2}f''(b)(x - b)^2$$

Quadratic error bound theorem

On interval $[a,b]$, if $|f'''(x)| \leq M$,
then $|f(x) - T_2(x)| \leq \frac{M}{6} |x - b|^3$.

Example:

Find the 2nd Taylor polynomial for
 $f(x) = x^{1/3}$ based at $b = 8$ and find
an error bound on the interval $[7,9]$.

Taylor Approximation Idea:

If two functions have **all** the same derivative values, then they are the same function (up to a constant).

Let's compare derivatives of $f(x)$ and $T_2(x)$ at b .

$$T_2(x) = f(b) + f'(b)(x - b) + \frac{1}{2}f''(b)(x - b)^2$$

$$T_2'(x) = 0 + f'(b) + f''(b)(x - b)$$

$$T_2''(x) = 0 + 0 + f''(b)$$

$$T_2'''(x) = 0$$

Now plug in $x = b$ to each of these.

- What do you see?
- Why did we need a $\frac{1}{2}$?
- What would $T_3(x)$ look like?
- What would $T_4(x)$ look like?
($T_5(x)$?, $T_6(x)$?...)

nth Taylor polynomial

$$f(b) + f'(b)(x - b) + \frac{1}{2}f''(b)(x - b)^2 + \frac{1}{3!}f'''(b)(x - b)^3 + \dots + \frac{1}{n!}f^{(n)}(b)(x - b)^n$$

In sigma notation:

$$T_n(x) = \sum_{k=0}^n \frac{1}{k!} f^{(k)}(b)(x - b)^k$$

Example: Find the 9th Taylor polynomial for $f(x) = e^x$ based at $b = 0$, and give an error bound on the interval $[-2, 2]$.


Taylor's Inequality (error bound):

on a given interval $[a, b]$,

if $|f^{(n+1)}(x)| \leq M$, then

$$|f(x) - T_n(x)| \leq \frac{M}{(n+1)!} |x - b|^{n+1}$$

$f(x) = e^x$ and
 $T_1(x), T_2(x), T_3(x), T_4(x), T_5(x)$



Example: Again consider,

$$f(x) = e^x \text{ based at } b = 0$$

Find the first value of n when

Taylor's inequality gives an error less than 0.0001 on $[-2,2]$.

Side Note:

For a fixed constant, a , the expression $\frac{a^k}{k!}$ goes to zero as k goes to infinity.

So the expression $\frac{1}{(n+1)!} |x - b|^{n+1}$, will always go to zero as n gets bigger.

Which means that the error goes to zero, unless something unusual is happening with M , which will see in examples later.

TN 4: Taylor Series

Def'n: The **Taylor Series** for $f(x)$ based at b is defined by

$$\sum_{k=0}^{\infty} \frac{1}{k!} f^{(k)}(b)(x - b)^k = \lim_{n \rightarrow \infty} T_n(x)$$

If the limit exists at x ,
then we say it **converges** at x .
(*i.e.* the error goes to zero at x)

Otherwise, we say it **diverges** at x .

The **open interval of convergence**
gives the largest open interval over
which the series converges.

Note: If

$$\lim_{n \rightarrow \infty} \frac{M}{(n + 1)!} |x - b|^{n+1} = 0$$

then x is in the open interval of
convergence.