

RC-Graphs and Schubert Polynomials

Nantel Bergeron and Sara Billey

CONTENTS

- 1. Introduction
- 2. Background on Schubert Polynomials
- 3. Constructing Schubert Polynomials From RC-Graphs
- 4. Double Schubert Polynomials
- 5. Monk's Rule
- 6. Conjectures
- Acknowledgements
- References

Using a formula of Billey, Jockusch and Stanley, Fomin and Kirillov have introduced a new set of diagrams that encode the Schubert polynomials. We call these objects rc-graphs. We define and prove two variants of an algorithm for constructing the set of all rc-graphs for a given permutation. This construction makes many of the identities known for Schubert polynomials more apparent, and yields new ones. In particular, we give a new proof of Monk's rule using an insertion algorithm on rc-graphs. We conjecture two analogs of Pieri's rule for multiplying Schubert polynomials. We also extend the algorithm to generate the double Schubert polynomials.

1. INTRODUCTION

Schubert polynomials (defined in Section 2) are of interest in algebraic geometry and in combinatorics. Their history goes back to the nineteenth century, but the modern notion is due to Lascoux and Schützenberger [1982; 1985], who developed a beautiful theory of these polynomials. A complete exposition can be found in [Macdonald 1991].

Our approach to computing Schubert polynomials is an algorithmic one. The idea is related to a conjecture originally due to Axel Kohnert [1990], saying that the Schubert polynomials could be constructed by applying a recursive algorithm on the diagram $D(w)$ defined from a permutation $w = [w_1, \dots, w_n]$ as the set of pairs (i, w_j) for which $j > i$ and $w_j < w_i$. Each diagram that appears in the recursion contributes a term to the Schubert polynomial. At this time, Kohnert's conjecture has not been proved except in the special case that w is a vexillary permutation (or 2143-avoiding). We have verified the conjecture for every permutation in the symmetric group S_7 .

Bergeron was supported by the National Science Foundation.

Billey was supported by the National Physical Science Consortium, IBM and UCSD.

In [Bergeron 1992] another algorithm, similar to Kohnert's, is given. This algorithm again starts with $D(w)$ but is computationally more complex. Furthermore, some identities about Schubert polynomials cannot readily be established using this method of computation. Therefore, we were driven to find yet another constructive method of computing Schubert polynomials from some other set of diagrams.

Fomin and Kirillov [1993] introduced a new set of diagrams that encode the Schubert polynomials. We call this object an rc-graph (for reduced-word compatible sequence graph). In the spirit of Kohnert's conjecture, we are interested in constructing Schubert polynomials by doing "moves" on rc-graphs. We will define and prove two variants of an algorithm for constructing the set of all rc-graphs for a given permutation in Section 3. This algorithm has been much more efficient in time and space than previously known ones. It can also be extended to generate the double Schubert polynomials as well. Using this construction, many of the identities known for Schubert polynomials become more apparent and new identities have emerged. In particular, Section 5 gives a new proof of Monk's rule, using an insertion algorithm on rc-graphs.

Experimentation on computers has greatly facilitated our work. We have gained an invaluable amount of intuition about Schubert polynomials by looking at data, we have been able to rule out false conjectures quickly, and we have found two very interesting conjectures. In Section 6, we conjecture two analogs of Pieri's rule for multiplying Schubert polynomials. We explain how we used computers to find these conjectures.

2. BACKGROUND ON SCHUBERT POLYNOMIALS

We will briefly review the basic terminology from [Macdonald 1991]. Let S_n be the symmetric group on n elements; we write a permutation $w \in S_n$ in one-line notation as $[w_1, w_2, \dots, w_n]$, for w_1, \dots, w_n in $\{1, 2, \dots, n\}$. Let s_i denote the transposition that interchanges the i -th and $(i + 1)$ -st entries

when acting from the right on a permutation. It is well known that s_1, \dots, s_{n-1} generate S_n , with the relations

$$\begin{aligned} s_i^2 &= 1, \\ s_i s_j &= s_j s_i \quad \text{for } |i - j| > 1, \\ s_i s_{i+1} s_i &= s_{i+1} s_i s_{i+1}. \end{aligned}$$

For $w \in S_n$, the length $l(w)$ of w is the length of a shortest string $s_{a_1} s_{a_2} \dots s_{a_p}$ representing w ; we call the p -tuple $a_1 a_2 \dots a_p$ a reduced word for w . Let $R(w)$ denote the set of all reduced words for a permutation w .

Let $\mathbb{Z}[x_1, x_2, \dots, x_n]$ denote the ring of polynomials in n variables with coefficients in \mathbb{Z} . We define an action of $w \in S_n$ on $f \in \mathbb{Z}[x_1, \dots, x_n]$ by $wf(x_1, x_2, \dots, x_n) = f(x_{w_1}, x_{w_2}, \dots, x_{w_n})$. From this we can define the divided difference operators

$$\partial_i f(x_1, x_2, \dots, x_n) = \frac{f(x_1, \dots, x_n) - s_i f(x_1, \dots, x_n)}{x_i - x_{i+1}}$$

for $1 \leq i \leq n - 1$. It is easy to check that $\partial_i^2 = 0$, $\partial_i \partial_j = \partial_j \partial_i$ if $|i - j| > 1$ and $\partial_i \partial_{i+1} \partial_i = \partial_{i+1} \partial_i \partial_{i+1}$. Therefore $\partial_w = \partial_{a_1} \partial_{a_2} \dots \partial_{a_p}$ does not depend on the choice of reduced word $a_1 a_2 \dots a_p \in R(w)$. In addition, $\partial_{a_1} \partial_{a_2} \dots \partial_{a_p} = 0$ if $a_1 a_2 \dots a_p$ is not reduced.

Definition. For each permutation $w \in S_n$, the Schubert polynomial \mathfrak{S}_w is

$$\mathfrak{S}_w = \partial_{w^{-1}w_0} x_1^{n-1} x_2^{n-2} \dots x_{n-1}^1 x_n^0,$$

where $w_0 = [n, n - 1, \dots, 1]$ is the longest element of S_n .

For $m > n$, we can associate with a permutation $w = [w_1, w_2, \dots, w_n] \in S_n$ the permutation $v = [w_1, w_2, \dots, w_n, n + 1, \dots, m] \in S_m$. Now v and w have the same set of reduced words (the generators being interpreted in S_n or S_m as appropriate), so their length is also the same. From now on we identify two such permutations, making S_n into a subgroup of S_m for $n < m$. We set $S_\infty = \bigcup S_n$, and still write $w \in S_\infty$ as $w = [w_1, w_2, \dots, w_n]$ if $w \in S_n$. The definition of a Schubert polynomial applies equally well to $w \in S_\infty$; it is not obvious at

this point that the result is independent of n , but this will follow from Theorem 2.1.

The theory of Schubert polynomials is closely intertwined with the study of reduced words. The key to understanding this relationship is Theorem 2.1, which amounts to an alternative definition of Schubert polynomials. This result was originally conjectured by Richard Stanley, and first proved in [Billey et al. 1993] and subsequently in [Fomin and Stanley 1991] in a very elegant way.

Definition. If $\mathbf{a} = a_1 \dots a_p \in R(w)$, we say that a p -tuple $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_p)$ of (strictly) positive integers is *\mathbf{a} -compatible* if

$$\begin{aligned} \alpha_1 &\leq \alpha_2 \leq \dots \leq \alpha_p, \\ \alpha_j &\leq a_j \quad \text{for } 1 \leq j \leq p, \\ \alpha_j &< \alpha_{j+1} \quad \text{if } a_j < a_{j+1}. \end{aligned}$$

Let $C(\mathbf{a})$ denote the set of all \mathbf{a} -compatible sequences.

Theorem 2.1 [Billey et al. 1993; Fomin and Stanley 1991]. *For any permutation $w \in S_\infty$,*

$$\mathfrak{S}_w = \sum_{\mathbf{a} \in R(w)} \sum_{\boldsymbol{\alpha} \in C(\mathbf{a})} x_{\alpha_1} x_{\alpha_2} \dots x_{\alpha_p}.$$

It is easy to compute a Schubert polynomial using this theorem, provided one can compute all reduced words. For example, for $w = [3, 1, 2, 5, 4]$ the reduced words are 214, 241 and 421. For each of these we compute all compatible sequences as follows:

$$\begin{array}{ccc} \underline{214} & \underline{241} & \underline{421} \\ 112 & & 111 \\ 113 & & \\ 114 & & \end{array}$$

Therefore $\mathfrak{S}_{[3,1,2,5,4]} = x_1^3 + x_1^2 x_2 + x_1^2 x_3 + x_1^2 x_4$.

3. CONSTRUCTING SCHUBERT POLYNOMIALS FROM RC-GRAPHS

In this section we define rc-graphs and an algorithm for computing Schubert polynomials. The goal of our algorithm is to start with a particular

rc-graph and apply a sequence of transformations, thereby obtaining all rc-graphs for a permutation. After proving several lemmas, we state and prove Theorem 3.7, our main theorem, which states that this algorithm constructs the Schubert polynomials.

There are in fact two flavors of the algorithm, dual in a certain sense to one another. The duality follows from Lemma 3.2, which gives a correspondence between the rc-graphs of w and those of w^{-1} .

We conclude this section with some corollaries that follow easily from Theorems 3.7.

Definition. Given a reduced word $\mathbf{a} = a_1 a_2 \dots a_p$ and an \mathbf{a} -compatible sequence $\boldsymbol{\alpha} = \alpha_1 \alpha_2 \dots \alpha_p$, the *reduced-word compatible sequence graph* or *rc-graph* $(\mathbf{a}, \boldsymbol{\alpha})$ is the subset $D(\mathbf{a}, \boldsymbol{\alpha})$ of $\{1, 2, \dots\} \times \{1, 2, \dots\}$ consisting of pairs

$$(\alpha_k, a_k - \alpha_k + 1) \quad \text{for } 1 \leq k \leq p.$$

We let $\mathcal{RC}(w) = \{D(\mathbf{a}, \boldsymbol{\alpha}) : \mathbf{a} \in R(w), \boldsymbol{\alpha} \in C(\mathbf{a})\}$.

We will depict $D(\mathbf{a}, \boldsymbol{\alpha})$ as an incidence table in upper triangular form, with rows and columns labeled $1, 2, \dots$. For example, one reduced word for $[3, 1, 4, 6, 5, 2]$ is $\mathbf{a} = 521345$, and $\boldsymbol{\alpha} = 111235$ is \mathbf{a} -compatible. The rc-graph $D(\mathbf{a}, \boldsymbol{\alpha})$ is given by

$$\begin{array}{cccccc} & 1 & 2 & 3 & 4 & 5 & 6 \\ 1 & + & + & \cdot & \cdot & + & \cdot \\ 2 & \cdot & + & \cdot & \cdot & \cdot & \\ 3 & \cdot & + & \cdot & \cdot & & \\ 4 & \cdot & \cdot & \cdot & & & \\ 5 & + & \cdot & & & & \\ 6 & \cdot & & & & & \end{array} \tag{3.1}$$

where a + represents a pair that belongs to the graph and a · one that does not.

Remarks 3.1. It will soon become clear why it is more convenient in our context to use the pairs $(\alpha_k, a_k - \alpha_k + 1)$ instead of the pairs (α_k, a_k) .

One can also define rc-graphs in the general case when \mathbf{a} is not a reduced word, but we will not need that.


Given an rc-graph $D(\mathbf{a}, \alpha)$, one can recover \mathbf{a} by reading out the numbers $j + i - 1$ for the occupied positions (i, j) from *right to left* in each row from top to bottom. The compatible sequence α is recovered by reading the row numbers of the occupied positions, in the same order. If $a_1 a_2 \dots a_p$ is the reduced word read from the rc-graph D , we let

$$\text{perm } D = s_{a_1} s_{a_2} \dots s_{a_p}$$

be the permutation such that $D \in \mathcal{RC}(\text{perm } D)$.

It follows from the definition of a compatible sequence that, if $\text{perm } D \in S_n$, all the elements (i, j) of D satisfy $i + j < n$. Conversely, any set of pairs of positive integers (i, j) that lies in the area $i + j < n$ and that gives a reduced word \mathbf{a} by means of the above reading is an rc-graph of a permutation of S_n , and the corresponding sequence of row numbers is \mathbf{a} -compatible.

The rc-graphs originally introduced by Fomin and Kirillov [1993] are a bit more elaborate. They represent a *planar history* of the inversions of $w = \text{perm } D$. To get from the definition above to this other one, we draw strands that cross at the positions $(i, j) \in D$ and avoid each other at the positions $(i, j) \notin D$. Here is the diagram for the rc-graph in (3.1):



It is easy to see that, for $D \in \mathcal{RC}(w)$, the strand that starts at the top of the w_i -th column wends its way down to the i -th row on the left side of the diagram. No two strands can cross more than once, since by removing two such crossings we would obtain a shorter representation for the permutation w , contradicting the fact that the given word is reduced. Thus $\mathcal{RC}(w)$ is the set of all such strand configurations with exactly $l(w)$ crossings. From

now on we will often think of our rc-graphs as strand diagrams, but we will draw the strands only when needed. Moreover we will not draw the “sea” of wavy strands past the crossings with $i + j$ maximal.

Lemma 3.2. *The transpose D^t of an rc-graph D in $\mathcal{RC}(w)$ is an rc-graph in $\mathcal{RC}(w^{-1})$.*

Proof. If $D \in \mathcal{RC}(w)$, the strands in D^t trace out the permutation w^{-1} . Furthermore, $l(w) = l(w^{-1})$. Therefore, the number of crossings of D^t is minimal. \square

We also write $\rho(D)$ for D^t , thus defining a bijection $\rho : \mathcal{RC}(w) \rightarrow \mathcal{RC}(w^{-1})$. Vic Reiner has suggested the same map using only reduced words and compatible sequences.

If we use the notation

$$x_D = \prod_{(i,j) \in D} x_i,$$


the following corollary is a simple consequence of Theorem 2.1. This was also noted in [Fomin and Kirillov 1993].

Corollary 3.3. *For any permutation $w \in S_\infty$,*

$$\mathfrak{S}_w = \sum_{D(\mathbf{a}, \alpha) \in \mathcal{RC}(w)} x_{D(\mathbf{a}, \alpha)}. \quad \square$$

We now introduce *moves* on rc-graphs that are analogous to the moves in Kohnert’s conjecture [Kohnert 1990].


For $w \in S_\infty$ and $D \in \mathcal{RC}(w)$, a *ladder move* L_{ij} is a change of the following type:



Here the two columns are adjacent and the number of rows involved is arbitrary. Formally, $L_{ij}(D) = D \cup \{(i - m, j + 1)\} \setminus \{(i, j)\}$, where the following conditions must be satisfied:

- $(i, j) \in D, (i, j + 1) \notin D,$
- $(i - m, j), (i - m, j + 1) \notin D$ for some $0 < m < i,$
and
- $(i - k, j), (i - k, j + 1) \in D$ for each $1 \leq k < m.$

A chute move C_{ij} is a change of the following type (we leave the formal description to the reader):



It is evident from these definitions and from Lemma 3.2 that:


Lemma 3.4. *Transposition conjugates chute moves to ladder moves, and vice versa. In symbols,*

$$\rho(L_{ij}(D)) = C_{ji}(\rho(D)),$$

for any rc-graph $D.$

Lemma 3.5. *Ladder and chute moves preserve the permutation associated with an rc-graph. In symbols, $\text{perm } C_{ij}(D) = \text{perm } D$ if D is an rc-graph to which C_{ij} can be applied, and likewise for $L_{ij}.$*

Proof. The strands in the region of a chute move look like this:



Clearly the move does not affect the way in which the strings are permuted. Transposition proves the case of a ladder move. \square


Next we give a criterion for an rc-graph to be the result of a chute move. Note that if the same chute

move is applied to two different rc-graphs, the results are also different; therefore we can talk about inverse chute (and ladder) moves.


Lemma 3.6. *An rc-graph $D \in \mathcal{RC}(w)$ is the result of a chute move—or, equivalently, it admits an inverse chute move—if and only if there is some pair $(i, j) \notin D$ such that $(i + 1, j) \in D.$*

Geometrically, the criterion for there *not* being an inverse chute move is that all the pluses in each column are clumped together at the top.

Proof. Suppose there is $(i, j) \notin D$ with $(i + 1, j) \in D.$ Look right along row $i + 1$ for the smallest $k > j$ such that $(i + 1, k) \notin D.$ There must be such a $k,$ since D contains only a finite number of points. The situation then is the following:



where the positions on the top row strictly between j and k may be filled with $+$ or $\cdot,$ but the position (i, k) must have a dot; if that position had a $+$ we would have



which is impossible because, as already remarked, two strands cannot cross twice. Now look to the left of (i, k) for the largest $m < k$ such that $(i, m) \notin D.$ Then $(i + 1, m)$ is able to move to (i, k) by an inverse chute move.

The converse also follows from this analysis. \square

To state our main theorem, we introduce two rc-graphs for $w \in S_\infty$ that are extremal in an appropriate sense:

$$D_{\text{bot}}(w) = \{(i, c) : c \leq m_i\},$$

where $m_i = \#\{j : j > i \text{ and } w_j < w_i\},$ arises from the largest reduced word in reverse lexicographic

order and the largest compatible sequence for this word in ordinary lexicographic order; and

$$D_{\text{top}}(w) = \{(c, j) : c \leq n_j\},$$

where $n_j = \#\{i : i < w_j^{-1} \text{ and } w_i > j\}$, arises from the smallest reduced word in reverse lexicographic order and the smallest compatible sequence for this word in ordinary lexicographic order. We have $D_{\text{top}}(w) = D_{\text{bot}}^t(w^{-1})$. Here are D_{top} and D_{bot} for the permutation of (3.1) on page 259:

	1	2	3	4	5		1	2	3	4	5
1	+	+	·	·	+	1	+	+	·	·	·
2	·	+	·	·		2	·	·	·	·	
3	·	+	·			3	+	·	·		
4	·	+				4	+	+			
5	·					5	+				

Let $\mathcal{C}(D)$ be the set of rc-graphs that can be derived from D by some sequence of (forward) chute moves. Define $\mathcal{L}(D)$ analogously, with “ladder” replacing “chute”.

Theorem 3.7. *Let $w \in S_\infty$.*

- (a) $D_{\text{top}}(w)$ does not admit an inverse chute.
- (b) Any element of $\mathcal{RC}(w)$ other than $D_{\text{top}}(w)$ admits an inverse chute.
- (c) $\mathcal{C}(D_{\text{top}}(w)) = \mathcal{RC}(w) = \mathcal{L}(D_{\text{bot}}(w))$.

(d)
$$\mathfrak{S}_w = \sum_{D \in \mathcal{C}(D_{\text{top}}(w))} x_D = \sum_{D \in \mathcal{L}(D_{\text{bot}}(w))} x_D.$$

Proof. (a) follows from Lemma 3.6, since by definition every column of $D_{\text{top}}(w)$ has an initial run of pluses and no others.

(b) Suppose that $w \in S_n$. Consider, for all $w' \in S_n$, all rc-graphs $D \in \mathcal{RC}(w')$ that don't admit inverse chutes. For such a graph, column j has between 0 and $n - j$ pluses clumped at the top, again by Lemma 3.6. This gives n choices for the first column, $n - 1$ for the second, and so on, so there are at most $n!$ such graphs. By part (a), there is at least one such graph, $D_{\text{top}}(w')$, for each permutation. Therefore all graphs of this type are of the form $D_{\text{top}}(w')$.

(c) Anything obtained from D_{top} by chute moves is in $\mathcal{RC}(w)$, by Lemma 3.5. Now an inverse chute move pushes up a +, so any graph $D \in \mathcal{RC}(w)$ must turn after finitely many inverse chutes into one that does not admit an inverse chute—that is, into $D_{\text{top}}(w)$, according to part (b). Reversing this sequence and applying chutes to $D_{\text{top}}(w)$ we recover D . This proves the first equality. The second follows from the first by duality: more precisely,

$$\begin{aligned} \mathcal{RC}(w) &= \rho(\mathcal{RC}(w^{-1})) = \rho(\mathcal{C}(D_{\text{top}}(w^{-1}))) \\ &= \rho(\mathcal{C}(\rho(D_{\text{bot}}(w)))) = \mathcal{L}(D_{\text{bot}}(w)), \end{aligned}$$

by Lemmas 3.2 and 3.4.

(d) follows from (c) and Corollary 3.3. □

As an example, Figure 1 shows the computation of $\mathfrak{S}_{[1432]}$ using ladder moves.

Chute and ladder moves define two partial orders on $\mathcal{RC}(w)$, with relations $L_{ij}(D) \prec D$ and $C_{ij}(D) \prec D$, respectively. These partial orders are dual to one another.

The following result restricts the relations among the elements in the partial ordered set. However, there are still multiple paths to some of the rc-graphs.

Corollary 3.8. *We can generate $\mathcal{RC}(w)$ using only chute moves C_{ij} such that i is the largest in column j , that is, those satisfying $(k, j) \notin D$ for all $k > j$. Similarly, we can generate $\mathcal{RC}(w)$ using only rightmost ladder moves L_{ij} , that is, those satisfying $(i, k) \notin D$ for all $k > i$.*

Proof. Given $D \in \mathcal{RC}(w)$ different from $D_{\text{top}}(w)$, choose an inverse chute move C_{ij}^{-1} such that i is as large as possible. The point (i, j) must be the lowest point in column j of $C_{ij}^{-1}(D)$, otherwise there exists a point $(k, j) \notin D$ with $(k + 1, j) \in D$, and hence another possible inverse chute move C_{kj}^{-1} with $k > i$. Next choose the lowest inverse chute move possible on $C_{ij}^{-1}(D)$. Continue applying the lowest move until there are no inverse chute moves possible. Reversing this sequence gives a sequence of chute moves C_{ij} such that (i, j) is the lowest point in column i that transforms $D_{\text{top}}(w)$ to D .




FIGURE 1. Computation of the Schubert polynomial of $w = [1, 4, 3, 2]$ using Theorem 3.7. Starting from $D_{\text{bot}}(w)$ (leftmost diagram) we apply all possible ladder moves and compute the monomial for each graph so obtained. The result is $\mathfrak{S}_{[1432]} = x_2^2 x_3 + x_1 x_2^2 + x_1 x_2 x_3 + x_1^2 x_3 + x_1^2 x_2$.

Transposing this proof, we get the result for ladder moves. \square

Algebraic proofs of Corollaries 3.9, 3.10, and 3.12 appear in [Macdonald 1991]. Corollary 3.11 first appeared in [Billey et al. 1993].

Corollary 3.9. *The Schubert polynomials, indexed by permutations in S_∞ , form an integral basis for*

$$\mathbb{Z}[x_1, x_2, \dots].$$

Proof. The leading term of each \mathfrak{S}_w , in reverse lexicographic order, is given by $D_{\text{bot}}(w)$. Each D_{bot} is unique, so each leading term is unique. Furthermore, given any monomial $x_1^{\alpha_1} \dots x_m^{\alpha_m}$, there exists a permutation w such that $x_{D_{\text{bot}}(w)} = x_1^{\alpha_1} \dots x_m^{\alpha_m}$: simply put α_1 pluses at the beginning of row 1, α_2 on row 2, and so on. \square

Corollary 3.10. *Given permutations $u \in S_m$ and $v \in S_n$, let $u \times v = [u_1, \dots, u_m, v_1 + m, \dots, v_n + m]$ and $1_m \times v = [1, \dots, m, v_1 + m, \dots, v_n + m]$. Then*

$$\mathfrak{S}_u \mathfrak{S}_{1_m \times v} = \mathfrak{S}_{u \times v}. \tag{3.2}$$

Proof. Every rc-graph in $\mathcal{RC}(u)$ is contained in $\{(i, j) : i + j < m\}$, and no rc-graph in $\mathcal{RC}(1_m \times v)$ contains points in $\{(i, j) : i + j \leq m\}$. No rc-graph in $\mathcal{RC}(u \times v)$ contains a point on the line $i + j = m$. Therefore, there is a bijection between $\mathcal{RC}(u) \times \mathcal{RC}(1_m \times v)$ and $\mathcal{RC}(u \times v)$, given by sending (D_1, D_2) to $D_1 \cup D_2$. \square

Let \downarrow denote the operation inverse to $v \mapsto 1_1 \times v$, in the notation of the preceding lemma, so that $\downarrow v = [v_2, v_3, \dots, v_n]$ if $v \in S_n$ with $v_1 = 1$ (and $\downarrow v$ is undefined if $v_1 \neq 1$).

Corollary 3.11. *For $w \in S_\infty$, we have*

$$\mathfrak{S}_w(x_1, x_2, \dots) = \sum x_1^{l(v)} \mathfrak{S}_{\downarrow vw}(x_2, x_3, \dots),$$

where the sum is over all permutations $v \in S_\infty$ such that $l(w) = l(vw) + l(v)$, $v = s_{i_1} s_{i_2} \dots s_{i_p}$ with $i_1 < i_2 < \dots < i_p$, and $(vw)_1 = 1$.

Proof. There is a bijection

$$\mathcal{RC}(w) \leftrightarrow \bigcup (v, \mathcal{RC}(\downarrow vw)),$$

where the union is over all permutations $v \in S_n$ such that $l(w) = l(vw) + l(v)$, $v = s_{i_1} s_{i_2} \dots s_{i_p}$ with $i_1 < i_2 < \dots < i_p$, and $(vw)_1 = 1$. The bijection is given by sending $D \in \mathcal{RC}(w)$ to (v, D') if $v = s_{i_1} s_{i_2} \dots s_{i_p}$, where the first row of D are points in columns i_1, i_2, \dots, i_p , and D' is the rc-graph obtained by removing the first row of D . \square

The next result is a generalization of Corollary 3.11.

Corollary 3.12. *For any fixed positive integer m and any $w \in S_n$, we have the decomposition*

$$\begin{aligned} \mathfrak{S}_w(x_1, \dots, x_n) &= \sum d_{uv}^w \mathfrak{S}_u(x_1, \dots, x_m) \mathfrak{S}_v(x_{m+1}, \dots, x_n), \end{aligned}$$

where the d_{uv}^w are non-negative integers.

Proof. Given a polynomial $f(x_1, x_2, \dots, x_m)$, let $\rho_m f = f(x_1, \dots, x_m, 0, 0, \dots)$. By an abuse of notation, we also let $\rho_m(\mathcal{RC}(w)) = \{D \in \mathcal{RC}(w) : \rho_m(x_D) = x_D\}$. For each $w \in S_\infty$ and each m , there exists a bijection

$$\mathcal{RC}(w) \leftrightarrow \bigcup \rho_m(\mathcal{RC}(u)) \times \mathcal{RC}(v),$$

where the union is over all permutations u, v such that $l(u) + l(v) = l(w)$ and $1_m \times v = u^{-1}w$. The bijection takes D to

$$\{(i, j) \in D : j \leq m\} \times \{(i, j - m) : (i, j) \in D, j > m\}.$$

Therefore,

$$\begin{aligned} \mathfrak{S}_w(x_1, \dots, x_n) &= \sum_{\substack{l(u)+l(v)=l(w) \\ 1_m \times v = u^{-1}w}} (\rho_m \mathfrak{S}_u(x_1, \dots, x_n)) \mathfrak{S}_v(x_{m+1}, \dots, x_n). \end{aligned}$$

Finally, we can expand $\rho_m \mathfrak{S}_u$ in a positive sum of Schubert polynomials by the transition equation (4.16) of [Macdonald 1991]. \square

4. DOUBLE SCHUBERT POLYNOMIALS


Double Schubert polynomials generalize the usual Schubert polynomials in two alphabets. The original definition, given by Lascoux and Schützenberger, was written in terms of divided difference operators. Our definition follows from [Macdonald 1991, eq. (6.3)]. We show that the double Schubert polynomials can also be represented graphically and that these graphs can be constructed by means of ladder moves on an initial graph.

Definition. We take two alphabets $X = \{x_1, x_2, \dots\}$ and $Y = \{y_1, y_2, \dots\}$. For $w \in S_\infty$, we define the *double Schubert polynomial* $\mathfrak{S}_w(X, Y)$ as

$$\mathfrak{S}_w(X, Y) = \sum_{\substack{v^{-1}u=w \\ l(u)+l(v)=l(w)}} (-1)^{l(v)} \mathfrak{S}_u(X) \mathfrak{S}_v(Y).$$

Definition. A *double rc-graph* E for a permutation w consists of the following data: a pair (u, v) of permutations such that $v^{-1}u = w$ and $l(w) = l(u) + l(v)$, and an *rc-graph* for each of u and v .

We usually think of the *rc-graph* for v as being placed upside down above that of u , and label its rows with negative numbers. For this reason we call the *rc-graph* for v the *upper half* of E , and the *rc-graph* for u the *lower half*. For example, for $w = [4, 3, 2, 1]$, one possible double *rc-graph* is



depending on whether or not we draw the strands. Here $u = [3, 4, 1, 2]$ and $v = [2, 1, 4, 3]$, and the lower and upper halves are



No two strands of a double *rc-graph* can cross twice. Two crossings in the same half are forbidden by the analogous result for simple *rc-graphs*, while one crossing in the upper half and one in the lower half would violate the condition $l(w) = l(u) + l(v)$, as their removal would yield a shorter representation for w .

There is a natural bijection between the double *rc-graphs* of w and those of w^{-1} , given by interchanging the roles of u and v such that $w = v^{-1}u$, and the respective halves. Graphically, this corresponds to reflecting across $i = 0$.

We denote the set of all double *rc-graphs* for w by $\widetilde{\mathcal{RC}}(w)$. Given a double *rc-graph* E with upper half D_- and lower half D_+ , we define an associated monomial

$$(xy)_E = (-1)^{\text{card} D_-} x_{D_+} y_{D_-} = \prod_{(i,j) \in D_+} x_i \prod_{(i,j) \in D_-} -y_i.$$

Corollary 3.3 has the following counterpart:

$$\mathfrak{S}_w(X, Y) = \sum_{E \in \widetilde{\mathcal{RC}}(w)} (xy)_E. \tag{4.1}$$


Indeed, by the definition of double Schubert polynomials and Corollary 3.3 we have

$$\mathfrak{S}_w(X, Y) = \sum_{\substack{v^{-1}u=w \\ l(u)+l(v)=l(w)}} \sum_{C \in \mathcal{RC}(u)} \sum_{D \in \mathcal{RC}(v)} (-1)^{l(v)} x_C y_D,$$


and this latter sum clearly equals $\sum_{E \in \widetilde{\mathcal{RC}}(w)} (xy)_E$.

We can extend the definition of a ladder or chute move to double rc-graphs. We will study ladder moves only and leave chutes to the reader. A ladder move on a double rc-graph E composed of an upper half D_- and a lower half D_+ is of one of three types:

- (a) A (forward) ladder move in D_+ .
- (b) An inverse ladder move in D_- (note that since the upper half is drawn upside down, this type of ladder move still moves crossings up).
- (c) A move of the form



We denote by $\mathcal{L}(E)$ be the set of double rc-graphs that can be obtained from E by ladder moves. As in the case of simple rc-graphs, ladder moves preserve the permutation w ; this follows from Lemma 3.5 for moves of types (a) and (b), and from the following picture for moves of type (c):



Note, however, that the permutations v and u of the upper and lower halves are not preserved in this case.

Let $E_{\text{bot}}(w)$ be the double rc-graph for w whose lower half is $D_{\text{bot}}(w)$ and whose upper half is trivial (no crossings).

Theorem 4.1. *Let $w \in S_\infty$. Then*

$$\widetilde{\mathcal{RC}}(w) = \mathcal{L}(E_{\text{top}}(w))$$

and


$$\mathfrak{S}_w(X, Y) = \sum_{E \in \mathcal{L}(E_{\text{bot}}(w))} (xy)_E.$$

Stanley first noted that double Schubert polynomials could be expressed in terms of generalized compatible sequences as in Theorem 2.1.

Proof. The second equation follows from the first and from (4.1). To prove the first equation, we reason much as in the proof of Theorem 3.7. First, as already observed, any double rc-graph obtained from $E_{\text{bot}}(w)$ by ladder moves is in $\widetilde{\mathcal{RC}}(w)$.

To show, conversely, that any $E \in \mathcal{RC}(w)$ is in $\mathcal{L}(E_{\text{bot}}(w))$, we proceed by induction on the number of crossings in the upper half D_- of E . If D_- has no crossings, E is in $\mathcal{L}(E_{\text{bot}}(w))$, by Theorem 3.7(c) applied to the lower half D_+ of E . Now suppose D_- has at least one crossing. Again using Theorem 3.7 on D_+ , we can assume that $D_+ = D_{\text{bot}}(u)$ for some permutation u . Using upside down ladder moves on D_- we may also assume that the row $i = -1$ is not empty.

Therefore we can find $j > 0$ such that $(-1, j) \in E$ and $(-1, j + 1) \notin E$, then $i > 0$ such that $(i, j) \notin E$ and $(k, j) \in E$ for $0 < k < i$. Since $D_+ = D_{\text{bot}}(u)$, we must have $(i, j + 1) \notin E$. We claim that we are allowed to move $(-1, j)$ to (i, j) by an inverse ladder move of type (c). We need only check that $(k, j + 1) \in E$ for all $0 < k < i$. If we assume this is not the case, we would have the configuration shown on the right, for $k > 0$ minimal with $(k, j + 1) \notin E$. This is impossible, since no two



We call the sequence of positions

$$(i_0, j_0), (i_1, j'_1), (i_1, j_1), \dots, (i_p, j'_p), (i_p, j_p)$$

the *insertion path*. The coordinates of an insertion path must satisfy the following inequalities:

$$\begin{aligned} i &= i_0 > i_1 > \dots > i_p \geq 1, \\ j_0 &< j'_1 > j_1 < \dots < j'_p > j_p. \end{aligned}$$

Let t_{kl} be the transposition that interchanges the elements in positions k and l when acting from the right on a permutation. Then $l(wt_{kl}) = l(w) + 1$ when $k < l, w_k < w_l$, and for no index m between k and l is there a w_m such that $w_k < w_m < w_l$. Note that the output (D', k, l) of the insertion algorithm satisfies $\text{perm } D' = (\text{perm } D)t_{kl}$. Indeed, after each add/delete step, the resulting graph is still in $\mathcal{RC}(\text{perm } D)$, since the added and deleted crossings involve the same pair of strands. The final step, when (i_p, j_p) is added, involves the strands numbered $k = s_p$ and $l = t_p$, so it corresponds to multiplying the permutation on the right by t_{kl} . In addition, $l(\text{perm } D') = l(\text{perm } D) + 1$, since D' is an rc-graph by construction, and it has one more crossing than D .

Algorithm 5.2 (Reverse Insertion). Given a permutation w , integers $l > r \geq k > 0$ with $l(wt_{kl}) = l(w) + 1$, and an rc-graph $D' \in \mathcal{RC}(wt_{kl})$, we can reverse the insertion algorithm to find a pair (D, i) , where $D \in \mathcal{RC}(w)$, $i \leq r$, and $(D', k, l) = I_r(D, i)$. This is done as follows:

There is a unique position $(i_0, j_0) \in D'$ where the strands k and l cross. Delete that crossing from D' . Now if there is no position to the right of (i_0, j_0) where the configuration is

$$t \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} s \\ t \end{array} \quad \text{with } s \leq r < t, \quad (5.2)$$

stop. Otherwise, let (i_0, j'_0) be the position minimizing $j'_0 > j_0$ and having the configuration (5.2). Let $s_1 = s$ and $t_1 = t$ be the strands there. By the observation preceding Algorithm 5.1, these strands s_1 and t_1 must cross at some point (i_1, j_1) where $i_1 > i_0$ and $j_1 < j'_0$. Add (i_0, j'_0) from D' , and

delete (i_1, j_1) . If possible, find (i_1, j'_1) satisfying (5.2) and such that $j'_1 > j_1$ is minimal, and continue in this way. All strands below row r have labels larger than r and $i_0 < i_1 < \dots < i_p \leq r$, so this process will eventually end. The last row i_p from which we remove a point becomes i , and the modified rc-graph D' becomes D .

Theorem 5.3 (Monk's Rule). Given $w \in S_\infty$ and a simple transposition s_r , we have

$$\mathfrak{S}_{s_r} \mathfrak{S}_w = \sum_{\substack{k \leq r < l \\ l(wt_{kl}) = l(w) + 1}} \mathfrak{S}_{wt_{kl}}.$$

Proof. There is a monomial-preserving bijection

$$\mathcal{RC}(w) \times \mathcal{RC}(s_r) \rightarrow \bigcup \mathcal{RC}(w'),$$

where the union is over all permutations $w' = wt_{kl}$ such that $k \leq r < l$ and $l(wt_{kl}) = l(w) + 1$. The bijection is given by $(D, i) \mapsto I_r(D, i)$, where we are using the fact that any rc-graph for s_r contains a single crossing, on some row $i \leq r$, to identify $\mathcal{RC}(s_r)$ with $\{1, \dots, r\}$.

The discussion immediately before Algorithm 5.2 shows that, if $D \in \mathcal{RC}(w)$, then $I_r(D, i) \in \mathcal{RC}(w')$ for some $w' = wt_{kl}$ such that $k \leq r < l$ and $l(wt_{kl}) = l(w) + 1$.

Conversely, assume that we are given an rc-graph $D' \in \mathcal{RC}(w')$ where $w' = wt_{kl}$ and $k \leq r < l$ and $l(wt_{kl}) = l(w) + 1$. Each D'_i for $0 \leq i \leq p$ in the reverse insertion sequence has the strand configuration of w and the number of points is $l(w)$, hence is in $\mathcal{RC}(w)$. Furthermore, a unique $i \leq r$ is given by the final row i_p . \square

Remark 5.4. A permutation w is *Grassmannian* if it has only one descent in the sequence w_1, w_2, \dots . When w is Grassmannian, the insertion algorithm corresponds to the usual Schensted insertion on semistandard tableaux. We leave it to the reader to check that one can find a weight-preserving bijection between $\mathcal{RC}(w)$ and the set of semistandard tableaux of shape $\lambda(w)$ filled with numbers $1, 2, \dots, r$, where r is the unique descent of w . (See [Billey et al. 1993, Theorem 2.5] and [Bergeron

1992, §2.) Then one can show that our insertion algorithm, keeping r fixed, corresponds (under this bijection) to Schensted insertion.

6. CONJECTURES

As we know (Corollary 3.9), Schubert polynomials form an integral basis for $\mathbb{Z}[x_1, x_2, \dots]$. A long-standing open problems in the theory of Schubert polynomials is to find a combinatorial proof that the coefficients c_{uv}^w are positive in the expansion

$$\mathfrak{S}_u \mathfrak{S}_v = \sum c_{uv}^w \mathfrak{S}_w.$$

The c_{uv}^w are the analogs of the Littlewood–Richardson coefficients in the theory of Schur functions. Monk’s rule is a special case of this problem. The two rules conjectured in this Section will be analogs of Pieri’s Rule, which was the predecessor of the Littlewood–Richardson rule.

Our investigation of the insertion algorithm in Section 5 suggested a generalization of Monk’s rule. We successively insert $b \geq i_1 \geq i_2 \geq \dots \geq i_d$ into a diagram. This led us to Conjecture 6.1. We were not able to prove this conjecture by generalizing the proof of Monk’s rule, that is, by successive insertions, but we still believe it is true! (Here is a counterexample, found by computer, that shows that successive insertion alone will not prove the conjectures: the multiplication of $\mathfrak{S}_{[1,2,5,4,3]}$ with $\mathfrak{S}_{[1,2,4,5,3]}$. For the insertions of $i_1 = 3$ and $i_2 = 3$ in the rc-graph $D = \{(1, 3), (1, 4), (2, 3)\}$, the second insertion *path* fails to remain weakly to the right of the first one.)

The second conjecture was found by computer investigation. We assumed that there would be an analogue of Conjecture 6.1 for permutations with diagram $D(w)$ a single column. We programmed several different patterns until a rule emerged. This computation would have been impossible to do by hand because we need to multiply Schubert polynomials and expand in the basis of Schubert polynomials many times for each example.

Let $r[b, d] = [1, 2, \dots, b - 1, b + d, b, b + 1, \dots]$. Note that $D_{\text{bot}}(r[b, d])$ is a single row. The per-

mutation $r[b, d]$ is Grassmannian, and its Schubert polynomial equals $h_d(x_1, x_2, \dots, x_b)$, the homogeneous symmetric function of degree d .

Conjecture 6.1. For any $w \in S_\infty$ and any positive integers b and d , we have

$$\mathfrak{S}_w \mathfrak{S}_{r[b,d]} = \sum \mathfrak{S}_{w'},$$

where the sum runs over all $w' = wt_{k_1 l_1} t_{k_2 l_2} \dots t_{k_d l_d}$ such that $k_i \leq b < l_i$ for $1 \leq i \leq d$, and if we let $w^{(i)} = w^{(i-1)} t_{k_i l_i}$ with $w^{(0)} = w$, then $l(w^{(i)}) = l(w^{(i-1)}) + 1$ and

$$w_{k_1}^{(1)} < w_{k_2}^{(2)} < \dots < w_{k_d}^{(d)}.$$

Remark 6.2. It is remarkable that this multiplication is multiplicity free!

Let $c[b, d] = [1, 2, \dots, b - d, b - d + 2, \dots, b + 1, b - d + 1, b + 2, b + 3, \dots]$. Note that $D_{\text{top}}(c[b, d])$ is a single column. The permutation $c[b, d]$ is also Grassmannian, and its Schubert polynomial equals $e_d(x_1, x_2, \dots, x_b)$, the elementary symmetric function.

Conjecture 6.3. For any $w \in S_\infty$ and any positive integers b and d , we have

$$\mathfrak{S}_w \mathfrak{S}_{c[b,d]} = \sum \mathfrak{S}_{w'},$$

where the sum runs over all $w' = wt_{k_1 l_1} t_{k_2 l_2} \dots t_{k_d l_d}$ such that $k_i \leq b < l_i$ for $1 \leq i \leq d$, and if we let $w^{(i)} = w^{(i-1)} t_{k_i l_i}$ with $w^{(0)} = w$, then


$$l(w^{(i)}) = l(w^{(i-1)}) + 1$$

and

$$w_{k_1}^{(1)} > w_{k_2}^{(2)} > \dots > w_{k_d}^{(d)} > 0.$$

The conjectures have been verified by computer for all permutations $w \in S_7$ and all values of b and d . We have found computations in S_8 to be beyond the capacity of our computers (Sun Sparc 10). The validity of these conjectures would greatly speed up any algorithm for expanding the products $\mathfrak{S}_w \mathfrak{S}_{r[b,d]}$ and $\mathfrak{S}_w \mathfrak{S}_{c[b,d]}$. To see why this is so, we look at an example.

Suppose that we want to expand $\mathfrak{S}_{r[3,2]} \mathfrak{S}_{[1,2,5,4,3]}$ in the basis of Schubert polynomials. Let $b = 3$ and $d = 2$, and construct a rooted tree in the following way:



The top of the tree is the initial permutation. We assume there are an infinite number of fixed points beyond what is written. We have inserted a vertical line after the position $b = 3$. To find the children of the root, we find all transpositions that switch numbers across the vertical line so that the lengths increase by exactly one. We label the edge from the root to a child by the smallest of the two numbers switched. Of course the smallest number will always come from the left. This constructs the first generation of the tree. For the next generation, we repeat the process above, but only allowing the transpositions for which the smallest number is bigger than the label on the edge of this node. We repeat the last step $d = 2$ times. The leaves of the tree are precisely the permutations w' that appear in the expansion in Conjecture 6.1.

ACKNOWLEDGEMENTS

We would like to thank several people for valuable suggestions, in particular Sergey Fomin, Adriano Garsia, Mark Haiman, Arun Ram, Richard Stanley, and Glenn Tesler.

Nantel Bergeron, Department of Mathematics, Harvard University, Cambridge, MA 02138
 (nantel@math.harvard.edu)

Sara Billey, Department of Mathematics, University of California at San Diego, La Jolla, CA 92093-0112
 (sara@macaulay.ucsd.edu)

Received October 29, 1993; accepted in revised form January 15, 1994

REFERENCES

[Bergeron 1992] N. Bergeron, “A combinatorial construction of the Schubert polynomials”, *J. Comb. Theory A* **60** (1992), 168–182.

[Bernstein et al. 1973] I. N. Bernstein, I. M. Gelfand and S. I. Gelfand, “Schubert cells and cohomology of the spaces G/P ”, *Russian Math. Surveys* **28** (1973), 1–26.

[Billey et al. 1993] S. Billey, W. Jockusch and R. Stanley, “Some Combinatorial Properties of Schubert Polynomials”, *J. Alg. Comb.* (2) **4** (1993), 345–374.

[Demazure 1974] M. Demazure, “Désingularization des variétés de Schubert généralisées”, *Ann. Sci. Éc. Norm. Sup.* (Paris) **7** (1974) 53–58.

[Fomin and Kirillov 1993] S. Fomin and A. N. Kirillov, “Yang-Baxter equation, symmetric functions, and Schubert polynomials”, submitted to the *Proceedings of the Conference on Power Series and Algebraic Combinatorics, Firenze, 1993*.

[Fomin and Stanley 1991] S. Fomin and R. Stanley, “Schubert Polynomials and the NilCoxeter algebra”, *Adv. Math.* **103** (1994), 196–207.

[Kohnert 1990] A. Kohnert, “Weintrauben, Polynome, Tableaux”, *Bayreuth. Math. Schrift.* **38** (1990), 1–97.

[Lascoux and Schützenberger 1982] A. Lascoux and M.-P. Schützenberger, “Polynômes de Schubert”, *C. R. Acad. Sci. Paris* **294** (1982) 447–450.

[Lascoux and Schützenberger 1985] A. Lascoux and M.-P. Schützenberger, “Schubert polynomials and the Littlewood–Richardson Rule”, *Lett. Math. Phys.* **10** (1985) 111–124.

[Macdonald 1991] I. G. Macdonald, *Notes on Schubert Polynomials*, Publications du LACIM **6**, Université du Québec à Montréal (1991).

[Schubert 1879] H. Schubert, *Kalkül der Abzählenden Geometrie*, Teubner, Leipzig, 1879.