Heat kernel estimates and parabolic Harnack inequalities for symmetric Dirichlet forms

Zhen-Qing Chen, Takashi Kumagai and Jian Wang

Abstract

In this paper, we consider the following symmetric Dirichlet forms on a metric measure space (M, d, μ) :

$$\mathcal{E}(f,g) = \mathcal{E}^{(c)}(f,g) + \int_{M \times M} (f(x) - f(y))(g(x) - g(y)) J(dx,dy),$$

where $\mathcal{E}^{(c)}$ is a strongly local symmetric bilinear form and J(dx, dy) is a symmetric Random measure on $M \times M$. Under general volume doubling condition on (M, d, μ) and some mild assumptions on scaling functions, we establish stability results for upper bounds of heat kernel (resp. two-sided heat kernel estimates) in terms of the jumping kernels, the cut-off Sobolev inequalities, and the Faber-Krahn inequalities (resp. the Poincaré inequalities). We also obtain characterizations of parabolic Harnack inequalities. Our results apply to symmetric diffusions with jumps even when the underlying spaces have walk dimensions larger than 2.

AMS 2010 Mathematics subject classification: Primary 60J35, 35K08, 60J60, 60J75; Secondary 31C25, 60J25, 60J45.

Keywords and phrases: symmetric Dirichlet form, metric measure space, heat kernel estimate, parabolic Harnack inequality, stability, cut-off Sobolev inequality, generalized capacity inequality, jumping kernel, Lévy system

Contents

1	Introdu	iction and main results	2
	1.1 Set	ting and some history	2
	1.2 Fur	nctional inequalities and heat kernel estimates	6
	1.3 Par	abolic Harnack inequalities	15
2	2 Preliminaries		
	2.1 Pro	perties of ϕ_c and ϕ_i	18
	2.2 EP	$\phi_{\phi,\leq,\varepsilon} \Longrightarrow \operatorname{Gcap}(\phi) \text{ and } \operatorname{Gcap}(\phi) + J_{\phi,\leq} \Longrightarrow \operatorname{CS}(\phi). \ldots \ldots$	19
	2.3 ρ -ti	runcated version of $CS(\phi)$	22
	2.4 Self	f-improvement of $CS(\phi)$.	24
	2.5 Con	nsequences of $CS(\phi)$: Caccioppoli and L^1 -mean value inequalities	25
3 Implications of heat kernel estimates		tions of heat kernel estimates	29
	3.1 UH	$\operatorname{IK}(\phi_c, \phi_i) + (\mathcal{E}, \mathcal{F})$ is conservative $\Longrightarrow \operatorname{J}_{\phi_i, \leq}$ and $\operatorname{HK}(\phi_c, \phi_i) \Longrightarrow \operatorname{J}_{\phi_i} \ldots \ldots \ldots \ldots$	29
	3.2 UH	$\operatorname{IK}(\phi_c, \phi_j)$ and $(\mathcal{E}, \mathcal{F})$ is conservative $\Longrightarrow \operatorname{Gcap}(\phi) \dots \dots \dots \dots \dots \dots \dots \dots \dots $	30

4	$\begin{array}{ll} \text{Implications of } FK(\phi), \ NDL(\phi), \ CS(\phi) \ \text{and} \ J_{\phi_j,\leq} \\ 4.1 & NDL(\phi) \Longrightarrow PI(\phi) + E_{\phi} \\ 4.2 & FK(\phi) + J_{\phi,\leq} + CS(\phi) \Longrightarrow E_{\phi} \ \text{and} \ FK(\phi) + J_{\phi,\leq} + E_{\phi} \Longrightarrow UHKD(\phi) \\ . \\ 4.3 & UHKD(\phi) + J_{\phi_j,\leq} + E_{\phi} \Longrightarrow UHK(\phi_c,\phi_j) \\ \end{array}$	32 32 32 32
5	Characterizations of two-sided heat kernel estimates 5.1 $\operatorname{PI}(\phi) + \operatorname{J}_{\phi_j} + \operatorname{CS}(\phi) \Longrightarrow \operatorname{HK}_{-}(\phi_c, \phi_j) \ldots \ldots$	43 44 46
6	Characterizations of parabolic Harnack inequalities	
7	Examples/Applications	50

1 Introduction and main results

1.1 Setting and some history

Let (M, d, μ) be a *metric measure space*; that is, (M, d) is a locally compact separable metric space, and μ is a positive Radon measure on M with full support. We assume that all balls are relatively compact and assume for simplicity that $\mu(M) = \infty$ throughout the paper. Note that we do not assume M to be connected nor (M, d) to be geodesic.

Consider a regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(M; \mu)$. The Beurling-Deny formula asserts that such a form can be decomposed into the strongly local term, the pure-jump term and the killing term (see [FOT, Theorem 4.5.2]). Throughout this paper, we consider the Dirichlet form $(\mathcal{E}, \mathcal{F})$ having both the strongly local term and the pure-jump term, and having no killing term. That is,

$$\mathcal{E}(f,g) = \mathcal{E}^{(c)}(f,g) + \int_{M \times M \setminus \text{diag}} (f(x) - f(y)(g(x) - g(y)) J(dx, dy))$$

$$= : \mathcal{E}^{(c)}(f,g) + \mathcal{E}^{(j)}(f,g), \qquad f,g \in \mathcal{F},$$

(1.1)

where $(\mathcal{E}^{(c)}, \mathcal{F})$ is the strongly local part of $(\mathcal{E}, \mathcal{F})$ (namely $\mathcal{E}^{(c)}(f, g) = 0$ for all $f, g \in \mathcal{F}$ having (f - c)g = 0 μ -a.e. on M for some constant $c \in \mathbb{R}$) and $J(\cdot, \cdot)$ is a symmetric Radon measure $M \times M \setminus \text{diag}$; see [CF, Theorems 4.3.3 and 4.3.11]. Here and in what follows, we will always take a quasi-continuous version when we pick a function on \mathcal{F} (see [FOT, Theorem 2.1.3] for the definition and existence of a quasi-continuous version of the element in \mathcal{F}). In this paper, we assume that neither $\mathcal{E}^{(c)}(\cdot, \cdot)$ nor $J(\cdot, \cdot)$ are identically zero.

Let $(\mathcal{L}, \mathcal{D}(\mathcal{L}))$ be the L^2 -generator of $(\mathcal{E}, \mathcal{F})$ on $L^2(M; \mu)$; namely, \mathcal{L} is the self-adjoint operator on $L^2(M; \mu)$ and $\mathcal{D}(\mathcal{L})$ is the domain. $f \in \mathcal{D}(\mathcal{L})$ if $f \in \mathcal{F}$ and there exists (unique) $u \in L^2(M; \mu)$ such that

$$\mathcal{E}(f,g) = -\langle u,g \rangle$$
 for all $g \in \mathcal{F}$,

where $\langle \cdot, \cdot \rangle$ is the inner product in $L^2(M; \mu)$. We write $\mathcal{L}f := u$. Let $\{P_t\}_{t \geq 0}$ be the associated semigroup. Given a regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(M; \mu)$, there is an associated μ -symmetric Hunt process $X = \{X_t, t \geq 0; \mathbb{P}^x, x \in M \setminus \mathcal{N}\}$ where $\mathcal{N} \subset M$ is a properly exceptional set for $(\mathcal{E}, \mathcal{F})$, and the Hunt process is unique up to a properly exceptional set — see [FOT, Theorem 4.2.8] for details. We fix X and \mathcal{N} , and write $M_0 = M \setminus \mathcal{N}$. For any bounded Borel measurable function f on M, we may set

$$P_t f(x) = \mathbb{E}^x f(X_t), \quad x \in M_0.$$

The heat kernel associated with the semigroup $\{P_t\}_{t\geq 0}$ (if it exists) is a measurable function $p(t, x, y) : (0, \infty) \times M_0 \times M_0 \to (0, \infty)$ that satisfies the following:

$$\mathbb{E}^{x}f(X_{t}) = P_{t}f(x) = \int p(t, x, y)f(y)\,\mu(dy) \quad \text{for all } x \in M_{0}, f \in L^{\infty}(M; \mu), \tag{1.2}$$

$$p(t, x, y) = p(t, y, x)$$
 for all $t > 0, x, y \in M_0$, (1.3)

$$p(s+t,x,z) = \int p(s,x,y)p(t,y,z)\,\mu(dy) \quad \text{for all } s > 0, t > 0, \ x,z \in M_0.$$
(1.4)

It is possible to regularize p(t, x, y) so that (1.2)-(1.4) hold for every point in M_0 , see [BBCK, Theorem 3.1] and [GT, Section 2.2] for details. Note that in some arguments of our paper, we can extend (without further mention) p(t, x, y) to all $x, y \in M$ by setting p(t, x, y) = 0 if either x or y is outside M_0 .

There is a long history on the heat kernel estimates and related topics for strongly local Dirichlet forms. Let us briefly mention some of the previous works which are related to our work. For diffusions on manifolds, Grigor'yan [Gr1] and Saloff-Coste [Sa1] independently proved that the following are equivalent: (i) Aronson-type Gaussian bounds for heat kernel, (ii) parabolic Harnack equality, and (iii) VD and Poincaré inequality. The results are later extended to strongly local Dirichlet forms on metric measure spaces in [BM, St1, St2] and to graphs in [De]. Detailed heat kernel estimates are heavily related to the control of harmonic and parabolic functions, and the origin of ideas and techniques used in this field goes back to the work by De Giorgi, Nash, Moser and Aronson. For more details, see, for example, [Gr2, Sa2] and the references therein. For anomalous diffusions on disordered media such as fractals (where the so-called walk dimension being larger than 2), the above equivalence still holds but one needs to replace (i) by (i') sub-Gaussian bounds for heat kernel, (iii) by (iii') VD, Poincaré inequality and a cut-off Sobolev inequality; see [AB, BB2, BBK1, GHL].

For heat kernel estimates of symmetric jump processes in general metric measure spaces, when $\alpha \in (0, 2)$ and the metric measure space M is a *d*-set, characterizations of α -stable-like heat kernel estimates were obtained in [CK1] which are stable under rough isometries. This result has later been extended to mixed stable-like processes in [CK2] under some growth condition on the rate function ϕ such as

$$\int_0^r \frac{s}{\phi(s)} \, ds \le \frac{c \, r^2}{\phi(r)} \quad \text{for all } r > 0 \tag{1.5}$$

with some constant c > 0. For α -stable-like processes where $\phi(r) = r^{\alpha}$, condition (1.5) corresponds exactly to $0 < \alpha < 2$. Some of the key methods used in [CK1] were inspired by a previous work [BL] on random walks on integer lattice \mathbb{Z}^d . A long standing open problem in this field is to find a characterization of heat kernel estimates, which is stable under rough isometries, for α -stable-like processes even with $\alpha \geq 2$ when the underlying spaces have walk dimensions larger than 2. This question has been resolved recently in [CKW1] under some mild volume growth condition. Actually, in [CKW1] we obtained stability of two-sided heat kernel estimates and upper bound heat kernel estimates for symmetric jump processes with Ahlfors *d*-set condition in the framework of metric measure spaces [GHH] and in the framework of infinite connected locally finite graphs [MS]. The readers can further refer to [CKW2] for the stability results of parabolic Harnack inequalities for symmetric pure jump Dirichlet forms, and for [CKW3] for various characterizations of elliptic Harnack inequalities.

In this paper, we consider symmetric regular Dirichlet forms that have both the strongly local term and the pure-jump term. As mentioned above, we can also consider the corresponding operators and Hunt processes (diffusions with jumps). We use the following example, which is a special case of our much more general results, to illustrate the novelty and strength of our main results.

Example 1.1. Let $U \subset \mathbb{R}^d$ be an unbounded global Lipschitz domain equipped with the Euclidean distance. Let $X = \{X_t\}_{t\geq 0}$ be a symmetric reflected diffusion with jumps on \overline{U} associated with the regular Dirichlet form $(\mathcal{E}, W^{1,2}(U))$ on $L^2(U; dx)$ given by

$$\mathcal{E}(u,v) = \int_{U} \nabla u(x) \cdot A(x) \nabla v(x) \, dx + \int_{U} \int_{U} (u(x) - u(y)) (v(x) - v(y)) \frac{c(x,y)}{|x - y|^{d + \alpha}} \, dx \, dy, \quad (1.6)$$

where $A(x) = (a_{ij}(x))_{1 \le i,j \le d}$ is a measurable uniformly elliptic and bounded $d \times d$ matrix-valued function on U, $0 < \alpha < 2$, and $c(\cdot, \cdot)$ is a symmetric measurable function on $U \times U$ that is bounded between two positive constants. Its L^2 -infinitesimal generator is of the form

$$\mathcal{L}u(x) = \sum_{i,j=1}^{d} \frac{\partial}{\partial x_i} \left(a_{ij}(x) \frac{\partial u(x)}{\partial x_j} \right) + 2 \lim_{\varepsilon \to 0} \int_{\{y \in U: |y-x| > \varepsilon\}} (u(y) - u(x)) \frac{c(x,y)}{|x-y|^{d+\alpha}} \, dy$$

with "Neumann" boundary condition. Then $\text{HK}(\phi_c, \phi_j)$ and $\text{PHI}(\phi)$ hold, where $\text{HK}(\phi_c, \phi_j)$ (reps. $\text{PHI}(\phi)$) is the detailed two-sided heat kernel estimates defined in (1.30) (resp. the parabolic Harnack inequality defined in (1.33)) with $\phi_c(r) = r^2$, $\phi_j(r) = r^{\alpha}$ and $\phi(r) = \phi_c(r) \land \phi_j(r)$.

To the best of our knowledge, even this result is new. When $U = \mathbb{R}^d$, these results are first obtained in [CK3]. See [CKKW2] for a different approach to this example without using the stability results of this paper. The proof of Example 1.1 will be given in Section 7.

Although it is very natural and important to study heat kernels for symmetric Dirichlet forms that have both the diffusive and jumping parts, there are very limited work in literature on this topic. For a Lévy process X that is the independent sum of a Brownian motion W and a symmetric α -stable process Y on Euclidean spaces, its transition density is the convolution of the transition densities of W and Y. In [SV], heat kernel estimates are derived for X by computing the convolution in four cases. In one of cases (the case of $|x|^2 < t < |x|^{\alpha} < 1$), the upper and lower bounds do not match. Nevertheless, parabolic Harnack inequality for X can be obtained from these estimates. In [CK3], sharp and comparable upper and lower bounds are obtained for a large class of symmetric diffusions with mixture stable-like jumps on Euclidean spaces. This sharp two-sided heat kernel estimates were new even for Lévy processes that are the independent sum of Brownian motion and symmetric stable processes. The results of [CK3] have been further extended to general metric measure spaces in [CKKW2] and to the cases where the jumping kernels can have exponential decay. One of the difficulties in obtaining fine properties for diffusions with jumps and associated operators is that it exhibits two different scales: the strongly local terms part has a diffusion scaling $r \mapsto \phi_c(r)$ while the pure jump part has a different type of scaling $r \mapsto \phi_i(r)$. On the other hand, as shown in [CKW2], in contrast to the cases of local operators/diffusions, for symmetric pure-jump processes, parabolic Harnack inequalities are no longer equivalent to (in fact weaker than) the two-sided heat kernel estimates. This discrepancy is caused by the heavy tail of the jumping kernel. This heavy tail phenomenon is also one of main sources of difficulties in analyzing non-local operators/jump processes. Diffusions with

jumps are even more complex than pure jumps case studied in [CKW2], in fact Theorem 1.17 of this paper asserts that, with $\phi(r) := \phi_c(r) \wedge \phi_j(r)$,

$$\operatorname{HK}_{-}(\phi_{c}, \phi_{j}) \iff \operatorname{PHI}(\phi) + \operatorname{J}_{\phi_{j}};$$

see Definition 1.11(ii) and (1.17) for definitions of $HK_{-}(\phi_c, \phi_j)$ and J_{ϕ_j} , respectively. In the pure jump case, it holds that

$$\operatorname{HK}(\phi_j) \iff \operatorname{PHI}(\phi_j) + \mathcal{J}_{\phi_j,>}$$

see [CKW2, Corollary 1.21]. Intuitively speaking, this discrepancy is due to the fact that the scale corresponding to the small time behavior of diffusions with jumps is dominated by the diffusive part and hence one can not recover information about the jumping scale function $\phi_j(r)$ for $r \leq 1$ from PHI(ϕ).

Due to the above difficulties and differences, obtaining the complete picture of heat kernel estimates, and the stability of heat kernel estimates and parabolic Harnack inequalities for symmetric Dirichlet forms including both local and non-local terms/diffusions with jumps requires new ideas. Our approach contains the following four key ingredients, and all of them are highly non-trivial:

- (i) We adopt the generalized capacity inequality formulation from the recent study on stablelike jumps processes under the Ahlfors *d*-set condition in the framework of metric measure spaces [GHH], and make use of the arguments depending on cut-off Sobolev inequality from [AB, CKW1] to derive some useful analytical properties of Dirichlet forms consisting of both strongly local terms and pure jumps terms. The generalized capacity condition $Gcap(\phi)$ is clean to state but it is not known whether it is stable under rough isometry or not, while the cut-off Sobolev inequality condition $CS(\phi)$ is lengthy to state but is clearly stable under rough isometry.
- (ii) We find a new self-improving argument for upper bounds for diffusions with jumps. The advantage of this technique is that it not only can take care of different scales both from the strongly local term part and the pure jump term, but also can treat the case that the volume of balls is not uniformly comparable.
- (iii) As mentioned above, different from the assertions for diffusions or symmetric pure jump processes, in the present setting parabolic Harnack inequalities are not equivalent to twosided heat kernel estimates. Moreover, the parabolic Harnack inequalities alone can not imply bounds (even upper bounds) of the jumping kernel. So, to obtain the characterizations of parabolic Harnack inequalities for diffusions with jumps we shall consider the weaker upper bounds of jumping kernels $J_{\phi,\leq}$ instead of the exact upper bounds of jumping kernels $J_{\phi_j,\leq}$. This indicates that only rough upper bounds of heat kernels UHK_{weak} together with NDL(ϕ) and UJS (see (4.10), Definitions 1.11(vi) and 1.16 for their definitions) is involved in the characterization of parabolic Harnack inequalities; see Theorem 1.17.
- (iv) Our results are obtained under some mild volume growth condition, called volume doubling and reverse volume doubling conditions; see Definition 1.2 for details. This is much weaker than the Ahlfors *d*-set condition and significant technical difficulties arise when working under this setting.

1.2 Functional inequalities and heat kernel estimates

In this paper, we are concerned with stable characterizations of both upper bounds and two-sided estimates on heat kernel, as well as of parabolic Harnack inequalities, for symmetric Dirichlet forms having both local and non-local terms on general metric measure spaces. To state our results for heat kernel estimates precisely, we need a number of definitions; some of them are taken from [CKW1]. Denote the ball centered at x with radius r by B(x,r) and $\mu(B(x,r))$ by V(x,r).

Definition 1.2. (i) We say that (M, d, μ) satisfies the volume doubling property (VD), if there exists a constant $C_{\mu} \ge 1$ such that for all $x \in M$ and r > 0,

$$V(x,2r) \le C_{\mu}V(x,r). \tag{1.7}$$

(ii) We say that (M, d, μ) satisfies the reverse volume doubling property (RVD), if there exist constants $l_{\mu}, c_{\mu} > 1$ such that for all $x \in M$ and r > 0,

$$V(x, l_{\mu}r) \ge c_{\mu}V(x, r). \tag{1.8}$$

VD condition (1.7) (resp. RVD condition (1.8)) is equivalent to the second (resp. the first) inequality in the following display:

$$\widetilde{c}_{\mu}\left(\frac{R}{r}\right)^{d_{1}} \leq \frac{V(x,R)}{V(x,r)} \leq \widetilde{C}_{\mu}\left(\frac{R}{r}\right)^{d_{2}} \quad \text{for all } x \in M \text{ and } R \geq r > 0,$$
(1.9)

where $d_1, d_2, \tilde{c}_{\mu}$ and \tilde{C}_{μ} are positive constants. Under RVD, $\mu(M) = \infty$ if and only if M has infinite diameter. If M is connected and unbounded, then VD implies RVD; see [GH, Proposition 5.1 and Corollary 5.3].

Let $\mathbb{R}_+ := [0, \infty)$, and $\phi_c : \mathbb{R}_+ \to \mathbb{R}_+$ (resp. $\phi_j : \mathbb{R}_+ \to \mathbb{R}_+$) be a strictly increasing continuous function with $\phi_c(0) = 0$ (resp. $\phi_j(0) = 0$), $\phi_c(1) = 1$ (resp. $\phi_j(1) = 1$) and satisfying that there exist constants $c_{1,\phi_c}, c_{2,\phi_c} > 0$ and $\beta_{2,\phi_c} \ge \beta_{1,\phi_c} > 1$ (resp. $c_{1,\phi_j}, c_{2,\phi_j} > 0$ and $\beta_{2,\phi_j} \ge \beta_{1,\phi_j} > 0$) such that

$$c_{1,\phi_c} \left(\frac{R}{r}\right)^{\beta_{1,\phi_c}} \leq \frac{\phi_c(R)}{\phi_c(r)} \leq c_{2,\phi_c} \left(\frac{R}{r}\right)^{\beta_{2,\phi_c}} \quad \text{for all } 0 < r \leq R.$$

$$\left(\text{resp. } c_{1,\phi_j} \left(\frac{R}{r}\right)^{\beta_{1,\phi_j}} \leq \frac{\phi_j(R)}{\phi_j(r)} \leq c_{2,\phi_j} \left(\frac{R}{r}\right)^{\beta_{2,\phi_j}} \quad \text{for all } 0 < r \leq R.\right)$$

$$(1.10)$$

Note that (1.10) is equivalent to the existence of constants c_{3,ϕ_c} , $l_{0,\phi_c} > 1$ such that

$$c_{3,\phi_c}^{-1}\phi_c(r) \le \phi_c(l_{0,\phi_c}r) \le c_{3,\phi_c}\phi_c(r)$$
 for all $r > 0$,

the same as $\phi_i(r)$. Throughout the paper, we assume that

$$\phi_c(r) \le \phi_j(r) \text{ for } r \in (0,1] \text{ and } \phi_c(r) \ge \phi_j(r) \text{ for } r \in [1,\infty).$$
 (1.11)

Since $\beta_{1,\phi_c} > 1$, by [BGT, Definition, p. 65; Definition, p. 66; Theorem 2.2.4 and its remark, p. 73], there exists a strictly increasing continuous function $\bar{\phi}_c(r) : \mathbb{R}_+ \to \mathbb{R}_+$ such that there are constants $c_2 \ge c_1 > 0$ so that

$$c_1 \frac{\phi_c(r)}{r} \le \bar{\phi}_c(r) \le c_2 \frac{\phi_c(r)}{r} \quad \text{for all } r > 0.$$

$$(1.12)$$

By (1.10), $\beta_{1,\phi_c} > 1$ and (1.12), we have $\lim_{r\to 0} \bar{\phi}_c(r) = 0$, $\lim_{r\to\infty} \bar{\phi}_c(r) = \infty$, and there are constants $c_{1,\bar{\phi}_c}, c_{2,\bar{\phi}_c} > 0$ such that

$$c_{1,\bar{\phi}_c} \left(\frac{R}{r}\right)^{\beta_{1,\phi_c}-1} \le \frac{\bar{\phi}_c(R)}{\bar{\phi}_c(r)} \le c_{2,\bar{\phi}_c} \left(\frac{R}{r}\right)^{\beta_{2,\phi_c}-1} \quad \text{for all } 0 < r \le R.$$
(1.13)

Given ϕ_c and ϕ_j satisfying (1.11), we set

$$\phi(r) := \phi_c(r) \land \phi_j(r) = \begin{cases} \phi_c(r), & r \in (0, 1], \\ \phi_j(r), & r \in [1, \infty). \end{cases}$$
(1.14)

It is clear that $\phi(r)$ is a strictly increasing function on \mathbb{R}_+ with $\phi(0) = 0$ and $\phi(1) = 1$, and satisfies that there exist constants $c_{1,\phi}, c_{2,\phi} > 0$ so that

$$c_{1,\phi} \left(\frac{R}{r}\right)^{\beta_{1,\phi}} \le \frac{\phi(R)}{\phi(r)} \le c_{2,\phi} \left(\frac{R}{r}\right)^{\beta_{2,\phi}} \quad \text{for all } 0 < r \le R,$$
(1.15)

where $\beta_{1,\phi} = \beta_{1,\phi_c} \wedge \beta_{1,\phi_j}$ and $\beta_{2,\phi} = \beta_{2,\phi_c} \vee \beta_{2,\phi_j}$. Throughout this paper, without any mention we will fix the notations for these three functions ϕ_c , ϕ_j and ϕ .

Definition 1.3. Let $\psi : \mathbb{R}_+ \to \mathbb{R}_+$. We say condition J_{ψ} holds if there exists a non-negative symmetric function J(x, y) on $M \times M$ so that for $\mu \times \mu$ -almost all $x, y \in M$,

$$J(dx, dy) = J(x, y) \,\mu(dx) \,\mu(dy), \tag{1.16}$$

and

$$\frac{c_1}{V(x, d(x, y))\psi(d(x, y))} \le J(x, y) \le \frac{c_2}{V(x, d(x, y))\psi(d(x, y))}.$$
(1.17)

We say that $J_{\psi,\leq}$ (resp. $J_{\psi,\geq}$) if (1.16) holds and the upper bound (resp. lower bound) in (1.17) holds.

Note that, without loss of generality, we may and do assume that in condition J_{ψ} $(J_{\psi,\geq})$ and $J_{\psi,\leq}$, respectively) that (1.17) (and the corresponding inequality) holds for every $x, y \in M$. Note also that, under VD, the bounds in condition (1.17) are consistent with the symmetry of J(x, y). See [CKW1, Remark 1.3] for more details.

Since $\phi(r) \leq \phi_j(r)$ for all r > 0, $J_{\phi_j,\leq}$ implies $J_{\phi,\leq}$; that is, condition $J_{\phi,\leq}$ is weaker than condition $J_{\phi_j,\leq}$. We will frequently use this fact in the paper.

Definition 1.4. Let $U \subset V$ be open sets of M with $U \subset \overline{U} \subset V$, and $\kappa \geq 1$. We say a non-negative bounded measurable function φ is a κ -cut-off function for $U \subset V$, if $\varphi \geq 1$ on U, $\varphi = 0$ on V^c and $0 \leq \varphi \leq \kappa$ on M. Any 1-cut-off function is simply referred to as a cut-off function.

It is obvious that for any κ -cut-off function φ for $U \subset V$, $1 \land \varphi$ is a cut-off function for $U \subset V$.

Motivated by [GHH] for pure jump Dirichlet forms, we formulate the generalized capacity condition for non-local Dirichlet forms that have diffusive parts. For this, we consider the following function space

$$\mathcal{F}'_b := \{ u + a : u \in \mathcal{F}_b, \, a \in \mathbb{R} \},\$$

where $\mathcal{F}_b := \mathcal{F} \cap L^{\infty}(M; \mu)$.

Definition 1.5. We say that the generalized capacity inequality $\text{Gcap}(\phi)$ holds, if there exist constants $\kappa \geq 1$ and C > 0 such that for every 0 < r < R, any $f \in \mathcal{F}'_b$ and for almost all $x_0 \in M$, there is a κ -cut-off function $\varphi \in \mathcal{F}_b$ for $B(x_0, R) \subset B(x_0, R+r)$ so that

$$\mathcal{E}(f^2\varphi,\varphi) \le \frac{C}{\phi(r)} \int_{B(x_0,R+r)} f^2 \, d\mu$$

Recall that for any subsets $A \subset B$, the relative capacity $\operatorname{cap}(A, B)$ is defined by

 $\operatorname{cap}(A,B) = \inf \{ \mathcal{E}(\varphi,\varphi) : \varphi \in \mathcal{F}_b \text{ is a cut-off function for } A \subset B \}.$

Following [GHH, Definition 1.7], for any subsets $A \subset B$, $f \in \mathcal{F}'_b$ and a constant $\kappa \ge 1$, we define the generalized relative capacity $\operatorname{cap}_f^{(\kappa)}(A, B)$ by

$$\operatorname{cap}_{f}^{(\kappa)}(A,B) = \inf \left\{ \mathcal{E}(f^{2}\varphi,\varphi) : \varphi \in \mathcal{F}_{b} \text{ is a } \kappa \text{-cut-off function for } A \subset B \right\}.$$

In particular, when f = 1 and $\kappa = 1$, $\operatorname{cap}_{f}^{(\kappa)}(A, B) = \operatorname{cap}(A, B)$. As mentioned in [GHH, Remarks 1.8 and 1.9], the quantity $\mathcal{E}(f^{2}\varphi, \varphi)$ in the definition of the generalized capacity is well defined, and the generalized capacity can take negative values. With this notation, the generalized capacity inequality $\operatorname{Gcap}(\phi)$ is equivalent to that there exist constants $\kappa \geq 1$ and C > 0 such that for every 0 < r < R, any $u \in \mathcal{F}'_{b}$ and almost all $x_{0} \in M$ so that

$$\operatorname{cap}_{u}^{(\kappa)}(B(x_0, R), B(x_0, R+r)) \le \frac{C}{\phi(r)} \int_{B(x_0, R+r)} u^2 d\mu.$$

Denote by $C_c(M)$ the space of continuous functions on M with compact support. It is well known that for any $f \in \mathcal{F}_b$, there exist unique positive Random measures $\Gamma(f, f)$ and $\Gamma_c(f, f)$ on M so that for every $g \in \mathcal{F} \cap C_c(M)$,

$$\int_M g \, d\Gamma(f, f) = \mathcal{E}(f, fg) - \frac{1}{2} \mathcal{E}(f^2, g),$$

and

$$\int_M g \, d\Gamma_c(f,f) = \mathcal{E}^{(c)}(f,fg) - \frac{1}{2} \mathcal{E}^{(c)}(f^2,g).$$

The energy measures $\Gamma(f, f)$ and $\Gamma_c(f, f)$ can be uniquely extended to any $f \in \mathcal{F}$ as the increasing limit of $\Gamma(f_n, f_n)$ and $\Gamma_c(f_n, f_n)$, respectively, where $f_n := ((-n) \lor f) \land n$. The measure $\Gamma(f, f)$ (resp. $\Gamma_c(f, f)$) is called the *energy measure* of f (which is also called the *carré* du champ in the literature) for \mathcal{E} (resp. its strongly local part $\mathcal{E}^{(c)}$).

To make use of the generalized capacity inequality $Gcap(\phi)$, we need to introduce a version of a cut-off Sobolev inequality that controls the energy of cut-off functions.

Definition 1.6. We say that condition $CS(\phi)$ holds if there exist constants $C_0 \in (0, 1]$ and $C_1, C_2 > 0$ such that for every $0 < r \leq R$, almost all $x_0 \in M$ and any $f \in \mathcal{F}$, there exists a

cut-off function $\varphi \in \mathcal{F}_b$ for $B(x_0, R) \subset B(x_0, R+r)$ so that the following holds:

$$\int_{B(x_0,R+(1+C_0)r)} f^2 d\Gamma(\varphi,\varphi)
\leq C_1 \left(\int_{B(x_0,R+r)} \varphi^2 d\Gamma_c(f,f)
+ \int_{B(x_0,R+r) \times B(x_0,R+(1+C_0)r)} \varphi^2(x) (f(x) - f(y))^2 J(dx,dy) \right)
+ \frac{C_2}{\phi(r)} \int_{B(x_0,R+(1+C_0)r)} f^2 d\mu.$$
(1.18)

- Remark 1.7. (i) Clearly, unlike $Gcap(\phi)$, condition $CS(\phi)$ is stable under rough isometry. $CS(\phi)$ is a combination of $CSA(\phi)$ for strongly local Dirichlet forms and $CSJ(\phi)$ for pure jump Dirichlet forms. $CSA(\phi)$ was introduced in [AB] for strongly local Dirichlet forms as a weaker version of the so called cut-off Sobolev inequality in [BB2, BBK1]; while $CSJ(\phi)$. as a counterpart of $CSA(\phi)$ for pure jump Dirichlet form, was given in [CKW1]. As pointed out in [CKW1, Remark 1.6(ii)], the main difference between $CSJ(\phi)$ and $CSA(\phi)$ is that the integrals in the left hand side and in the second term of the right hand side of the inequality (1.18) are over $B(x_0, R + (1 + C_0)r)$ instead of over $B(x_0, R + r)$ for [AB]. Note that the integral over $B(x_0, R+r)^c$ is zero in the left hand side of (1.18) for the case of strongly local Dirichlet forms. As we see in [CKW1] for the arguments of the stability of heat kernel estimates for jump processes, it is important to enlarge the ball $B(x_0, R+r)$ and integrate over $B(x_0, R + (1 + C_0)r)$ rather than over $B(x_0, R + r)$. In the present setting, we will deal with Dirichlet forms having both local and non-local parts (i.e., the associated Hunt process have both the diffusive and jumping parts), and so it is natural to use the formula similar to CSJ(ϕ) of [CKW1]. As supp [φ] $\subset B(x_0, R+r)$, we could replace $B(x_0, R+r)$ by $B(x_0, R+(1+C_0)r)$ in the integral region of the first term on the right hand side of (1.18).
 - (ii) Denote by \mathcal{F}_{loc} the space of functions locally in \mathcal{F} ; that is, $f \in \mathcal{F}_{loc}$ if and only if for any relatively compact open set $U \subset M$ there exists $g \in \mathcal{F}$ such that $f = g \mu$ -a.e. on U. Since each ball is relatively compact and (1.18) uses the property of f on $B(x_0, R + (1 + C_0)r)$ only, $CS(\phi)$ also holds for any $f \in \mathcal{F}_{loc}$.
- (iii) As mentioned in [GHL, page 1492] and [CKW1, Remark 1.7], if the non-negative cut-off function φ in CS(ϕ) can be chosen as a Lipschitz continuous function, then CS(ϕ) always holds under VD, (1.10) and $J_{\phi,\leq}$. For example, this is the case that any geodesically complete Riemannian manifold with $\phi_c(r) = r^2$ and $\phi_j(r) = r^{\alpha}$ with some $\alpha \in (0, 2)$. See also Example 1.1.

For $\alpha > 0$, define

$$\mathcal{E}_{\alpha}(f,g) = \mathcal{E}(f,g) + \alpha \int_{M} f(x)g(x)\,\mu(dx) \quad \text{for } f,g \in \mathcal{F}.$$

We next introduce the Faber-Krahn inequality. For any open set $D \subset M$, let \mathcal{F}^D be the \mathcal{E}_1 -closure of $\mathcal{F} \cap C_c(D)$ in \mathcal{F} . Define

$$\lambda_1(D) = \inf \left\{ \mathcal{E}(f, f) : f \in \mathcal{F}^D \text{ with } \|f\|_2 = 1 \right\},$$
(1.19)

the bottom of the Dirichlet spectrum of the corresponding self-adjoint operator in D.

Definition 1.8. We say that the *Faber-Krahn inequality* $FK(\phi)$ holds if there exist positive constants C and ν such that for any ball B(x, r) and any open set $D \subset B(x, r)$,

$$\lambda_1(D) \ge \frac{C}{\phi(r)} (V(x,r)/\mu(D))^{\nu}.$$
 (1.20)

Since $V(x,r) \ge \mu(D)$ for $D \subset B(x,r)$, if (1.20) holds for some $\nu = \nu_0 > 0$, it then holds for every $\nu \in (0, \nu_0)$. So without loss of generality, we may and do assume $0 < \nu < 1$.

Definition 1.9. We say that the (weak) Poincaré inequality $PI(\phi)$ holds if there exist constants C > 0 and $\kappa \ge 1$ such that for any ball $B_r = B(x, r)$ with $x \in M$ and for any $f \in \mathcal{F}_b$,

$$\int_{B_r} (f - \overline{f}_{B_r})^2 d\mu \le C\phi(r) \left(\int_{B_{\kappa r}} \Gamma_c(f, f) + \int_{B_{\kappa r} \times B_{\kappa r}} (f(y) - f(x))^2 J(dx, dy) \right), \quad (1.21)$$

where Γ_c is the energy measure of local bilinear form of $(\mathcal{E}, \mathcal{F})$, and $\overline{f}_{B_r} = \frac{1}{\mu(B_r)} \int_{B_r} f d\mu$ is the average value of f on B_r .

If the integral on the right hand side of (1.21) is over $B_r \times B_r$ (i.e. $\kappa = 1$), then it is called strong Poincaré inequality. If the metric is geodesic, it is known that (weak) Poincaré inequality implies strong Poincaré inequality (see for instance [Sa1, Section 5.3]), but in general they are not the same. In this paper, we only use weak Poincaré inequality. Note also that the left hand side of (1.21) is equal to $\inf_{a \in \mathbb{R}} \int_{B_r} (f-a)^2 d\mu$.

Recall that $X = \{X_t\}_{t\geq 0}$ is the Hunt process associated with the regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(M; \mu)$ with properly exceptional set \mathcal{N} , and $M_0 := M \setminus \mathcal{N}$. For a set $A \subset M$, define the exit time $\tau_A = \inf\{t > 0 : X_t \notin A\}$.

Definition 1.10. (i) We say that E_{ϕ} holds if there is a constant $c_1 > 1$ such that for all r > 0 and all $x \in M_0$,

$$c_1^{-1}\phi(r) \le \mathbb{E}^x[\tau_{B(x,r)}] \le c_1\phi(r).$$

We say that $E_{\phi,\leq}$ (resp. $E_{\phi,\geq}$) holds if the upper bound (resp. lower bound) in the inequality above holds.

(ii) We say $EP_{\phi,\leq}$ holds if there is a constant c > 0 such that for all r, t > 0 and all $x \in M_0$,

$$\mathbb{P}^x(\tau_{B(x,r)} \le t) \le \frac{ct}{\phi(r)}.$$

We say $EP_{\phi,\leq,\varepsilon}$ holds, if there exist constants $\varepsilon, \delta \in (0,1)$ such that for any $x_0 \in M$ and r > 0,

$$\mathbb{P}^x(\tau_{B(x_0,r)} \leq \delta\phi(r)) \leq \varepsilon$$
 for all $x \in B(x_0,r/4) \cap M_0$.

It is clear that $EP_{\phi,\leq}$ implies $EP_{\phi,\leq,\varepsilon}$. It is also easy to see that, under (1.10), E_{ϕ} implies $EP_{\phi,<,\varepsilon}$; see Proposition 2.4 below.

We use $\phi_c^{-1}(t)$ (resp. $\phi_j^{-1}(t)$) to denote the inverse function of the strictly increasing function $t \mapsto \phi_c(t)$ (resp. $t \mapsto \phi_j(t)$). Throughout the paper, we write $f(s,x) \simeq g(s,x)$, if there exist constants $c_1, c_2 > 0$ such that $c_1g(s,x) \leq f(s,x) \leq c_2g(s,x)$ for the specified range of the argument (s,x). Similarly, we write $f(s,x) \approx g(s,x)$, if there exist constants $c_k > 0$, $k = 1, \dots, 4$, such that $c_1g(c_2s,x) \leq f(s,x) \leq c_3f(c_4s,x)$ for the specified range of (s,x).

We consider the following two-sided estimates of heat kernel for the local Dirichlet forms. Define

$$p^{(c)}(t,x,y) := \frac{1}{V(x,\phi_c^{-1}(t))} \exp\left(-\sup_{s>0}\left\{\frac{d(x,y)}{s} - \frac{t}{\phi_c(s)}\right\}\right), \quad t > 0, x, y \in M_0.$$
(1.22)

This kernel arises in the two-sided estimates of heat kernel for strongly local Dirichlet forms; see e.g. [AB]. In the literature (see [GT, HK]), there is another expression of two-sided heat kernel estimates for the local Dirichlet forms given by

$$p^{(c)}(t,x,y) = \frac{1}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right), \quad t > 0, x, y \in M_0.$$
(1.23)

Note that

$$m(t,r) := \frac{r}{\bar{\phi}_c^{-1}(t/r)}$$
(1.24)

is the unique solution of

$$\bar{\phi}_c\left(\frac{r}{m(t,r)}\right) = \frac{t}{r}, \quad t,r > 0.$$
(1.25)

We will show that (1.22) and (1.23) are equivalent to each other in our setting in the sense that there are constants $c_k > 0$, $k = 1, \dots, 4$, so that for $p^{(c)}(t, x, y)$ given by (1.22),

$$\frac{c_1}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{c_2 d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right) \le p^{(c)}(t,x,y) \\
\le \frac{c_3}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{c_4 d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right)$$
(1.26)

for every t > 0 and $x, y \in M_0$. See Lemma 2.2 and Corollary 2.3 for the proofs. On the other hand, we note that, in all the literature we know, for example [BB1], [HK] and [GT, Page 1217– 1218], the lower bound in the estimate (1.23) (more explicitly, the lower bound of off-diagonal estimate in (1.23)) was established under assumptions that include (M, d, μ) being connected and satisfying the chain condition; that is, there exists a constant C > 0 such that, for any $x, y \in M$ and for any $n \in \mathbb{N}$, there exists a sequence $\{x_i\}_{i=0}^n \subset M$ such that $x_0 = x, x_n = y$ and $d(x_i, x_{i+1}) \leq Cd(x, y)/n$ for all $i = 0, 1, \dots, n-1$.

In the following, we designate (1.23) as the expression of $p^{(c)}(t, x, y)$. Note that

$$\frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))} \preceq 1 \quad \text{if and only if} \quad \phi_c(d(x,y))/t \preceq 1$$

So for each fixed a > 0,

$$p^{(c)}(t, x, y) \simeq 1/V(x, \phi_c^{-1}(t))$$
 when $d(x, y) \le a\phi_c^{-1}(t)$. (1.27)

Set

$$p^{(j)}(t,x,y) := \frac{1}{V(x,\phi_j^{-1}(t))} \wedge \frac{t}{V(x,d(x,y))\phi_j(d(x,y))}.$$
(1.28)

It is easy to see that for each fixed a > 0,

$$p^{(j)}(t,x,y) \simeq \frac{1}{V(x,\phi_j^{-1}(t))}$$
 when $d(x,y) \le a\phi_j^{-1}(t)$. (1.29)

Definition 1.11. (i) We say that $HK(\phi_c, \phi_j)$ holds if there exists a heat kernel p(t, x, y) of the semigroup $\{P_t\}_{t\geq 0}$ associated with $(\mathcal{E}, \mathcal{F})$ and the following estimates hold for all t > 0 and all $x, y \in M_0$,

$$c_{1}\left(\frac{1}{V(x,\phi_{c}^{-1}(t))}\wedge\frac{1}{V(x,\phi_{j}^{-1}(t))}\wedge\left(p^{(c)}(c_{2}t,x,y)+p^{(j)}(t,x,y)\right)\right)$$

$$\leq p(t,x,y) \tag{1.30}$$

$$\leq c_{3}\left(\frac{1}{V(x,\phi_{c}^{-1}(t))}\wedge\frac{1}{V(x,\phi_{j}^{-1}(t))}\wedge\left(p^{(c)}(c_{4}t,x,y)+p^{(j)}(t,x,y)\right)\right),$$

where $c_k > 0$, $k = 1, \dots, 4$, are constants independent of $x, y \in M_0$ and t > 0. For simplicity and by abusing the notation, we abbreviate the two-sided estimate (1.30) as

$$p(t, x, y) \asymp \frac{1}{V(x, \phi_c^{-1}(t))} \wedge \frac{1}{V(x, \phi_j^{-1}(t))} \wedge \left(p^{(c)}(t, x, y) + p^{(j)}(t, x, y) \right).$$

(ii) We say $HK_{-}(\phi_c, \phi_j)$ holds if the upper bound in (1.30) holds but the lower bound is replaced by the following: there are constants $c_0, c_1 > 0$ so that for all $x, y \in M_0$,

$$p(t, x, y) \ge c_0 \left(\frac{1}{V(x, \phi^{-1}(t))} \mathbf{1}_{\{d(x, y) \le c_1 \phi^{-1}(t)\}} + \frac{t}{V(x, d(x, y)) \phi_j(d(x, y))} \mathbf{1}_{\{d(x, y) > c_1 \phi^{-1}(t)\}} \right).$$

$$(1.31)$$

- (iii) We say UHK(ϕ_c, ϕ_j) (resp. LHK(ϕ_c, ϕ_j)) holds if the upper bound (resp. the lower bound) in (1.30) holds.
- (iv) We say UHKD(ϕ) holds if there is a constant c > 0 such that for all t > 0 and all $x \in M_0$,

$$p(t, x, x) \le \frac{c}{V(x, \phi^{-1}(t))}.$$

(v) We say a near-diagonal lower bound heat kernel estimate $NL(\phi)$ holds if there are constants $c_1, c_2 > 0$ such that for all t > 0 and all $x, y \in M_0$ with $d(x, y) \le c_1 \phi^{-1}(t)$,

$$p(t, x, y) \ge \frac{c_2}{V(x, \phi^{-1}(t))}.$$

(vi) Denote by $(P_t^D)_{t\geq 0}$ the (Dirichlet) semigroups of $(\mathcal{E}, \mathcal{F}^D)$, and by $p^D(t, x, y)$ the corresponding (Dirichlet) heat kernel. We say that a near diagonal lower bounded estimate for Dirichlet heat kernel NDL(ϕ) holds, if there exist $\varepsilon \in (0, 1)$ and $c_1 > 0$ such that for any $x_0 \in M, r > 0, 0 < t \le \phi(\varepsilon r)$ and $B = B(x_0, r)$,

$$p^{B}(t, x, y) \ge \frac{c_{1}}{V(x_{0}, \phi^{-1}(t))}, \quad x, y \in B(x_{0}, \varepsilon \phi^{-1}(t)) \cap M_{0}.$$
 (1.32)

Remark 1.12. We have five remarks about this definition.



Figure 1: Dominant term in the heat kernel estimates $HK(\phi_c, \phi_j)$ for p(t, x, y)

(i) Note that the scaling of $\text{HK}(\phi_c, \phi_j)$ (also $\text{HK}_-(\phi_c, \phi_j)$, $\text{UHK}(\phi_c, \phi_j)$ and $\text{LHK}(\phi_c, \phi_j)$) is NOT completely determined by ϕ . Indeed, it also includes the information of $\phi_j(r)$ for $0 < r \leq 1$. Yet, we use this notation since ϕ gives us the space-time relation of the heat kernel estimates. Furthermore, it follows from Theorem 1.17 that $\text{HK}_-(\phi_c, \phi_j)$ (and so $\text{HK}(\phi_c, \phi_j)$) is stronger than $\text{PHI}(\phi)$, and, consequently, $\text{PHR}(\phi)$; see Definition 1.15(ii) and Definition 5.1(i) for precise definitions. In particular, this implies that $\text{HK}_-(\phi_c, \phi_j)$ and $\text{HK}(\phi_c, \phi_j)$ hold for all $x, y \in M$ (not only for all $x, y \in M_0$).

(ii) Since
$$\phi(r) := \phi_c(r) \land \phi_j(r), \ \phi^{-1}(r) = \phi_c^{-1}(r) \lor \phi_j^{-1}(r)$$
 and so

$$\frac{1}{V(x,\phi^{-1}(t))} = \frac{1}{V(x,\phi_c^{-1}(t))} \wedge \frac{1}{V(x,\phi_j^{-1}(t))}.$$

It follows from (1.27) and (1.29) that under $HK_{-}(\phi_{c}, \phi_{j})$, for each fixed c > 0,

$$p(t, x, y) \simeq 1/V(x, \phi^{-1}(t))$$
 when $d(x, y) \le c\phi^{-1}(t)$.

In particular, $NL(\phi)$ holds under $HK_{-}(\phi_c, \phi_j)$, and so under $HK(\phi_c, \phi_j)$. The off-diagonal estimates of $HK(\phi_c, \phi_j)$ is expressed by the factor $p^{(c)}(c_k t, x, y) + p^{(j)}(t, x, y)$. In particular, $HK(\phi_c, \phi_j)$ is equivalent to

$$\begin{split} & p(t,x,y) \\ & \asymp \frac{1}{V(x,\phi^{-1}(t))} \wedge \left(\frac{t}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{1}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))} \right) \right) \\ & \asymp \frac{1}{V(x,\phi^{-1}(t))} \wedge \left(\frac{1}{V(x,d(x,y))} \left(\frac{t}{\phi_j(d(x,y))} + \exp\left(-\frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))} \right) \right) \right). \end{split}$$

(iii) We can express $HK(\phi_c, \phi_j)$ in the following way. For $0 < t \le 1$,

$$p(t,x,y) \asymp \begin{cases} \frac{1}{V(x,\phi_c^{-1}(t))}, & d(x,y) \le c_1\phi_c^{-1}(t), \\ \frac{t}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{1}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right), & d(x,y) \ge c_1\phi_c^{-1}(t); \end{cases}$$

for $t \geq 1$,

$$p(t, x, y) \simeq p^{(j)}(t, x, y) \simeq \begin{cases} \frac{1}{V(x, \phi_j^{-1}(t))}, & d(x, y) \le c_2 \phi_j^{-1}(t), \\ \frac{t}{V(x, d(x, y)) \phi_j(d(x, y))}, & d(x, y) \ge c_2 \phi_j^{-1}(t). \end{cases}$$

In particular, for $t \geq 1$, heat kernel estimates are dominated by the non-local part of Dirichlet form $(\mathcal{E}, \mathcal{F})$. Furthermore, for $0 < t \leq 1$, we have the following more explicit expression for $\mathrm{HK}(\phi_c, \phi_i)$:

$$p(t,x,y) \asymp \begin{cases} \frac{1}{V(x,\phi_c^{-1}(t))}, & d(x,y) \le c_1\phi_c^{-1}(t), \\ \frac{1}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right), & c_1\phi_c^{-1}(t) \le d(x,y) \le t_*, \\ \frac{t}{V(x,d(x,y))\phi_j(d(x,y))}, & d(x,y) \ge t_*, \end{cases}$$

where t_* satisfies that

$$c_{3}\phi_{c}^{-1}(t)\log^{(\beta_{1,\phi_{c}}-1)/\beta_{2,\phi_{c}}}(\phi_{c}^{-1}(t)/\phi_{j}^{-1}(t)) \leq t_{*} \leq c_{4}\phi_{c}^{-1}(t)\log^{(\beta_{2,\phi_{c}}-1)/\beta_{1,\phi_{c}}}(\phi_{c}^{-1}(t)/\phi_{j}^{-1}(t)),$$

and β_{1,ϕ_c} and β_{2,ϕ_c} are given in (1.10). See the proof of Proposition 4.6 for more details. Figure 1 indicates which term is the dominant one for the estimate of p(t, x, y) in each region.

- (iv) For any $D \subset M$, it holds for $x, y \in M_0$ and t > 0 that $p(t, x, y) \geq p^D(t, x, y)$, and so NDL(ϕ) is stronger than NL(ϕ). Furthermore, under VD and (1.10) we can prove that NL(ϕ) together with UHK(ϕ_c, ϕ_j) implies NDL(ϕ), see Lemma 5.7. We also note that, under VD, $V(x_0, \phi^{-1}(t))$ in the definition of NDL(ϕ) can be replaced by either $V(x, \phi^{-1}(t))$ or $V(y, \phi^{-1}(t))$. Under (1.10), we may also replace $\phi(\varepsilon r)$ and $\varepsilon \phi^{-1}(t)$ in the definition of NDL(ϕ) by $\varepsilon \phi(r)$ and $\phi^{-1}(\varepsilon t)$, respectively.
- (v) If in the lower bound for the definition of $HK_{-}(\phi_{c}, \phi_{j})$, we assume

$$p(t, x, y) \ge c_0 \left(\frac{1}{V(x, \phi_c^{-1}(t))} \land p^{(j)}(t, x, y)\right)$$

instead of (1.31), then we only have

$$p(t,x,y) \ge c_2 \left(\frac{1}{V(x,\phi_c^{-1}(t))} \land \frac{1}{V(x,\phi_j^{-1}(t))} \right) = \frac{c_2}{V(x,\phi^{-1}(t))} \quad \text{for } d(x,y) \le c_1 \phi_j^{-1}(t)$$

Note that as $\phi^{-1}(t) = \mathbf{1}_{[0,1]}(t)\phi_c^{-1}(t) + \mathbf{1}_{(1,\infty)}\phi_j^{-1}(t)$ and $\phi_c^{-1}(t) \ge \phi_j^{-1}(t)$ on [0,1], the above inequality is weaker than $\operatorname{NL}(\phi)$ (for instance when $\phi_c(r) = r^2$ and $\phi_j(r) = r^{\alpha}$ with $0 < \alpha < 2$).

We say $(\mathcal{E}, \mathcal{F})$ is conservative if its associated Hunt process X has infinite lifetime. This is equivalent to $P_t 1 = 1$ a.e. on M_0 for every t > 0. It follows from [CKW1, Proposition 3.1(ii)] that VD and NL(ϕ) imply that $(\mathcal{E}, \mathcal{F})$ is conservative.

Theorem 1.13. Assume that the metric measure space (M, d, μ) satisfies VD and RVD, and that the scale functions ϕ_c and ϕ_j satisfy (1.10) and (1.11). Let $\phi := \phi_c \wedge \phi_j$. The following are equivalent:

- (i) $\operatorname{HK}_{-}(\phi_c, \phi_j)$.
- (ii) UHK (ϕ_c, ϕ_j) , NL (ϕ) and J $_{\phi_i}$.
- (iii) UHKD(ϕ), NDL(ϕ) and J_{ϕ_i}.

(iv) $PI(\phi)$, J_{ϕ_i} and $Gcap(\phi)$.

(v) $PI(\phi)$, J_{ϕ_i} and $CS(\phi)$.

If, additionally, (M, d, μ) is connected and satisfies the chain condition, then all the conditions above are equivalent to:

(vi) $\text{HK}(\phi_c, \phi_j)$.

In the process of establishing Theorem 1.13, we also obtain the following characterizations for UHK (ϕ_c, ϕ_i) .

Theorem 1.14. Assume that the metric measure space (M, d, μ) satisfies VD and RVD, and that the scale functions ϕ_c and ϕ_j satisfy (1.10) and (1.11). Let $\phi := \phi_c \wedge \phi_j$. Then the following are equivalent:

- (i) UHK (ϕ_c, ϕ_i) and $(\mathcal{E}, \mathcal{F})$ is conservative.
- (ii) UHKD(ϕ), $J_{\phi_j,\leq}$ and E_{ϕ} .
- (iii) FK(ϕ), J_{ϕ_i , \leq} and Gcap(ϕ).
- (iv) FK(ϕ), J_{ϕ_i ,< and CS(ϕ).}

The proof of Theorem 1.14 is given at the end of Section 4, while the proof of Theorem 1.13 is given at the end of Section 5. We point out that $\text{UHK}(\phi_c, \phi_j)$ alone does not imply the conservativeness of the associated Dirichlet form $(\mathcal{E}, \mathcal{F})$. See [CKW1, Proposition 3.1 and Remark 3.2] for more details. Under VD, RVD and (1.10), NDL(ϕ) implies E_{ϕ} (see Proposition 4.1(ii) below), and so (iii) in Theorem 1.13 is stronger than (ii) in Theorem 1.14. We also note that RVD is only used in the implications of UHKD(ϕ) \Longrightarrow FK(ϕ) and PI(ϕ) \Longrightarrow FK(ϕ); see Proposition 3.1 below. In particular, (iii) \Longrightarrow (iv) \Longrightarrow (i) in Theorem 1.14 holds true under VD and (1.10). See Figure 2 below for various relations among UHK(ϕ_c, ϕ_j), HK₋(ϕ_c, ϕ_j) and HK(ϕ_c, ϕ_j).

We emphasize again that the connectedness and the chain condition of the underlying metric measure space (M, d, μ) are only used to derive optimal lower bounds off-diagonal estimates for heat kernel when the time is small (i.e., from $HK_{-}(\phi_c, \phi_j)$ to $HK(\phi_c, \phi_j)$), while for other statements in the two main results above, the metric measure space (M, d, μ) is only assumed to satisfy the general VD and RVD; that is, neither do we assume M to be connected nor (M, d)to be geodesic. Furthermore, we do not assume the uniform comparability of volume of balls; that is, we do not assume the existence of a non-decreasing function V on $[0, \infty)$ with V(0) = 0so that $\mu(B(x, r)) \simeq V(r)$ for all $x \in M$ and r > 0.

1.3 Parabolic Harnack inequalities

Let $Z := \{V_s, X_s\}_{s \ge 0}$ be the space-time process corresponding to X, where $V_s = V_0 - s$. The augmented filtration generated by Z satisfying the usual conditions will be denoted by $\{\widetilde{\mathcal{F}}_s; s \ge 0\}$. The law of the space-time process $s \mapsto Z_s$ starting from (t, x) will be denoted by $\mathbb{P}^{(t,x)}$. For every open subset D of $[0, \infty) \times M$, define $\tau_D = \inf\{s > 0 : Z_s \notin D\}$.

Definition 1.15. (i) We say that a Borel measurable function u(t, x) on $[0, \infty) \times M$ is parabolic (or caloric) on $D = (a, b) \times B(x_0, r)$ for the process X if there is a properly exceptional set \mathcal{N}_u associated with the process X so that for every relatively compact open subset U of D, $u(t, x) = \mathbb{E}^{(t,x)}u(Z_{\tau_U})$ for every $(t, x) \in U \cap ([0, \infty) \times (M \setminus \mathcal{N}_u))$.



Figure 2: Diagram for heat kernel estimates

(ii) We say that the parabolic Harnack inequality (PHI(ϕ)) holds for the process X, if there exist constants $0 < C_1 < C_2 < C_3 < C_4$, $C_5 > 1$ and $C_6 > 0$ such that for every $x_0 \in M$, $t_0 \geq 0$, R > 0 and for every non-negative function u = u(t, x) on $[0, \infty) \times M$ that is parabolic on cylinder $Q(t_0, x_0, \phi(C_4R), C_5R) := (t_0, t_0 + \phi(C_4R)) \times B(x_0, C_5R)$,

$$\operatorname{ess\,sup}_{Q_{-}} u \le C_6 \operatorname{ess\,inf}_{Q_{+}} u, \tag{1.33}$$

where
$$Q_{-} := (t_0 + \phi(C_1R), t_0 + \phi(C_2R)) \times B(x_0, R)$$
 and $Q_{+} := (t_0 + \phi(C_3R), t_0 + \phi(C_4R)) \times B(x_0, R)$.

The above $PHI(\phi)$ is called a weak parabolic Harnack inequality in [BGK], in the sense that (1.33) holds for some $C_1, \dots, C_5 > 0$. It is called a parabolic Harnack inequality in [BGK] if (1.33) holds for any choice of positive constants C_1, \dots, C_5 with $C_6 = C_6(C_1, \dots, C_5) < \infty$. Since our underlying metric measure space may not be geodesic, one can not expect to deduce parabolic Harnack inequalities from weak parabolic Harnack inequalities.

The following definition was initially introduced in [BBK2] in the setting of graphs. See [CKK2] for the general setting of metric measure spaces.

Definition 1.16. We say that UJS holds if there is a non-negative symmetric function J(x, y) on $M \times M$ so that for $\mu \times \mu$ almost all $x, y \in M$, (1.16) holds, and that there is a constant c > 0 such that for μ -a.e. $x, y \in M$ with $x \neq y$,

$$J(x,y) \le \frac{c}{V(x,r)} \int_{B(x,r)} J(z,y) \,\mu(dz) \quad \text{for every } 0 < r \le d(x,y)/2. \tag{1.34}$$

The following are the main results for parabolic Harnack inequalities. See Section 6 and Remark 4.9 for notations appeared in the statement.

Theorem 1.17. Suppose that the metric measure space (M, d, μ) satisfies VD and RVD, and that the scale functions ϕ_c and ϕ_j satisfy (1.10) and (1.11). Let $\phi := \phi_c \wedge \phi_j$. Then the following statements are equivalent.

(i) $PHI(\phi)$.

- (ii) $\text{UHK}_{weak}(\phi) + \text{NDL}(\phi) + \text{UJS}.$
- (iii) $PHR(\phi) + E_{\phi} + UJS + J_{\phi,\leq}$.
- (iv) $EHR + E_{\phi} + UJS + J_{\phi,\leq}$.
- (v) $\operatorname{PI}(\phi) + \operatorname{J}_{\phi,\leq} + \operatorname{Gcap}(\phi) + \operatorname{UJS}.$
- (vi) $PI(\phi) + J_{\phi,<} + CS(\phi) + UJS.$

Consequently, we have

$$\operatorname{HK}_{-}(\phi_{c},\phi_{j}) \Longleftrightarrow \operatorname{PHI}(\phi) + \operatorname{J}_{\phi_{j}}.$$
(1.35)

If in additional, the metric measure space (M, d, μ) is connected and satisfies the chain condition, then

$$\mathrm{HK}(\phi_c, \phi_j) \Longleftrightarrow \mathrm{PHI}(\phi) + \mathrm{J}_{\phi_j}.$$
 (1.36)

The equivalence between (i) and (ii) will be proved in Theorem 6.3, the equivalence between (i), (iii) and (iv) will be established in Theorem 6.4, while the equivalence between (i), (v) and (vi) will be given in Theorem 6.5. The last two assertions of Theorem 1.17 follow from the equivalence between (i), (v) and Theorem 1.13.

We emphasize that, different from the purely non-local setting as studied in [CKW2], PHI(ϕ) alone can only imply $J_{\phi,<}$ but not the stronger $J_{\phi_i,<}$. See Example 7.1 for a counterexample.

The rest of the paper is organized as follows. In the next section, we present some preliminary results about $\text{Gcap}(\phi)$. We show in Proposition 2.5 that $\text{Gcap}(\phi)$ along with $J_{\phi,\leq}$ yields $\text{CS}(\phi)$. This immediately yields (iv) \implies (v) in Theorem 1.13 and (iii) \implies (iv) in Theorem 1.14. Furthermore, $\text{CS}(\phi)$ enjoys the self-improving property, and enables us to make full use of the ideas in [CKW1, CKW2]. For example, via them we can obtain the L^1 -mean value inequalities in the present setting, which play a key tool to obtain E_{ϕ} . In Section 3, we investigate consequences of UHK(ϕ_c, ϕ_j), and establish (i) \implies (iii) of Theorem 1.14. Section 4 is devoted to obtaining UHK(ϕ_c, ϕ_j), which is the most difficult part of the paper. The crucial step is to apply rough tail probability estimates to derive sharp UHK(ϕ_c, ϕ_j), which requires detailed analysis of the roles of the local and non-local parts in different time and space regions. The proof of Theorem 1.14 is given at the end of Section 4. Section 5 is devoted to the two-sided heat kernel estimates and the proof of Theorem 1.13. Various characterizations of PHI(ϕ) are given in Section 6. In the last section, some examples are shown to illustrate the applications of our results, and a counterexample is also given to indicate that alone PHI(ϕ) does not imply $J_{\phi_i,\leq}$.

Throughout this paper, we will use c, with or without subscripts, to denote strictly positive finite constants whose values are insignificant and may change from line to line. For $p \in [1, \infty]$, we will use $||f||_p$ to denote the L^p -norm in $L^p(M; \mu)$. For $a, b \in \mathbb{R}_+$, $a \wedge b =: \min\{a, b\}$ and $a \vee b =: \max\{a, b\}$. For $B = B(x_0, r)$ and a > 0, we use aB to denote the ball $B(x_0, ar)$, and $\overline{B} := \{x \in M : d(x, x_0) \leq r\}$. For any subset D of M, D^c denotes its complement in M.

2 Preliminaries

In this section, we mainly present some preliminary results about $\text{Gcap}(\phi)$. For our later use to establish the characterizations of parabolic Harnack inequalities, we always assume that $J_{\phi,\leq}$ is

satisfied in this section. Since $\phi(r) \leq \phi_j(r)$ for all r > 0, $J_{\phi,\leq}$ is weaker than $J_{\phi_j,\leq}$, and so all the results in this section still hold true with $J_{\phi,\leq}$ replaced by $J_{\phi_j,\leq}$.

2.1 Properties of ϕ_c and ϕ_j

We recall the following statement from [CKW1].

Lemma 2.1. ([CKW1, Lemma 2.1]) Assume that VD and (1.10) hold. If $J_{\phi_j,\leq}$ (resp. $J_{\phi,\leq}$) holds, then there exists a constant $c_1 > 0$ such that

$$\int_{B(x,r)^c} J(x,y)\,\mu(dy) \le \frac{c_1}{\phi_j(r)} \quad \text{for every } x \in M \text{ and } r > 0.$$

$$\left(\text{resp. } \int_{B(x,r)^c} J(x,y)\,\mu(dy) \le \frac{c_1}{\phi(r)} \quad \text{for every } x \in M \text{ and } r > 0.\right)$$

The following lemma is concerned with the exponential function in estimates for $p^{(c)}(t, x, y)$.

Lemma 2.2. Under (1.10), for any $c_0 > 0$ there exists a constant $C := C(c_0) \ge 1$ so that for any t, r > 0,

$$\frac{r}{C\bar{\phi}_c^{-1}(t/r)} \le \sup_{s>0} \left\{ \frac{r}{s} - \frac{c_0 t}{\phi_c(s)} \right\} \le \frac{Cr}{\bar{\phi}_c^{-1}(t/r)}.$$

Proof. We fix $c_0 > 0$ throughout the proof. Let $c_2 \ge c_1 > 0$ be the positive constants in (1.12). For any $b > c_0/c_2 + 1$ and t, r > 0,

$$\begin{split} \sup_{s>0} \left\{ \frac{r}{s} - \frac{c_0 t}{\phi_c(s)} \right\} &\geq \frac{r}{\bar{\phi}_c^{-1}(bt/r)} - \frac{c_0 t}{\phi_c(\bar{\phi}_c^{-1}(bt/r))} \\ &\geq \frac{r}{\bar{\phi}_c^{-1}(bt/r)} - \frac{c_0 r}{c_2 b \bar{\phi}_c^{-1}(bt/r)} = \frac{(bc_2 - c_0) r}{bc_2 \bar{\phi}_c^{-1}(bt/r)} \\ &\geq \frac{1}{c_{2,\bar{\phi}_c} b^{1+1/(\beta_{1,\phi_c}-1)}} \frac{r}{\bar{\phi}_c^{-1}(t/r)}, \end{split}$$

where we used (1.12) in the second inequality and (1.13) in the last inequality. On the other hand, for any $0 < a \le 1$ and t, r > 0,

$$\begin{split} \sup_{s>0} \left\{ \frac{r}{s} - \frac{c_0 t}{\phi_c(s)} \right\} &\leq \sup_{0 < s \le a\bar{\phi}_c^{-1}(t/r)} \left\{ \frac{r}{s} - \frac{c_0 t}{\phi_c(s)} \right\} + \sup_{s>a\bar{\phi}_c^{-1}(t/r)} \left\{ \frac{r}{s} - \frac{c_0 t}{\phi_c(s)} \right\} \\ &\leq \sup_{0 < s \le a\bar{\phi}_c^{-1}(t/r)} \left\{ \frac{r}{s} - \frac{c_0 r\bar{\phi}_c(s/a)}{\phi_c(s)} \right\} + \frac{r}{a\bar{\phi}_c^{-1}(t/r)} \\ &\leq \sup_{0 < s \le a\bar{\phi}_c^{-1}(t/r)} \left\{ \frac{r}{s} \left(1 - \frac{ac_0 c_1 \phi_c(s/a)}{\phi_c(s)} \right) \right\} + \frac{r}{a\bar{\phi}_c^{-1}(t/r)} \\ &\leq \sup_{0 < s \le a\bar{\phi}_c^{-1}(t/r)} \left\{ \frac{r}{s} \left(1 - c_0 c_1 c_{1,\phi_c} a^{-\beta_{1,\phi_c}+1} \right) \right\} + \frac{r}{a\bar{\phi}_c^{-1}(t/r)} \end{split}$$

where we used (1.12) in the third inequality and (1.10) in the last inequality. Taking $a \in (0, 1]$ such that $a = a_0 := (1 + (1/(c_0c_1c_{1,\phi_c})))^{-1/(\beta_{1,\phi_c}-1)}$ in the inequality above, we find that

$$\sup_{s>0}\left\{\frac{r}{s}-\frac{c_0t}{\phi_c(s)}\right\} \le \frac{r}{a_0\bar{\phi}_c^{-1}(t/r)}.$$

The desired assertion follows from the above two conclusions.

Corollary 2.3. Under (1.10), the expressions (1.22) and (1.23) for $p^{(c)}(t, x, y)$ are equivalent in the sense that (1.26) holds.

Proof. This is a direct consequence of Lemma 2.2.

2.2
$$\operatorname{EP}_{\phi,\leq,\varepsilon} \Longrightarrow \operatorname{Gcap}(\phi)$$
 and $\operatorname{Gcap}(\phi) + J_{\phi,\leq} \Longrightarrow \operatorname{CS}(\phi)$.

Recall that $\Gamma(f, f)$ (resp. $\Gamma_c(f, f)$) is the *energy measure* of f for \mathcal{E} (resp. its strongly local part $\mathcal{E}^{(c)}$). For any $f, g \in \mathcal{F}$, the signed measure $\Gamma(f, g)$ is defined by

$$\Gamma(f,g) = \frac{1}{2} \big(\Gamma(f+g,f+g) - \Gamma(f,f) - \Gamma(g,g) \big).$$

Similar definition applies to $\Gamma_c(f,g)$. The measure $\Gamma(f,g)$ is symmetric and bilinear in (f,g). The following Cauchy-Schwarz inequality holds:

$$\left| \int_{M} fg \, d\Gamma(u, v) \right| \leq \left(\int_{M} f^{2} \, d\Gamma(u, u) \right)^{1/2} \left(\int_{M} g^{2} \, d\Gamma(v, v) \right)^{2}$$
$$\leq \frac{1}{2\lambda} \int_{M} f^{2} \, d\Gamma(u, u) + \frac{\lambda}{2} \int_{M} g^{2} \, d\Gamma(v, v)$$
(2.1)

for all $u, v \in \mathcal{F}$, bounded f, g on M, and $\lambda > 0$. When the Dirichlet form $(\mathcal{E}, \mathcal{F})$ admits no killings as the one given by (1.1),

$$\mathcal{E}(f,g) = \Gamma(f,g)(M) \text{ for } f,g \in \mathcal{F}.$$

The following Leibniz and chain rules hold for the energy measure Γ_c for the strongly local part $\mathcal{E}^{(c)}$ of the Dirichlet form $(\mathcal{E}, \mathcal{F})$: for all $f, g, h \in \mathcal{F}_b$,

$$\Gamma_c(fg,h)(dx) = g(x)\Gamma_c(f,h)(dx) + f(x)\Gamma_c(g,h)(dx)$$

and

$$\Gamma_c(\Phi(f),g)(dx) = \Phi'(f)(x)\Gamma_c(f,g)(dx),$$

where $\Phi : \mathbb{R} \to \mathbb{R}$ is any smooth function with $\Phi(0) = 0$. The measure Γ_c has the strong local property in the sense that if f is constant on a set F, then

$$\Gamma_c(f, f)(F) = 0. \tag{2.2}$$

see [CF, Theorem 4.3.8 and Exercise 4.3.12]. For $f, g \in \mathcal{F}$, define

It is easy to check that the following chain rule holds:

$$\int_{M} d\Gamma_{j}(fg,h) = \int_{M} f \, d\Gamma_{j}(g,h) + \int_{M} g \, d\Gamma_{j}(f,h) \quad \text{for } f, g, h \in \mathcal{F}_{b}.$$

See [CKS, Lemma 3.5 and Theorem 3.7] for more details.

The following proposition extends [GHH, Lemma 2.8] from strongly local Dirichet forms on metric measure spaces to symmetric Dirichlet forms having both local and nonlocal terms on general metric measure spaces.

Proposition 2.4. Suppose that (1.10) holds. Then $EP_{\phi,\leq,\varepsilon} \Longrightarrow Gcap(\phi)$. Consequently,

$$\mathrm{EP}_{\phi,\leq} \Longrightarrow \mathrm{EP}_{\phi,\leq,\varepsilon} \Longrightarrow \mathrm{Gcap}(\phi),$$

and

$$E_{\phi} \Longrightarrow EP_{\phi,\leq,\varepsilon} \Longrightarrow Gcap(\phi).$$

Proof. The proof of $EP_{\phi,\leq,\varepsilon} \Longrightarrow Gcap(\phi)$ is the same as that of [GHH, Lemma 2.8]. We note that while [GHH, Leamm 2.8] concerns with purely non-local Dirichlet forms, its proof does not use any character of pure jump Dirichlet forms and works for general symmetric Dirichlet forms. Clearly, $EP_{\phi,\leq}$ implies $EP_{\phi,\leq,\varepsilon}$. By the same proof as that of [CKW1, Lemma 4.16], we have $E_{\phi} \Longrightarrow EP_{\phi,\leq,\varepsilon}$. This establishes the last assertion.

The next proposition extends [GHH, Lemma 2.4] from strongly local Dirichet forms on metric measure spaces satisfying the *d*-set upper bound condition to symmetric Dirichlet forms having both local and nonlocal terms on general metric measure spaces with VD condition. It in particular gives the implication (iv) \implies (v) in Theorem 1.13 and (iii) \implies (iv) in Theorem 1.14.

Proposition 2.5. Under VD and (1.10),

$$\operatorname{Gcap}(\phi) \text{ and } J_{\phi,\leq} \Longrightarrow \operatorname{CS}(\phi).$$

To prove Proposition 2.5, we need the following lemma.

Lemma 2.6. (i) For each $f, g \in \mathcal{F}_b$ and $\eta > 0$,

$$(1-\eta^{-1})\int f^2 d\Gamma_c(g,g) \leq \int d\Gamma_c(g,gf^2) + \eta \int g^2 d\Gamma_c(f,f).$$

(ii) For each $f, g \in \mathcal{F}_b$, $\eta > 0$ and any subset $D \subset M$,

$$(1 - \eta^{-1}) \int_{D \times D} f^{2}(x)(g(x) - g(y))^{2} J(dx, dy)$$

$$\leq \int_{D \times D} (g(x)f^{2}(x) - g(y)f^{2}(y))(g(x) - g(y)) J(dx, dy)$$

$$+ \eta \int_{D \times D} g^{2}(x)(f(x) - f(y))^{2} J(dx, dy).$$
(2.3)

Proof. (2.3) has been proved in [CKW1, Lemma 3.5], and so we only need to show (i). For any $f, g \in \mathcal{F}_b$ and $\eta > 0$, by Leibniz and chain rules and the Cauchy-Schwarz inequality (2.1),

$$\int d\Gamma_c(g, f^2g) = 2 \int gf \, d\Gamma_c(f, g) + \int f^2 \, d\Gamma_c(g, g)$$
$$\geq -\eta^{-1} \int f^2 \, d\Gamma_c(g, g) - \eta \int g^2 \, d\Gamma_c(f, f) + \int f^2 \, d\Gamma_c(g, g),$$

proving the assertion (i).

Proof of Proposition 2.5. For fixed $x_0 \in M$, $0 < r \leq R$ and $C_0 \in (0, 1]$, set $B_0 = B(x_0, R)$, $B_1 = B(x_0, R + r/2)$, $B_2 = B(x_0, R + r)$ and $B_3 = B(x_0, R + (1 + C_0)r)$. For any $f \in \mathcal{F}$, by $\operatorname{Gcap}(\phi)$ there is a κ -cut-off function φ for $B_0 \subset B_1$ such that

$$\mathcal{E}(f^2\varphi,\varphi) \le \frac{C_1}{\phi(r)} \int_{B_1} f^2 \, d\mu.$$
(2.4)

Since $\varphi = 0$ on B_1^c , we have

$$\begin{split} \mathcal{E}(f^{2}\varphi,\varphi) &= \mathcal{E}^{(c)}(f^{2}\varphi,\varphi) + \mathcal{E}^{(j)}(f^{2}\varphi,\varphi) \\ &= \mathcal{E}^{(c)}(f^{2}\varphi,\varphi) \\ &+ \left(\int_{B_{2}\times B_{2}} + \int_{B_{2}^{c}\times B_{2}} + \int_{B_{2}\times B_{2}^{c}}\right)(\varphi(x) - \varphi(y))(f^{2}(x)\varphi(x) - f^{2}(y)\varphi(y)) J(dx,dy) \\ &\geq \int_{B_{2}} d\Gamma_{c}(f^{2}\varphi,\varphi) + \int_{B_{2}\times B_{2}}(\varphi(x) - \varphi(y))(f^{2}(x)\varphi(x) - f^{2}(y)\varphi(y)) J(dx,dy). \end{split}$$

This along with Lemma 2.6 with $\eta = 2$ and $\text{Gcap}(\phi)$ yields that

$$\begin{split} &\int_{B_2} f^2 \, d\Gamma_c(\varphi,\varphi) + \int_{B_2 \times B_2} f^2(x)(\varphi(x) - \varphi(y))^2 \, J(dx,dy) \\ &\leq 2 \left(\int_{B_2} d\Gamma_c(f^2\varphi,\varphi) + \int_{B_2 \times B_2} (\varphi(x) - \varphi(y))(f^2(x)\varphi(x) - f^2(y)\varphi(y)) \, J(dx,dy) \right) \\ &\quad + 4 \left(\int_{B_2} \varphi^2 \, d\Gamma_c(f,f) + \int_{B_2 \times B_2} \varphi^2(x)(f(x) - f(y))^2 \, J(dx,dy) \right) \\ &\leq 2 \mathcal{E}(f^2\varphi,\varphi) + 4 \left(\int_{B_2} \varphi^2 \, d\Gamma_c(f,f) + \int_{B_2 \times B_2} \varphi^2(x)(f(x) - f(y))^2 \, J(dx,dy) \right) \\ &\leq \frac{2C_1}{\phi(r)} \int_{B_1} f^2 \, d\mu + 4 \left(\int_{B_2} \varphi^2 \, d\Gamma_c(f,f) + \int_{B_2 \times B_2} \varphi^2(x)(f(x) - f(y))^2 \, J(dx,dy) \right). \end{split}$$

Replacing [GHH, (2.4) on page 447] with the inequality above, and following the proof of [GHH, Lemma 2.4] from [GHH, (2.4) on page 447] to the end, we can obtain that

$$\begin{split} &\int_{B_3} f^2 \, d\Gamma_c(\varphi,\varphi) + \int_{B_3 \times B_3} f^2(x)(\varphi(x) - \varphi(y))^2 \, J(dx,dy) \\ &= \int_{B_2} f^2 \, d\Gamma_c(\varphi,\varphi) + \int_{B_3 \times B_3} f^2(x)(\varphi(x) - \varphi(y))^2 \, J(dx,dy) \\ &\leq \frac{C_2}{\phi(r)} \int_{B_3} f^2 \, d\mu + 4 \left(\int_{B_2} \varphi^2 \, d\Gamma_c(f,f) + \int_{B_2 \times B_2} \varphi^2(x)(f(x) - f(y))^2 \, J(dx,dy) \right). \end{split}$$

(Indeed, this can be seen by replacing B and Ω in the proof of [GHH, Lemma 2.4] with B_2 and B_3 , respectively.) We note that for the argument above, we used Lemma 2.1. Again by Lemma 2.1, we can further improve the inequality above into

$$\begin{split} \int_{B_3} f^2 \, d\Gamma(\varphi,\varphi) &= \int_{B_3} f^2 \, d\Gamma_c(\varphi,\varphi) + \int_{B_3 \times B_3} f^2(x)(\varphi(x) - \varphi(y))^2 \, J(dx,dy) \\ &+ \int_{B_3 \times B_3^c} f^2(x)(\varphi(x) - \varphi(y))^2 \, J(dx,dy) \end{split}$$

$$\leq \int_{B_3} f^2 d\Gamma_c(\varphi,\varphi) + \int_{B_3 \times B_3} f^2(x)(\varphi(x) - \varphi(y))^2 J(dx,dy) + \int_{B_1 \times B_3^c} f^2(x) J(dx,dy) \leq 4 \left(\int_{B_2} \varphi^2 d\Gamma_c(f,f) + \int_{B_2 \times B_2} \varphi^2(x)(f(x) - f(y))^2 J(dx,dy) \right) + \frac{C_3}{\phi(r)} \int_{B_3} f^2 d\mu$$
desired assertion. \Box

This proves the desired assertion.

2.3 ρ -truncated version of $CS(\phi)$

To deal with processes with long rang jumps, we will frequently use the truncation. Fix $\rho > 0$ and define a bilinear form $(\mathcal{E}^{(\rho)}, \mathcal{F})$ by

$$\mathcal{E}^{(\rho)}(u,v) = \mathcal{E}^{(c)}(u,v) + \int_{M \times M} (u(x) - u(y))(v(x) - v(y)) \mathbf{1}_{\{d(x,y) \le \rho\}} J(dx,dy)$$

= : $\mathcal{E}^{(c)}(u,v) + \mathcal{E}^{(\rho)}_{j}(u,v).$

Clearly, the form $\mathcal{E}^{(\rho)}(u, v)$ is well defined for $u, v \in \mathcal{F}$, and $\mathcal{E}^{(\rho)}(u, u) \leq \mathcal{E}(u, u)$ for all $u \in \mathcal{F}$. Assume that VD, (1.10) and $J_{\phi,\leq}$ hold. Then we have by Lemma 2.1 that for all $u \in \mathcal{F}$,

$$\begin{split} \mathcal{E}(u,u) - \mathcal{E}^{(\rho)}(u,u) &= \int_{M \times M} (u(x) - u(y))^2 \mathbf{1}_{\{d(x,y) > \rho\}} J(dx,dy) \\ &\leq 4 \int_M u^2(x) \, \mu(dx) \int_{B(x,\rho)^c} J(x,y) \, \mu(dy) \leq \frac{c_0 ||u||_2^2}{\phi(\rho)} \end{split}$$

Thus $\mathcal{E}_1(u, u)$ is equivalent to $\mathcal{E}_1^{(\rho)}(u, u) := \mathcal{E}^{(\rho)}(u, u) + ||u||_2^2$ for every $u \in \mathcal{F}$. Hence $(\mathcal{E}^{(\rho)}, \mathcal{F})$ is a regular Dirichlet form on $L^2(M; \mu)$. Throughout this paper, we call $(\mathcal{E}^{(\rho)}, \mathcal{F})$ ρ -truncated Dirichlet form. The Hunt process associated with $(\mathcal{E}^{(\rho)}, \mathcal{F})$, denoted by $(X_t^{(\rho)})_{t\geq 0}$, can be identified in distribution with the Hunt process of the original Dirichlet form $(\mathcal{E}, \mathcal{F})$ by removing those jumps of size larger than ρ .

Define $J(x, dy) = J(x, y) \mu(dy)$. Let $J^{(\rho)}(dx, dy) = \mathbf{1}_{\{d(x,y) \le \rho\}} J(dx, dy), J^{(\rho)}(x, dy) = \mathbf{1}_{\{d(x,y) \le \rho\}} J(x, dy)$, and $\Gamma_j^{(\rho)}(f, g)$ be the carré du champ of the non-local part $\mathcal{E}_j^{(\rho)}$ for the ρ -truncated Dirichlet form $(\mathcal{E}^{(\rho)}, \mathcal{F})$; namely,

$$\mathcal{E}_{j}^{(\rho)}(f,g) = \int_{M} \mu(dx) \int_{M} (f(x) - f(y))(g(x) - g(y)) J^{(\rho)}(x,dy) =: \int_{M} d\Gamma_{j}^{(\rho)}(f,g)$$

We also set

$$\Gamma^{(\rho)}(f,g) := \Gamma_c(f,g) + \Gamma_j^{(\rho)}(f,g).$$

Lemma 2.7. Under VD, (1.10) and $J_{\phi,\leq}$, if $CS(\phi)$ holds, then $CS^{(\rho)}(\phi)$ holds too, i.e., there exist constants $C_0 \in (0,1]$ and $C_1, C_2 > 0$ such that for every $0 < r \leq R$, almost all $x_0 \in M$ and any $f \in \mathcal{F}$, there exists a cut-off function $\varphi \in \mathcal{F}_b$ for $B(x_0, R) \subset B(x_0, R+r)$ so that the following holds for all $\rho \in (0,\infty]$:

$$\int_{B(x_0,R+(1+C_0)r)} f^2 \, d\Gamma^{(\rho)}(\varphi,\varphi)$$

$$\leq C_1 \left(\int_{B(x_0, R+r)} \varphi^2 \, d\Gamma_c(f, f) + \int_{B(x_0, R+r) \times B(x_0, R+(1+C_0)r)} \varphi^2(x) (f(x) - f(y))^2 \, J^{(\rho)}(dx, dy) \right) \\ + \frac{C_2}{\phi(r \wedge \rho)} \int_{B(x_0, R+(1+C_0)r)} f^2 \, d\mu.$$

Proof. The proof is based on Lemma 2.1, and we omit it here. See [CKW1, Proposition 2.3(1)].

We also note that, by the proof of [CKW1, Proposition 2.3(4)], under VD, (1.10) and $J_{\phi,\leq}$, if $CS(\phi)$ (resp. $CS^{(\rho)}(\phi)$) holds for some $C_0 \in (0, 1]$, then for any $C'_0 \in (C_0, 1]$, there exist constants $C_1, C_2 > 0$ (where C_2 depends on C'_0) such that $CS(\phi)$ (resp. $CS^{(\rho)}(\phi)$) holds for C'_0 .

Proposition 2.8. Assume that VD, (1.10) and $J_{\phi,\leq}$ hold. If $CS(\phi)$ holds, then there is a constant $c_0 > 0$ such that for every $0 < r \leq R$, $\rho > 0$ and almost all $x \in M$,

$$\operatorname{Cap}^{(\rho)}(B(x,R),B(x,R+r)) \le c_0 \frac{V(x,R+r)}{\phi(r \wedge \rho)}.$$

In particular, we have

$$\operatorname{Cap}(B(x,R), B(x,R+r)) \le c_0 \frac{V(x,R+r)}{\phi(r)}.$$
(2.5)

Proof. According to Lemma 2.7 and the remark below its proof, $CS^{(\rho)}(\phi)$ holds for every $\rho > 0$ and we may and do take $C_0 = 1$ in (1.18). Fix $x_0 \in M$ and write $B_s := B(x_0, s)$ for $s \ge 0$. Let $f \in \mathcal{F}$ with $0 \le f \le 1$ such that $f|_{B_{R+2r}} = 1$ and $f|_{B_{R+3r}} = 0$. For any $0 < r \le R$, let $\varphi \in \mathcal{F}_b$ be the cut-off function for $B_R \subset B_{R+r}$ associated with f in $CS^{(\rho)}(\phi)$. Then for any $\rho > 0$,

$$\begin{aligned} \operatorname{Cap}^{(\rho)}(B_{R}, B_{R+r}) &\leq \int_{B_{R+2r}} d\Gamma^{(\rho)}(\varphi, \varphi) + \int_{B_{R+2r}^{c}} d\Gamma^{(\rho)}(\varphi, \varphi) \\ &= \int_{B_{R+2r}} f^{2} d\Gamma^{(\rho)}(\varphi, \varphi) + \int_{B_{R+2r}^{c}} d\Gamma^{(\rho)}(\varphi, \varphi) \\ &\leq c_{1} \left(\int_{B_{R+r}} d\Gamma_{c}(f, f) + \int_{B_{R+r} \times B_{R+2r}} (f(x) - f(y))^{2} J^{(\rho)}(dx, dy) \right) \\ &\quad + \frac{c_{2}}{\phi(r \wedge \rho)} \int_{B_{R+2r}} f^{2} d\mu + \int_{B_{R+2r}^{c}} \mu(dx) \int_{B_{R+r}} \varphi^{2}(y) J(x, y) \, \mu(dy) \\ &\leq 0 + 0 + \frac{c_{2}\mu(B_{R+2r})}{\phi(r \wedge \rho)} + \frac{c_{3}\mu(B_{R+r})}{\phi(r)} \\ &\leq \frac{c_{4}\mu(B_{R+r})}{\phi(r \wedge \rho)}, \end{aligned}$$

where we used $CS^{(\rho)}(\phi)$ in the second inequality, and applied Lemma 2.1 and VD in the third inequality.

Now let f_{ρ} be the potential whose $\mathcal{E}^{(\rho)}$ -norm gives the capacity. Then the Cesàro mean of a subsequence of f_{ρ} converges in \mathcal{E}_1 -norm, say to f, and $\mathcal{E}(f, f)$ is no less than the capacity corresponding to $\rho = \infty$. So (2.5) is proved.

2.4 Self-improvement of $CS(\phi)$

We next show that the leading constant in $CS^{(\rho)}(\phi)$ is self-improving in the following sense.

Proposition 2.9. Suppose that VD, (1.10) and $J_{\phi,\leq}$ hold. If $CS^{(\rho)}(\phi)$ holds, then there exists a constant $C_0 \in (0,1]$ so that for any $\varepsilon > 0$, there exists a constant $c_1(\varepsilon) > 0$ such that for every $0 < r \leq R$, almost all $x_0 \in M$ and any $f \in \mathcal{F}$, there exists a cut-off function $\varphi \in \mathcal{F}_b$ for $B(x_0, R) \subset B(x_0, R+r)$ so that the following holds for all $\rho > 0$:

$$\int_{B(x_0,R+(1+C_0)r)} f^2 d\Gamma^{(\rho)}(\varphi,\varphi)
\leq \varepsilon \left(\int_{B(x_0,R+r)} \varphi^2 d\Gamma_c(f,f)
+ \int_{B(x_0,R+r)\times B(x_0,R+(1+C_0)r)} \varphi^2(x)(f(x) - f(y))^2 J^{(\rho)}(dx,dy) \right)
+ \frac{c_1(\varepsilon)}{\phi(r \wedge \rho)} \int_{B(x_0,R+(1+C_0)r)} f^2 d\mu.$$
(2.6)

Proof. By the remark before Proposition 2.8, we may and do assume that $CS^{(\rho)}(\phi)$ holds with $C_0 = 1$. Fix $x_0 \in M$, $0 < r \leq R$ and $f \in \mathcal{F}$. For s > 0, set $B_s = B(x_0, s)$. The goal is to construct a cut-off function $\varphi \in \mathcal{F}_b$ for $B_R \subset B_{R+r}$ so that (2.6) holds. Without loss of generality, in the following we may and do assume that $\int_{B_{R+2r}} f^2 d\mu > 0$; otherwise, (2.6) holds trivially.

For $\lambda > 0$ whose exact value to be determined later, let

$$s_n = c_0 r e^{-n\lambda/(2\beta_{2,\phi})},$$

where $c_0 := c_0(\lambda)$ is chosen so that $\sum_{n=1}^{\infty} s_n = r$, and $\beta_{2,\phi} := \beta_{2,\phi_c} \vee \beta_{2,\phi_j}$ is given in (1.15). Set $r_0 = 0$ and

$$r_n = \sum_{k=1}^n s_k, \quad n \ge 1.$$

Clearly, $R < R + r_1 < R + r_2 < \cdots < R + r$. For any $n \ge 0$, define $U_n := B_{R+r_{n+1}} \setminus B_{R+r_n}$, and $U_n^* := B_{R+r_{n+1}+s_{n+1}} \setminus B_{R+r_n-s_{n+1}}$. Let $\theta > 0$, whose value also to be determined later, and define $f_{\theta} := |f| + \theta$. By $\operatorname{CS}^{(\rho)}(\phi)$ (with $R = R + r_n$, $r = r_{n+1} - r_n = s_{n+1}$), there exists a cut-off function φ_n for $B_{R+r_n} \subset B_{R+r_{n+1}}$ such that

$$\begin{split} &\int_{B_{R+r_{n+1}+s_{n+1}}} f_{\theta}^2 \, d\Gamma^{(\rho)}(\varphi_n,\varphi_n) \\ &\leq C_1 \left(\int_{B_{R+r_{n+1}}} \varphi_n^2 \, d\Gamma_c(f_{\theta},f_{\theta}) + \int_{B_{R+r_{n+1}}\times B_{R+r_{n+1}+s_{n+1}}} \varphi_n^2(x) (f_{\theta}(x) - f_{\theta}(y))^2 \, J^{(\rho)}(dx,dy) \right) \\ &\quad + \frac{C_2}{\phi(s_{n+1}\wedge\rho)} \int_{B_{R+r_{n+1}+s_{n+1}}} f_{\theta}^2 \, d\mu, \end{split}$$

where C_1, C_2 are positive constants independent of f_{θ} and φ_n . Here, we mention that since $(\mathcal{E}, \mathcal{F})$ is a regular Dirichlet form on $L^2(M, \mu)$, $f_{\theta} \in \mathcal{F}_{loc}$, and so, by Remark 1.7(ii), $CS^{(\rho)}(\phi)$ can be applied to f_{θ} .

Let $b_n = e^{-n\lambda}$ and define

$$\varphi = \sum_{n=1}^{\infty} (b_{n-1} - b_n)\varphi_n.$$
(2.7)

Then φ is a cut-off function for $B_R \subset B_{R+r}$, because $\varphi = 1$ on B_R and $\varphi = 0$ on B_{R+r}^c . Hence, combining the proof of [AB, Lemma 5.1] with that of [CKW1, Proposition 2.4], we can verify that the function φ defined by (2.7) satisfies (2.6) with $C_0 = 1$ and $\varphi \in \mathcal{F}_b$. (In particular, by [AB, (5.7) in the proof of Lemma 5.1], we can insert the function φ^2 in front of $d\Gamma_c(f, f)$.) The details are omitted here.

As a direct consequence of Lemma 2.7 and Proposition 2.9, we have the following corollary.

Corollary 2.10. Suppose that VD, (1.10), $J_{\phi,\leq}$ and $CS(\phi)$ hold. Then there exists a constant $c_1 > 0$ such that for every $0 < r \leq R$, almost all $x_0 \in M$ and any $f \in \mathcal{F}$, there exists a cut-off function $\varphi \in \mathcal{F}_b$ for $B(x_0, R) \subset B(x_0, R+r)$ so that the following holds for all $\rho \in (0, \infty]$,

$$\int_{B(x_0,R+2r)} f^2 d\Gamma^{(\rho)}(\varphi,\varphi) \\
\leq \frac{1}{8} \left(\int_{B(x_0,R+r)} \varphi^2 d\Gamma_c(f,f) + \int_{B(x_0,R+r) \times B(x_0,R+2r)} \varphi^2(x) (f(x) - f(y))^2 J^{(\rho)}(dx,dy) \right) \quad (2.8) \\
+ \frac{c_1}{\phi(r \wedge \rho)} \int_{B(x_0,R+2r)} f^2 d\mu.$$

2.5 Consequences of $CS(\phi)$: Caccioppoli and L^1 -mean value inequalities

In this subsection, we establish mean value inequalities for subharmonic functions. For stability results for heat kernel estimates, we only need mean value inequalities for the ρ -truncated Dirichlet form $(\mathcal{E}^{(\rho)}, \mathcal{F})$, while the mean value inequalities for the original Dirichlet form $(\mathcal{E}, \mathcal{F})$ will be used as one of the key tools in the study of characterizations of parabolic Harnack inequalities. We will first present these inequalities for subharmonic functions of the original Dirichlet form $(\mathcal{E}, \mathcal{F})$ and then indicate similar inequalities for subharmonic functions of the ρ -truncated Dirichlet form $(\mathcal{E}^{(\rho)}, \mathcal{F})$.

Definition 2.11. Let D be an open subset of M.

(i) We say that a bounded nearly Borel measurable function u on M is \mathcal{E} -subharmonic (resp. \mathcal{E} -harmonic, \mathcal{E} -superharmonic) in D if $u \in \mathcal{F}_{loc}^D$ satisfies

$$\mathcal{E}(u,\varphi) \le 0 \quad (\text{resp.} = 0, \ge 0)$$

for any $0 \leq \varphi \in \mathcal{F}^D$.

(ii) A nearly Borel measurable function u on M is said to be subharmonic (resp. harmonic, superharmonic) in D (with respect to the process X) if for any relatively compact subset $U \subset D$, $t \mapsto u(X_{t \wedge \tau_U})$ is a uniformly integrable submartingale (resp. martingale, supermartingale) under \mathbb{P}^x for q.e. $x \in M$.

The following result is established in [C, Theorem 2.11 and Lemma 2.3] first for harmonic functions, and then extended in [ChK, Theorem 2.9] to subharmonic functions.

Theorem 2.12. Let D be an open subset of M, and let u be a bounded function. Then u is \mathcal{E} -harmonic (resp. \mathcal{E} -subharmonic) in D if and only if u is harmonic (resp. subharmonic) in D.

To establish the Caccioppoli inequality, we also need the following definition.

Definition 2.13. Let $\psi : \mathbb{R}_+ \to \mathbb{R}_+$. For a Borel measurable function u on M, we define its *nonlocal tail* in the ball $B(x_0, r)$ with respect to the function ψ by

$$\operatorname{Tail}_{\psi}(u; x_0, r) = \psi(r) \int_{B(x_0, r)^c} \frac{|u(z)|}{V(x_0, d(x_0, z))\psi(d(x_0, z))} \,\mu(dz).$$
(2.9)

Suppose that VD and (1.10) hold. By Lemma 2.1, both $\operatorname{Tail}_{\phi_j}(u; x_0, r)$ and $\operatorname{Tail}_{\phi}(u; x_0, r)$ are finite if u is bounded.

We first show that $CS(\phi)$ enables us to prove a Caccioppoli inequality for \mathcal{E} -subharmonic functions.

Lemma 2.14. (Caccioppoli inequality) Suppose that VD, (1.10), $CS(\phi)$ and $J_{\phi,\leq}$ hold. For $x_0 \in M$ and s > 0, let $B_s = B(x_0, s)$. For 0 < r < R, let u be an \mathcal{E} -subharmonic function on B_{R+r} for the Dirichlet form $(\mathcal{E}, \mathcal{F})$, and $v := (u - \theta)^+$ for $\theta > 0$. Let φ be the cut-off function for $B_{R-r} \subset B_R$ associated with v in $CS(\phi)$. Then there exists a constant c > 0 independent of x_0, R, r and θ such that

$$\int_{B_{R+r}} d\Gamma(v\varphi, v\varphi) \le \frac{c}{\phi(r)} \left[1 + \frac{1}{\theta} \left(1 + \frac{R}{r} \right)^{d_2 + \beta_{2,\phi} - \beta_{1,\phi}} \operatorname{Tail}_{\phi}(u; x_0, R+r) \right] \int_{B_{R+r}} u^2 \, d\mu, \quad (2.10)$$

where $\beta_{1,\phi}$ and $\beta_{2,\phi}$ are given in (1.15).

Proof. (i) Since u is \mathcal{E} -subharmonic in B_{R+r} for the Dirichlet form $(\mathcal{E}, \mathcal{F})$ and $\varphi^2 v \in \mathcal{F}^{B_R}$, we have $u \in \mathcal{F}^{loc}_{B_{R+r}}$ and $\mathcal{E}(u, \varphi^2 v) = \mathcal{E}^{(c)}(u, \varphi^2 v) + \mathcal{E}^{(j)}(u, \varphi^2 v) \leq 0$.

As $u - v = u \mathbb{1}_{\{u \le \theta\}} + \theta \mathbb{1}_{\{u > \theta\}}$ and v = 0 on $\{u \le \theta\}$, we have by (2.2) that $\Gamma_c(u - v, v) = 0$ on M. Hence by the Leibniz and chain rules as well as the Cauchy-Schwarz inequality (2.1),

$$\begin{split} \mathcal{E}^{(c)}(u,\varphi^2 v) &= \int_M \varphi^2 \, d\Gamma_c(u,v) + 2 \int_M \varphi v \, d\Gamma_c(u,\varphi) \\ &= \int_M \varphi^2 \, d\Gamma_c(v,v) + 2 \int_M \varphi v \, d\Gamma_c(v,\varphi) \\ &\geq \int_M \varphi^2 \, d\Gamma_c(v,v) - \left(4 \int_M v^2 \, d\Gamma_c(\varphi,\varphi) + \frac{1}{4} \int_M \varphi^2 \, d\Gamma_c(v,v)\right) \\ &\geq \frac{3}{4} \int_M \varphi^2 \, d\Gamma_c(v,v) - 4 \int_M v^2 \, d\Gamma_c(\varphi,\varphi) \\ &= \frac{3}{4} \int_{B_R} \varphi^2 \, d\Gamma_c(v,v) - 4 \int_{B_R} v^2 \, d\Gamma_c(\varphi,\varphi). \end{split}$$

On the other hand, by [CKW1, (4.5)], we have

$$\mathcal{E}^{(j)}(u,\varphi^2 v) \ge \int_{B_R \times B_{R+r}} \varphi^2(x)(v(x) - v(y))^2 J(dx, dy) - 4 \int_{B_{R+r}} v^2 d\Gamma_j(\varphi, \varphi)$$

$$-\frac{c_1}{\theta\phi(r)}\left[\left(1+\frac{R}{r}\right)^{d_2+\beta_{2,\phi}-\beta_{1,\phi}}\operatorname{Tail}_{\phi}\left(u;x_0,R+r\right)\right]\int_{B_R}u^2\,d\mu.$$

Combining all the estimates above, we arrive at

$$0 \leq 4 \int_{B_{R+r}} v^2 d\Gamma(\varphi, \varphi) - \frac{3}{4} \Big(\int_{B_R} \varphi^2 d\Gamma_c(v, v) + \int_{B_R \times B_{R+r}} \varphi^2(x) (v(x) - v(y))^2 J(dx, dy) \Big)$$
$$+ \frac{c_1}{\theta \phi(r)} \Big[\left(1 + \frac{R}{r} \right)^{d_2 + \beta_{2,\phi} - \beta_{1,\phi}} \operatorname{Tail}_{\phi} (u; x_0, R+r) \Big] \int_{B_R} u^2 d\mu.$$
(2.11)

(ii) It is easy to see from the Leibniz rule and the Cauchy-Schwarz inequality (2.1) that

$$\int_{B_{R+r}} d\Gamma_c(v\varphi, v\varphi) \le 2 \int_{B_{R+r}} v^2 \, d\Gamma_c(\varphi, \varphi) + 2 \int_{B_{R+r}} \varphi^2 \, d\Gamma_c(v, v).$$

According to [CKW1, (4.6)],

$$\begin{split} \int_{B_{R+r}} d\Gamma_j(v\varphi, v\varphi) &\leq 2 \int_{B_{R+r}} v^2 \, d\Gamma_j(\varphi, \varphi) + 2 \int_{B_R \times B_{R+r}} \varphi^2(x) (v(x) - v(y))^2 \, J(dx, dy) \\ &+ \frac{c_2}{\phi(r)} \int_{B_R} u^2 \, d\mu. \end{split}$$

Hence,

$$\int_{B_{R+r}} d\Gamma(v\varphi, v\varphi) \leq 2 \int_{B_{R+r}} v^2 d\Gamma(\varphi, \varphi) + 2 \int_{B_R \times B_{R+r}} \varphi^2(x) (v(x) - v(y))^2 J(dx, dy) + 2 \int_{B_R} \varphi^2 d\Gamma_c(v, v) + \frac{c_2}{\phi(r)} \int_{B_R} u^2 d\mu.$$

$$(2.12)$$

Combining (2.11) with (2.12), we have for a > 0,

$$a \int_{B_{R+r}} d\Gamma(v\varphi, v\varphi)$$

$$\leq (2a+4) \int_{B_{R+r}} v^2 d\Gamma(\varphi, \varphi)$$

$$+ \left(2a - \frac{3}{4}\right) \left(\int_{B_R} \varphi^2 d\Gamma_c(v, v) + \int_{B_R \times B_{R+r}} \varphi^2(x)(v(x) - v(y))^2 J(dx, dy)\right)$$

$$+ \frac{c_3(1+a)}{\phi(r)} \left[1 + \frac{1}{\theta} \left(1 + \frac{R}{r}\right)^{d_2 + \beta_{2,\phi} - \beta_{1,\phi}} \operatorname{Tail}_{\phi}(u; x_0, R+r)\right] \int_{B_R} u^2 d\mu.$$
(2.13)

Next by using (2.8) for v with $\rho = \infty$, we have

$$\begin{split} &\int_{B_{R+r}} v^2 \, d\Gamma(\varphi,\varphi) \\ &\leq \frac{1}{8} \left[\int_{B_R} \varphi^2 \, d\Gamma_c(v,v) + \int_{B_R \times B_{R+r}} \varphi^2(x) (v(x) - v(y))^2 \, J(dx,dy) \right] + \frac{c_0}{\phi(r)} \int_{B_{R+r}} v^2 \, d\mu \end{split}$$

$$\leq \frac{1}{8} \left[\int_{B_R} \varphi^2 \, d\Gamma_c(v,v) + \int_{B_R \times B_{R+r}} \varphi^2(x) (v(x) - v(y))^2 \, J(dx,dy) \right] + \frac{c_0}{\phi(r)} \int_{B_{R+r}} u^2 \, d\mu.$$

Plugging this into (2.13) with a = 1/9 (so that (4+2a)/8 + (2a - (3/4)) = 0), we obtain

$$\frac{1}{9} \int_{B_{R+r}} d\Gamma(v\varphi, v\varphi) \le \frac{c_4}{\phi(r)} \left[1 + \frac{1}{\theta} \left(1 + \frac{R}{r} \right)^{d_2 + \beta_{2,\phi} - \beta_{1,\phi}} \operatorname{Tail}_{\phi} \left(u; x_0, R+r \right) \right] \int_{B_{R+r}} u^2 \, d\mu,$$
ch proves the desired assertion.

which proves the desired assertion.

Proposition 2.15. (L²-mean value inequality) Assume VD, (1.10), $FK(\phi)$, $CS(\phi)$ and $J_{\phi,\leq}$ hold. For $x_0 \in M$ and R > 0, let u be a bounded \mathcal{E} -subharmonic function in $B(x_0, R)$. Then for any $\delta > 0$,

ess sup
$$_{B(x_0,R/2)} u \le c_1 \left[\left(\frac{(1+\delta^{-1})^{1/\nu}}{V(x_0,R)} \int_{B(x_0,R)} u^2 d\mu \right)^{1/2} + \delta \operatorname{Tail}_{\phi} (u;x_0,R/2) \right],$$

where ν is the constant appearing in the FK(ϕ) inequality (1.20), and $c_1 > 0$ is a constant independent of x_0 , R and δ . In particular, there is a constant c > 0 independent of x_0 and R so that

ess
$$\sup_{B(x_0, R/2)} u \le c \left[\left(\frac{1}{V(x_0, R)} \int_{B(x_0, R)} u^2 d\mu \right)^{1/2} + \operatorname{Tail}_{\phi} (u; x_0, R/2) \right].$$

Proof. With the aid of (2.10), one can see that the comparison results over balls as stated in [CKW1, Lemma 4.8] still hold true. We can then follow the proof of [CKW1, Proposition 4.10] line by the line to obtain the desired assertion. We omit details here.

In the following, we consider L^2 and L^1 mean value inequalities for \mathcal{E} -subharmonic functions for truncated Dirichlet forms. In the truncated situation we can no longer use the nonlocal tail of subharmonic functions. The remedy is to enlarge the integral regions in the right hand side of the mean value inequalities. Since the proof is almost the same as these of [CKW1, Proposition 4.11 and Corollary 4.12 (with some necessary modifications as done in the proof of Lemma 2.14), we omit it here.

Proposition 2.16. (L^2 and L^1 mean value inequalities for ρ -truncated Dirichlet forms) Assume VD, (1.10), $FK(\phi)$, $CS(\phi)$ and $J_{\phi,<}$ hold. There are positive constants $c_1, c_2 > 0$ so that for $x_0 \in M$, $\rho, R > 0$, and for any bounded $\mathcal{E}^{(\rho)}$ -subharmonic function u on $B(x_0, R)$ for the ρ -truncated Dirichlet form $(\mathcal{E}^{(\rho)}, \mathcal{F})$, we have

ess sup
$$_{B(x_0,R/2)}u^2 \le \frac{c_1}{V(x_0,R)} \left(1+\frac{\rho}{R}\right)^{d_2/\nu} \left(1+\frac{R}{\rho}\right)^{\beta_{2,\phi}/\nu} \int_{B(x_0,R+\rho)} u^2 d\mu;$$

(ii)

ess
$$\sup_{B(x_0, R/2)} u \le \frac{c_2}{V(x_0, R)} \left(1 + \frac{\rho}{R}\right)^{d_2/\nu} \left(1 + \frac{R}{\rho}\right)^{\beta_{2,\phi}/\nu} \int_{B(x_0, R+\rho)} u \, d\mu.$$
 (2.14)

Here, ν is the constant in FK(ϕ), d_2 and $\beta_{2,\phi}$ are the exponents in (1.9) from VD and (1.15) respectively.

3 Implications of heat kernel estimates

First we note that by the same proof of [CKW1, Proposition 7.6], we have the following.

Proposition 3.1. Under VD, RVD and (1.10),

 $\mathrm{UHKD}(\phi) \Longrightarrow \mathrm{FK}(\phi) \qquad and \qquad \mathrm{PI}(\phi) \Longrightarrow \mathrm{FK}(\phi).$

Denote by ζ the lifetime of the Hunt process X associated with the regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(M; \mu)$. We have the following two facts.

Proposition 3.2. (i) Under VD, $NL(\phi) \Longrightarrow \zeta = \infty$.

(ii) Assume that VD, (1.10), E_{ϕ} and $J_{\phi,\leq}$ hold. Then $(\mathcal{E},\mathcal{F})$ is conservative; that is, $\zeta = \infty$.

Proof. (i) is taken directly from [CKW1, Proposition 3.1(ii)], which holds for any symmetric Markov process.

(ii) can be proved by exactly the same argument as that of [CKW1, Lemma 4.21]. The details are omitted here. $\hfill \Box$

3.1 UHK $(\phi_c, \phi_j) + (\mathcal{E}, \mathcal{F})$ is conservative $\Longrightarrow J_{\phi_j, \leq}$ and HK $_-(\phi_c, \phi_j) \Longrightarrow J_{\phi_j}$

Proposition 3.3. Under VD and (1.10),

UHK
$$(\phi_c, \phi_i)$$
 and $(\mathcal{E}, \mathcal{F})$ is conservative $\Longrightarrow J_{\phi_i, <}$

and

$$\operatorname{HK}_{-}(\phi_{c},\phi_{j}) \Longrightarrow \operatorname{J}_{\phi_{j}}$$

In particular, $HK(\phi_c, \phi_j) \Longrightarrow J_{\phi_j}$.

Proof. We only prove that case that $HK_{-}(\phi_{c}, \phi_{j}) \Longrightarrow J_{\phi_{j}}$, and the other two cases can be verified similarly. For t > 0, consider the bilinear form $\mathcal{E}^{(t)}(f,g) := \langle f - P_{t}f, g \rangle / t$. Since $(\mathcal{E}, \mathcal{F})$ is conservative by Proposition 3.2(i), we can write

$$\mathcal{E}^{(t)}(f,g) = \frac{1}{2t} \int_M \int_M (f(x) - f(y))(g(x) - g(y))p(t,x,y)\,\mu(dx)\,\mu(dy).$$

It is well known that $\lim_{t\to 0} \mathcal{E}^{(t)}(f,g) = \mathcal{E}(f,g)$ for all $f,g \in \mathcal{F}$. Let A, B be disjoint compact sets, and take $f,g \in \mathcal{F}$ such that $\operatorname{supp} f \subset A$ and $\operatorname{supp} g \subset B$. Then in view of the strongly local property of $\mathcal{E}^{(c)}(f,g)$,

$$\mathcal{E}^{(t)}(f,g) = -\frac{1}{t} \int_A \int_B f(x)g(y)p(t,x,y)\,\mu(dy)\,\mu(dx) \xrightarrow{t \to 0} - \int_A \int_B f(x)g(y)\,J(dx,dy).$$

Let $r_0 := d(A, B)$. For any $0 < t \le \phi_c(r_0)$, by VD, (1.10), (1.12) and (1.13),

$$\sup_{x \in A, y \in B} \frac{1}{V(x, \phi_c^{-1}(t))} \exp\left(-c_1 \frac{d(x, y)}{\bar{\phi}_c^{-1}(t/d(x, y))}\right)$$

$$\begin{split} &\leq c_2 \sup_{x \in A, y \in B} \frac{1}{V(x, d(x, y))} \left(\frac{d(x, y))}{\phi_c^{-1}(t)}\right)^{d_2} \exp\left(-c_3 \left(\frac{\phi_c(d(x, y))}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\leq c_4 \sup_{x \in A, y \in B} \frac{1}{V(x, d(x, y))} \left(\frac{\phi_c(d(x, y)))}{t}\right)^{d_2\beta_{2,\phi_c}} \exp\left(-c_3 \left(\frac{\phi_c(d(x, y))}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\leq c_5 \sup_{x \in A, y \in B} \frac{1}{V(x, d(x, y))} \exp\left(-\frac{c_3}{2} \left(\frac{\phi_c(d(x, y))}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\leq c_6 \sup_{x \in A} \frac{1}{V(x, r_0)} \exp\left(-\frac{c_3}{2} \left(\frac{\phi_c(r_0))}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right), \end{split}$$

where in the third inequality we used the fact that

$$r^{d_2\beta_{2,\phi_c}} \le c_7 \exp\left(\frac{c_3}{2}r^{1/(\beta_{2,\phi_c}-1)}\right), \quad r \ge 1.$$

The inequality above yields that

$$\lim_{t \to 0} \frac{1}{t} \int_A \int_B f(x)g(y)p^{(c)}(t,x,y)\,\mu(dy)\,\mu(dx) = 0.$$

Hence, using $HK_{-}(\phi_c, \phi_j)$, we obtain

$$\int_A \int_B f(x)g(y) J(dx, dy) \asymp \int_A \int_B \frac{f(x)g(y)}{V(x, d(x, y))\phi_j(d(x, y))} \, \mu(dy) \, \mu(dx)$$

for all $f, g \in \mathcal{F}$ such that supp $f \subset A$ and supp $g \subset B$. Since A, B are arbitrary disjoint compact sets, it follows that J(dx, dy) is absolutely continuous w.r.t. $\mu(dx) \mu(dy)$, and J_{ϕ_j} holds. \Box

3.2 UHK(ϕ_c, ϕ_i) and $(\mathcal{E}, \mathcal{F})$ is conservative \Longrightarrow Gcap(ϕ)

In this subsection, we give the proof that $UHK(\phi_c, \phi_j)$ and the conservativeness of $(\mathcal{E}, \mathcal{F})$ imply $Gcap(\phi)$.

We begin with the following lemma.

Lemma 3.4. Assume that VD, (1.10) and UHK(ϕ_c, ϕ_j) hold and that $(\mathcal{E}, \mathcal{F})$ is conservative. Then $\text{EP}_{\phi,\leq}$ holds.

Proof. We first verify that there is a constant $c_1 > 0$ such that for each t, r > 0 and for almost all $x \in M$,

$$\int_{B(x,r)^c} p(t,x,y)\,\mu(dy) \le \frac{c_1t}{\phi(r)}.$$

Indeed, we only need to consider the case that $\phi(r) > t$; otherwise, the inequality above holds trivially with $c_1 = 1$. According to UHK (ϕ_c, ϕ_j) , VD, (1.10), (1.12) and (1.13), for any t, r > 0 with $\phi(r) > t$ and almost all $x \in M$,

$$\int_{B(x,r)^c} p(t,x,y)\,\mu(dy)$$

$$\begin{split} &= \sum_{i=0}^{\infty} \int_{B(x,2^{i+1}r) \setminus B(x,2^{i}r)} p(t,x,y) \, \mu(dy) \\ &\leq \sum_{i=0}^{\infty} \frac{c_2 V(x,2^{i+1}r)}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{c_3 2^i r}{\bar{\phi}_c^{-1}(t/(2^i r))}\right) + \sum_{i=0}^{\infty} \frac{c_2 t V(x,2^{i+1}r)}{V(x,2^{i}r)\phi_j(2^i r)} \\ &\leq c_4 \sum_{i=0}^{\infty} \left(\frac{2^i r}{\phi_c^{-1}(t)}\right)^{d_2} \exp\left(-\frac{c_5 2^i r}{\bar{\phi}_c^{-1}(t/r)}\right) + \frac{c_4 t}{\phi_j(r)} \sum_{i=0}^{\infty} 2^{-i\beta_{1,\phi_j}} \\ &\leq c_6 \sum_{i=0}^{\infty} \left(2^i \left(\frac{\phi_c(r)}{t}\right)^{1/\beta_{1,\phi_c}}\right)^{d_2} \exp\left(-c_7 2^i \left(\frac{\phi_c(r)}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) + \frac{c_6 t}{\phi_j(r)} \\ &\leq c_6 \sum_{i=0}^{\infty} \left(2^{i(\beta_{2,\phi_c}-1)} \frac{\phi_c(r)}{t}\right)^{d_2(1+1/(\beta_{2,\phi_c}-1))} \exp\left(-c_7 \left(2^{i(\beta_{2,\phi_c}-1)} \frac{\phi_c(r)}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) + \frac{c_6 t}{\phi_j(r)} \\ &\leq c_8 \sum_{i=0}^{\infty} \exp\left(-\frac{c_7}{2} \left(2^{i(\beta_{2,\phi_c}-1)} \frac{\phi_c(r)}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) + \frac{c_6 t}{\phi_j(r)} \\ &\leq c_9 t \left(\frac{1}{\phi_c(r)} + \frac{1}{\phi_j(r)}\right) \leq \frac{c_{10} t}{\phi(r)}, \end{split}$$

where in the arguments above we used $\phi_c(r) \wedge \phi_j(r) = \phi(r)$ for all r > 0, in the second inequality we used the fact that $\bar{\phi}_c(r)$ is increasing on $(0, \infty)$, in the fourth inequality we used the fact that $\beta_{1,\phi_c} > 1$, and in the fifth and the sixth inequalities we used the facts that there is a constant $c_{11} > 0$ such that

$$s^{d_2(1+1/(\beta_{2,\phi_c}-1))} \le c_{11} \exp\left(\frac{c_7}{2}s^{1/(\beta_{2,\phi_c}-1)}\right), \quad s \ge 1$$

and

$$\sum_{i=0}^{\infty} \exp\left(-\frac{c_7}{2}(2^i s)^{1/(\beta_{2,\phi_c}-1)}\right) \le c_{11}/s, \quad s \ge 1,$$

respectively.

Now, since $(\mathcal{E}, \mathcal{F})$ is conservative, by the strong Markov property, for any each t, r > 0 and for almost all $x \in M$,

$$\mathbb{P}^{x}(\tau_{B(x,r)} \leq t) = \mathbb{P}^{x}(\tau_{B(x,r)} \leq t, X_{2t} \in B(x, r/2)^{c}) + \mathbb{P}^{x}(\tau_{B(x,r)} \leq t, X_{2t} \in B(x, r/2)) \\
\leq \mathbb{P}^{x}(X_{2t} \in B(x, r/2)^{c}) + \sup_{\substack{z \notin B(x,r)^{c}, s \leq t}} \mathbb{P}^{z}(X_{2t-s} \in B(z, r/2)^{c}) \\
\leq \frac{c_{13}t}{\phi(r)},$$

which yields $\text{EP}_{\phi,\leq}$. (Note that the conservativeness of $(\mathcal{E},\mathcal{F})$ is used in the equality above. Indeed, without the conservativeness, there must be an extra term $\mathbb{P}^x(\tau_{B(x,r)} \leq t, \zeta \leq 2t)$ in the right hand side of the above equality, where ζ is the lifetime of X.)

Proposition 3.5. Suppose that VD, (1.10) and UHK(ϕ_c, ϕ_j) hold, and that $(\mathcal{E}, \mathcal{F})$ is conservative. Then Gcap(ϕ) holds.

Proof. According to Lemma 3.4, $EP_{\phi,\leq}$ hold true. Thus the desired assertion follows from Proposition 2.4.

4 Implications of $FK(\phi)$, $NDL(\phi)$, $CS(\phi)$ and $J_{\phi_i,\leq}$

4.1 $NDL(\phi) \Longrightarrow PI(\phi) + E_{\phi}$

Proposition 4.1. Assume that VD and (1.10) hold.

(i) NDL(ϕ) \Longrightarrow PI(ϕ) + E $_{\phi,\geq}$.

(ii) If in addition RVD holds, then $NDL(\phi) \Longrightarrow E_{\phi}$.

Proof. The proof is the same as that of [CKW2, Proposition 3.5], so it is omitted. \Box

4.2 $FK(\phi) + J_{\phi,<} + CS(\phi) \Longrightarrow E_{\phi}$ and $FK(\phi) + J_{\phi,<} + E_{\phi} \Longrightarrow UHKD(\phi)$

The next two propositions can be proved by following the arguments in [CKW1], which are based on the probabilistic ideas and are valid for general symmetric Dirichlet forms.

Proposition 4.2. Assume VD, (1.10), $FK(\phi)$, $J_{\phi,\leq}$ and $CS(\phi)$ hold. Then E_{ϕ} holds.

Proof. According to [CKW1, Lemma 4.14], $E_{\phi,\leq}$ holds under VD, (1.10) and FK(ϕ). On the other hand, by Proposition 2.16, we have the L^1 -mean value inequality (2.14) under the assumptions of Proposition 4.2. Then by the proofs of [CKW1, Lemmas 4.15, 4.16 and 4.17], we obtain $E_{\phi,\geq}$.

Proposition 4.3. Suppose that VD, (1.10), $FK(\phi)$, E_{ϕ} and $J_{\phi,\leq}$ hold. Then $UHKD(\phi)$ is satisfied, i.e., there is a constant c > 0 such that for all $x \in M_0$ and t > 0,

$$p(t, x, x) \le \frac{c}{V(x, \phi^{-1}(t))}.$$

Proof. The proof is the same as that for [CKW1, Theorem 4.25], and so we omit the details here. \Box

4.3 UHKD $(\phi) + J_{\phi_j,\leq} + E_{\phi} \Longrightarrow$ UHK (ϕ_c, ϕ_j)

First we have the following by the proof of [CKW1, Lemma 5.2].

Lemma 4.4. Under VD and (1.10), if UHKD(ϕ), $J_{\phi,\leq}$ and E_{ϕ} hold, then the ρ -truncated Dirichlet form ($\mathcal{E}^{(\rho)}, \mathcal{F}$) has the heat kernel $q^{(\rho)}(t, x, y)$, and it satisfies that for any t > 0 and all $x, y \in M_0$,

$$q^{(\rho)}(t,x,y) \le c_1 \left(\frac{1}{V(x,\phi^{-1}(t))} + \frac{1}{V(y,\phi^{-1}(t))} \right) \exp\left(c_2 \frac{t}{\phi(\rho)} - c_3 \frac{d(x,y)}{\rho} \right),$$

where c_1, c_2, c_3 are positive constants independent of ρ . Consequently, for any t > 0 and all $x, y \in M_0$,

$$q^{(\rho)}(t,x,y) \le \frac{c_4}{V(x,\phi^{-1}(t))} \left(1 + \frac{d(x,y)}{\phi^{-1}(t)}\right)^{d_2} \exp\left(c_2 \frac{t}{\phi(\rho)} - c_3 \frac{d(x,y)}{\rho}\right).$$

From now till the end of this subsection, we will assume that $J_{\phi_j,\leq}$ is satisfied. Note that since $\phi(r) \leq \phi_j(r)$ for all r > 0, condition $J_{\phi_j,\leq}$ implies $J_{\phi,\leq}$.

Recall that there is close relation between p(t, x, y) and $q^{(\rho)}(t, x, y)$ via Meyer's decomposition, e.g. see [CKW1, Section 7.2]. In particular, according to [CKW1, (4.34) and Proposition 4.24 (and its proof)], for any $t, \rho > 0$ and all $x, y \in M_0$,

$$p(t,x,y) \le q^{(\rho)}(t,x,y) + \frac{c_1 t}{V(x,\rho)\phi_j(\rho)} \exp\left(\frac{c_1 t}{\phi(\rho)}\right),\tag{4.1}$$

where c_1 is independent of $t, \rho > 0$ and $x, y \in M_0$.

The following lemma is essentially taken from [BKKL, Lemma 4.3], which is partly motivated by the proof of [CKW1, Proposition 5.3]. For the sake of the completeness and for further applications, we spell out its proof here.

Lemma 4.5. Assume that VD, (1.10), UHKD(ϕ), $J_{\phi_j,\leq}$ and E_{ϕ} hold. Let $f : \mathbb{R}_+ \times \mathbb{R}_+ \to \mathbb{R}_+$ be a measurable function such that $t \mapsto f(r,t)$ is non-increasing for all r > 0, and that $r \mapsto f(r,t)$ is non-decreasing for all t > 0. Suppose that the following hold:

- (i) For each b > 0, $\sup_{t>0} f(b\phi^{-1}(t), t) < \infty$.
- (ii) There exist constants $\eta \in (0, \beta_{1,\phi_i}]$ and $a_1, c_1 > 0$ such that for all $x \in M_0$ and r, t > 0,

$$\int_{B(x,r)^c} p(t,x,y) \,\mu(dy) \le c_1 \left(\frac{\phi_j^{-1}(t)}{r}\right)^{\eta} + c_1 \exp\left(-a_1 f(r,t)\right)$$

Then there exist constants $k, c_0 > 0$ such that for all $x, y \in M_0$ and t > 0,

$$p(t,x,y) \leq \frac{c_0 t}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{c_0}{V(x,\phi^{-1}(t))} \left(1 + \frac{d(x,y)}{\phi^{-1}(t)}\right)^{d_2} \exp\left(-a_1 k f(d(x,y)/(16k),t)\right)$$

Furthermore, the conclusion still holds true for any $t \in (0,T]$ or $t \in [T,\infty)$ with some T > 0, if assumptions (i) and (ii) above are restricted on the corresponding time interval.

Proof. We only consider the case that $t \in (0, \infty)$, since the other cases can be treated similarly. We first note that, by E_{ϕ} and $J_{\phi_j,\leq}$, the Dirichlet form $(\mathcal{E}, \mathcal{F})$ is conservative by Proposition 3.2(ii). For fixed $x_0 \in M$, set $B_s := B(x_0, s)$ for all s > 0. By the conservativeness of the Dirichlet form $(\mathcal{E}, \mathcal{F})$, the strong Markov property, assumption (ii) and the fact that $t \mapsto f(r, t)$ is non-increasing, we have that for any $x \in B_{r/4} \cap M_0$ and $t \ge 0$,

$$\mathbb{P}^{x}(\tau_{B_{r}} \leq t) \leq c_{1} \left(\frac{\phi_{j}^{-1}(t)}{r}\right)^{\eta} + c_{1} \exp\left(-a_{1}f(r/4, t)\right);$$

see the end of the proof for Lemma 3.4. For any $\rho > 0$ and any subset $D \subset M$, denote by $\tau_D^{(\rho)} = \inf\{t > 0 : X_t^{(\rho)} \notin D\}$, where $(X_t^{(\rho)})_{t \ge 0}$ is the symmetric Hunt process associated with the ρ -truncated Dirichlet form $(\mathcal{E}^{(\rho)}, \mathcal{F})$. By [CKW1, Lemma 7.8], $J_{\phi_j,\leq}$ and Lemma 2.1, we have that for all $x \in B_{r/4} \cap M_0$ and r > 0,

$$\mathbb{P}^{x}(\tau_{B_{r}}^{(r)} \leq t) \leq c_{1} \left(\frac{\phi_{j}^{-1}(t)}{r}\right)^{\eta} + c_{1} \exp\left(-a_{1}f(r/4,t)\right) + \frac{c_{2}t}{\phi_{j}(r)} =: \Phi(r,t).$$

This together with [CKW1, Lemma 7.1] yields that for all $t > 0, x \in M_0$ and $k \ge 1$,

$$\int_{B(x,2kr)^c} q^{(\rho)}(t,x,y)\,\mu(dy) \le \Phi(r,t)^k.$$
(4.2)

Let $k = [(\beta_{2,\phi_j} + 2d_2)/\eta] + 1$. For any $x, y \in M_0$ and t > 0 with $4k\phi^{-1}(t) \ge d(x,y)$, it follows from the assumption that $r \mapsto f(r,t)$ is non-decreasing and assumption (i) that $f(d(x,y)/(16k),t) \le f(\phi^{-1}(t)/4,t) \le A < \infty$. Thus, for any $x, y \in M_0$ and t > 0 with $4k\phi^{-1}(t) \ge d(x,y)$, according to UHKD(ϕ) and VD,

$$p(t, x, y) \leq c_3 \left(\frac{1}{V(x, \phi^{-1}(t))} + \frac{1}{V(y, \phi^{-1}(t))} \right)$$
$$\leq \frac{c_4 e^{a_1 k A}}{V(x, \phi^{-1}(t))} \left(1 + \frac{d(x, y)}{\phi^{-1}(t)} \right)^{d_2} \exp\left(-a_1 k f(d(x, y)/(16k), t)\right).$$

Next we consider the case that $x, y \in M_0$ and t > 0 with $4k\phi^{-1}(t) \le d(x, y)$. Letting r = d(x, y) and $\rho = r/(4k)$, it holds

$$\begin{split} q^{(\rho)}(t,x,y) &= \int_{M} q^{(\rho)}(t/2,x,z) q^{(\rho)}(t/2,z,y) \, \mu(dz) \\ &\leq \left(\int_{B(x,r/2)^{c}} + \int_{B(y,r/2)^{c}} \right) q^{(\rho)}(t/2,x,z) q^{(\rho)}(t/2,z,y) \, \mu(dz) \\ &\leq \frac{c_{5}}{V(x,\phi^{-1}(t))} \left(1 + \frac{d(x,y)}{\phi^{-1}(t)} \right)^{d_{2}} \Phi(\rho,t/2)^{k}. \end{split}$$

Here, the last inequality follows from (4.2) and VD as well as

$$q^{(\rho)}(t,x,z) \le \frac{c_0}{V(x,\phi^{-1}(t))} e^{c_0 t/\phi(\rho)} \le \frac{c_0 e^{c_0}}{V(x,\phi^{-1}(t))}, \quad x,z \in M_0, t > 0 \text{ with } \phi^{-1}(t) \le \rho,$$

which follows from [CKW1, Lemma 5.1] (based on UHKD(ϕ) and E_{ϕ}) and [CKW1, Lemma 7.8] (based on $J_{\phi_j,\leq}$ and the fact that $\phi_j(r) \geq \phi(r)$ for all r > 0). Since $\rho \geq \phi^{-1}(t) \geq \phi_j^{-1}(t)$ and $k\beta_{1,\phi_j} \geq k\eta \geq \beta_{2,\phi_j} + 2d_2$, by (1.10),

$$\left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{\eta k} + \left(\frac{t}{\phi_j(\rho)}\right)^k \le c_6 \left[\left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{\beta_{2,\phi_j} + 2d_2} + \left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{k\beta_{1,\phi_j}} \right] \le c_7 \left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{\beta_{2,\phi_j} + 2d_2}$$

Thus for all $x, y \in M_0$ and $t, \rho > 0$ with $\rho \ge \phi^{-1}(t) \ge \phi_j^{-1}(t)$,

$$\begin{split} &\frac{1}{V(x,\phi^{-1}(t))} \left(1 + \frac{d(x,y)}{\phi^{-1}(t)}\right)^{d_2} \left[\left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{\eta k} + \left(\frac{t}{\phi_j(\rho)}\right)^k \right] \\ &\leq \frac{c_7}{V(x,\phi^{-1}(t))} \left(1 + \frac{d(x,y)}{\phi^{-1}(t)}\right)^{d_2} \left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{\beta_{2,\phi_j} + 2d_2} \leq \frac{c_8}{V(x,\rho)} \left(\frac{\rho}{\phi_j^{-1}(t)}\right)^{2d_2} \left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{\beta_{2,\phi_j} + 2d_2} \\ &\leq \frac{c_9 t}{V(x,\phi_j^{-1}(t))\phi_j(d(x,y))}, \end{split}$$

where the second inequality follows from VD and the fact that $\rho \ge \phi^{-1}(t) \ge \phi_j^{-1}(t)$, while in the last inequality we used (1.10). Hence, for any $x, y \in M_0$ and t > 0 with $4k\phi^{-1}(t) \le d(x, y)$,

$$\begin{split} &q^{(\rho)}(t,x,y) \\ &\leq \frac{c_{10}}{V(x,\phi^{-1}(t))} \left(1 + \frac{d(x,y)}{\phi^{-1}(t)}\right)^{d_2} \left(\left(\frac{\phi_j^{-1}(t)}{\rho}\right)^{\eta k} + \left(\frac{t}{\phi_j(\rho)}\right)^k + \exp\left(-a_1 k f(\rho/4,t)\right)\right) \\ &\leq \frac{c_{11} t}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{c_{11}}{V(x,\phi^{-1}(t))} \left(1 + \frac{d(x,y)}{\phi^{-1}(t)}\right)^{d_2} \exp\left(-a_1 k f(d(x,y)/(16k),t)\right). \end{split}$$

The desired assertion now follows from (4.1), $J_{\phi_i,\leq}$ and Lemma 2.1.

Proposition 4.6. Under VD and (1.10), if UHKD(ϕ), $J_{\phi_j,\leq}$ and E_{ϕ} hold, then we have UHK(ϕ_c, ϕ_j).

To prove Proposition 4.6, we will use Lemma 4.5 and need the following lemma.

Lemma 4.7. Assume that VD, (1.10), UHKD(ϕ), $J_{\phi_j,\leq}$ and E_{ϕ} hold. Then there exist constants a, c > 0 and $N \in \mathbb{N}$ such that for all $x, y \in M_0$ and t > 0,

$$p(t, x, y) \le \frac{ct}{V(x, d(x, y))\phi_j(d(x, y))} + \frac{c}{V(x, \phi^{-1}(t))} \exp\left(-\frac{ad(x, y)^{1/N}}{\phi^{-1}(t)^{1/N}}\right).$$

Proof. We claim that there exist $a_1, c_1 > 0$ such that for all $x \in M_0$ and t, r > 0

$$\int_{B(x,r)^c} p(t,x,y)\,\mu(dy) \le c_1 \left(\frac{\phi_j^{-1}(t)}{r}\right)^\eta + c_1 \exp\left(-\frac{a_1 r^{1/N}}{\phi^{-1}(t)^{1/N}}\right),\tag{4.3}$$

where $\eta = \beta_{1,\phi_j} - ((d_2 + \beta_{1,\phi_j})/N) \in (0,\beta_{1,\phi_j}]$ (by taking N large enough). If (4.3) holds, then the assertion follows from Lemma 4.5 by taking $f(r,t) = (r/\phi^{-1}(t))^{1/N}$.

When $r \leq \phi^{-1}(t)$, (4.3) holds trivially with $c_1 = e^{a_1}$. So it suffices to consider the case that $r \geq \phi^{-1}(t)$. According to (4.1) and Lemma 4.4, for any $t, \rho > 0$ and $x, y \in M_0$,

$$p(t, x, y) \le \frac{c_2}{V(x, \phi^{-1}(t))} \left(1 + \frac{d(x, y)}{\phi^{-1}(t)} \right)^{d_2} \exp\left(\frac{c_2 t}{\phi(\rho)} - \frac{c_3 d(x, y)}{\rho}\right) + \frac{c_2 t}{V(x, \rho)\phi_j(\rho)} \exp\left(\frac{c_2 t}{\phi(\rho)}\right).$$

Take $\alpha \in (d_2/(d_2 + \beta_{1,\phi_j}), 1)$, and define

$$\rho_n := \rho_n(r, t) = 2^{n\alpha} r^{1 - 1/N} \phi^{-1}(t)^{1/N}, \quad n \in \mathbb{N} \cup \{0\}$$

Since $r \ge \phi^{-1}(t)$, $\phi^{-1}(t) \le \rho_n \le 2^{n\alpha}r$ for all $n \in \mathbb{N} \cup \{0\}$. In particular, $t \le \phi(\rho_n)$ for all $n \in \mathbb{N} \cup \{0\}$. Thus, for any $x \in M_0$ and t, r > 0,

$$\begin{split} \int_{B(x,r)^c} p(t,x,y) \, \mu(dy) &= \sum_{n=0}^{\infty} \int_{B(x,2^{n+1}r) \setminus B(x,2^n r)} p(t,x,y) \, \mu(dy) \\ &\leq \frac{c_2 e^{c_2}}{V(x,\phi^{-1}(t))} \sum_{n=0}^{\infty} \left(1 + \frac{2^{n+1}r}{\phi^{-1}(t)}\right)^{d_2} \exp\left(-\frac{c_3 2^n r}{\rho_n}\right) V(x,2^{n+1}r) \end{split}$$

$$+ c_2 e^{c_2} t \sum_{n=0}^{\infty} \frac{V(x, 2^{n+1}r)}{V(x, \rho_n)\phi_j(\rho_n)}$$

=: $I_1 + I_2$.

On the one hand, by the definition of ρ_n and VD,

$$I_{1} = \frac{c_{2}e^{c_{2}}}{V(x,\phi^{-1}(t))} \sum_{n=0}^{\infty} \left(1 + \frac{2^{n+1}r}{\phi^{-1}(t)}\right)^{d_{2}} \exp\left(-c_{3}2^{n(1-\alpha)}\left(\frac{r}{\phi^{-1}(t)}\right)^{1/N}\right) V(x,2^{n+1}r)$$

$$\leq c_{4} \sum_{n=0}^{\infty} \left(1 + \frac{2^{n+1}r}{\phi^{-1}(t)}\right)^{2d_{2}} \exp\left(-c_{3}2^{n(1-\alpha)}\left(\frac{r}{\phi^{-1}(t)}\right)^{1/N}\right)$$

$$\leq c_{5} \sum_{n=0}^{\infty} \exp\left(-\frac{c_{3}}{2}2^{n(1-\alpha)}\left(\frac{r}{\phi^{-1}(t)}\right)^{1/N}\right) \leq c_{6} \exp\left(-c_{7}\left(\frac{r}{\phi^{-1}(t)}\right)^{1/N}\right),$$

where in the second inequality we used the fact that there is a constant $c_8 > 0$ such that for all $n \ge 1$ and $s \ge 1$,

$$(1+2^{n+1}s)^{2d_2} \le c_8 \exp\left(\frac{c_3}{2}2^{n(1-\alpha)}s^{1/N}\right).$$

On the other hand, according to VD and (1.10),

$$I_{2} \leq c_{9} \sum_{n=0}^{\infty} \left(\frac{2^{n}r}{\rho_{n}}\right)^{d_{2}} \left(\frac{\phi_{j}^{-1}(t)}{\rho_{n}}\right)^{\beta_{1,\phi_{j}}}$$
$$\leq c_{10} \sum_{n=0}^{\infty} 2^{n((1-\alpha)d_{2}-\alpha\beta_{1,\phi_{j}})} \left(\frac{\phi_{j}^{-1}(t)}{r}\right)^{\beta_{1,\phi_{j}}-((d_{2}+\beta_{1,\phi_{j}})/N)} \leq c_{11} \left(\frac{\phi_{j}^{-1}(t)}{r}\right)^{\eta}.$$

where the first and the second inequalities follow from the facts that $\phi(t) \leq \phi_j(t)$ and so $\phi^{-1}(t) \geq \phi_j^{-1}(t)$, and in the last inequality we used $\eta = \beta_{1,\phi_j} - ((d_2 + \beta_{1,\phi_j})/N)$ and $\alpha \in (d_2/(d_2 + \beta_{1,\phi_j}), 1)$. Combining estimates for I_1 and I_2 , we obtain (4.3). The proof is complete.

Now, we give the

Proof of Proposition 4.6. The proof is split into two cases.

(1) We first consider the case that $t \ge 1$. By UHKD(ϕ), we only need to check the case that $x, y \in M_0$ and $t \ge 1$ with $d(x, y) \ge \phi_j^{-1}(t)$. First, according to Lemma 4.7, there exist constants a, c > 0 and $N \in \mathbb{N}$ such that for all $x, y \in M_0$ and $t \ge 1$,

$$p(t, x, y) \le \frac{ct}{V(x, d(x, y))\phi_j(d(x, y))} + \frac{c}{V(x, \phi_j^{-1}(t))} \exp\left(-\frac{ad(x, y)^{1/N}}{\phi_j^{-1}(t)^{1/N}}\right).$$

Furthermore, we find by VD and (1.10) that

$$\begin{aligned} \frac{1}{V(x,\phi_j^{-1}(t))} \exp\left(-\frac{ad(x,y)^{1/N}}{\phi_j^{-1}(t)^{1/N}}\right) &\leq \frac{c_1}{V(x,d(x,y))} \left(\frac{d(x,y)}{\phi_j^{-1}(t)}\right)^{d_2} \exp\left(-\frac{ad(x,y)^{1/N}}{\phi_j^{-1}(t)^{1/N}}\right) \\ &\leq \frac{c_2}{V(x,d(x,y))} \exp\left(-\frac{a}{2} \frac{d(x,y)^{1/N}}{\phi_j^{-1}(t)^{1/N}}\right) \end{aligned}$$

$$\leq \frac{c_3}{V(x, d(x, y))} \left(\frac{\phi_j^{-1}(t)}{d(x, y)}\right)^{\beta_{2, \phi_j}} \\ \leq \frac{c_3 t}{V(x, d(x, y))\phi_j(d(x, y))},$$
(4.4)

where in the second and the third inequalities we used the facts that

$$r^{d_2} \le c_4 e^{ar^{1/N}/2}$$
 and $e^{-ar^{1/N}/2} \le c_5 r^{-\beta_{2,\phi_j}}$ for $r \ge 1$,

respectively. Hence for all $x, y \in M_0$ and $t \ge 1$ with $d(x, y) \ge \phi_j^{-1}(t)$,

$$p(t, x, y) \le \frac{c_6 t}{V(x, d(x, y))\phi_j(d(x, y))}$$

We claim that for any $t \ge 1$ and $x, y \in M_0$ with $d(x, y) \ge \phi_j^{-1}(t)$,

$$\frac{1}{V(x,\phi_c^{-1}(t))}\exp\left(-c_7\frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right) \le \frac{c_{11}t}{V(x,d(x,y))\phi_j(d(x,y))}.$$

Indeed, since $\phi_c(t) \ge \phi_j(t)$ for all $t \ge 1$, for any $t \ge 1$ and $x, y \in M_0$ with $d(x, y) \ge \phi_j^{-1}(t)$,

$$\begin{split} &\frac{1}{V(x,\phi_c^{-1}(t))} \exp\left(-c_7 \frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right) \\ &\leq \frac{c_8}{V(x,d(x,y))} \left(1 + \frac{d(x,y)}{\phi_c^{-1}(t)}\right)^{d_2} \exp\left(-c_7 \frac{d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right) \\ &\leq \frac{c_8}{V(x,d(x,y))} \left(1 + \frac{\phi_c(d(x,y))}{t}\right)^{d_2/\beta_{1,\phi_c}} \exp\left(-c_9 \left(\frac{\phi_c(d(x,y))}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\leq \frac{c_{10}}{V(x,d(x,y))} \exp\left(-\frac{c_9}{2} \left(\frac{\phi_c(d(x,y))}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\leq \frac{c_{10}}{V(x,d(x,y))} \exp\left(-\frac{c_9}{2} \left(\frac{\phi_j(d(x,y))}{t}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\leq \frac{c_{11}t}{V(x,d(x,y))\phi_j(d(x,y))}, \end{split}$$

where in the first inequality we used VD, the second inequality follows from (1.10), (1.12) and (1.13), and in the third and the last inequalities we applied the following two inequalities

$$(1+r)^{d_2/\beta_{1,\phi_c}} \le c_{12} \exp\left(\frac{c_9}{2}r^{1/(\beta_{2,\phi_c}-1)}\right), \quad r \ge 1$$

and

$$\exp\left(-\frac{c_9}{2}r^{1/(\beta_{2,\phi_c}-1)}\right) \le c_{13}r^{-1}, \quad r \ge 1,$$

respectively. This establishes $\text{UHK}(\phi_c, \phi_j)$ for the case that $t \geq 1$.

(2) We next consider the case of $t \leq 1$. It suffices to consider the case when $x, y \in M_0$ and $0 < t \leq 1$ with $d(x, y) \geq \phi_c^{-1}(t)$. By Lemma 4.5, it is enough to show that there exist constants $\eta \in (0, \beta_{1,\phi_j}]$ and $c_1, c_2 > 0$ such that for any $x \in M_0$, $0 < t \leq 1$ and r > 0,

$$\int_{B(x,r)^c} p(t,x,y) \,\mu(dy) \le c_1 \left(\frac{\phi_j^{-1}(t)}{r}\right)^{\eta} + c_1 \exp\left(-c_2 \frac{r}{\bar{\phi}_c^{-1}(t/r)}\right). \tag{4.5}$$

Indeed, by (4.5) and Lemma 4.5 with $f(r,t) = r/\bar{\phi}_c^{-1}(t/r)$, we have that for any $x, y \in M_0$ and $0 < t \leq 1$,

$$p(t,x,y) \le c_3 \left(\frac{t}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{1}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{c_0^*d(x,y)}{\bar{\phi}_c^{-1}(t/d(x,y))}\right) \right).$$

The desired assertion follows. In the following, we will prove (4.5). For this, we will consider four different cases.

(i) If $r \leq C_0 \phi_c^{-1}(t)$ for some constant $C_0 > 0$ whose exact valued will be determined in the step (ii) below, then by the non-decreasing property of ϕ_c , (1.12) and (1.13),

$$\exp\left(c_2 \frac{r}{\bar{\phi}_c^{-1}(t/r)}\right) \le \exp\left(c_2 \frac{C_0 \phi_c^{-1}(t)}{\bar{\phi}_c^{-1}(t/(C_0 \phi_c^{-1}(t)))}\right) \le e^{c_4}.$$

Hence, (4.5) holds trivially by taking $c_1 = e^{c_4}$.

(ii) Suppose that $C_0\phi_c^{-1}(t) \leq r \leq r_*(t)$, where $r_*(t)$ is to be determined later. For any $n \in \mathbb{N} \cup \{0\}$, define

$$\rho_n = c_* 2^{n\alpha} \bar{\phi}_c^{-1} \left(\frac{t}{r}\right)$$

with $d_2/(d_2 + \beta_{1,\phi_j}) < \alpha < 1$, where c_* is also determined later. Then by (4.1) and Lemma 4.4, for any $x, y \in M_0$ with $2^n r \leq d(x, y) \leq 2^{n+1}r$ and $0 < t \leq 1$,

$$p(t,x,y) \leq \frac{c_1}{V(x,\phi_c^{-1}(t))} \left(1 + \frac{2^{n+1}r}{\phi_c^{-1}(t)}\right)^{d_2} \exp\left(\frac{c_1t}{\phi(\rho_n)} - \frac{c_22^n r}{\rho_n}\right) + \frac{c_1t}{V(x,\rho_n)\phi_j(\rho_n)} \exp\left(\frac{c_1t}{\phi(\rho_n)}\right)$$
$$\leq \frac{c_3}{V(x,\phi_c^{-1}(t))} \left(1 + \frac{2^{n+1}r}{\phi_c^{-1}(t)}\right)^{d_2} \exp\left(\frac{c_1t}{\phi_c(\rho_n)} - \frac{c_22^n r}{\rho_n}\right) + \frac{c_3t}{V(x,\rho_n)\phi_j(\rho_n)} \exp\left(\frac{c_1t}{\phi_c(\rho_n)}\right)$$

where in the last inequality we used the fact that if $\rho_n \ge 1$, then $\phi(\rho_n) = \phi_j(\rho_n) \ge \phi_j(1) = 1 \ge t$ for all $t \in (0, 1]$. Hence, for any $x \in M_0$, $0 < t \le 1$ and $C_0 \phi_c^{-1}(t) \le r \le r_*(t)$,

$$\begin{split} &\int_{B(x,r)^c} p(t,x,y)\,\mu(dy) \\ &= \sum_{n=0}^{\infty} \int_{B(x,2^{n+1}r)\setminus B(x,2^nr)} p(t,x,y)\,\mu(dy) \\ &\leq \sum_{n=0}^{\infty} \frac{c_4 V(x,2^{n+1}r)}{V(x,\phi_c^{-1}(t))} \left(1 + \frac{2^{n+1}r}{\phi_c^{-1}(t)}\right)^{d_2} \exp\left(\frac{c_5 t}{c_*^{\beta_{1,\phi_c}} 2^{n\alpha\beta_{1,\phi_c}}\phi_c(\bar{\phi}_c^{-1}(t/r))} - \frac{c_2 2^{n(1-\alpha)}r}{c_*\bar{\phi}_c^{-1}(t/r)}\right) \\ &+ \sum_{n=0}^{\infty} \frac{c_4 t V(x,2^{n+1}r)}{V(x,\rho_n)\phi_j(\rho_n)} \exp\left(\frac{c_5 t}{c_*^{\beta_{1,\phi_c}} 2^{n\alpha\beta_{1,\phi_c}}\phi_c(\bar{\phi}_c^{-1}(t/r))}\right) \\ &=: I_1 + I_2, \end{split}$$

where the inequality above follows from (1.10), and c_1, \dots, c_5 are independent of c_* . On the one hand, by taking $c_* = (1 + (2c_2^*c_5/c_2))^{1/(\beta_{1,\phi_c}-1)}$ (with c_2^* being the constant c_2 in (1.12)), we find by VD and (1.12) that

$$I_1 \le c_6 \sum_{n=0}^{\infty} \left(\frac{2^n r}{\phi_c^{-1}(t)}\right)^{2d_2} \exp\left(-c_7 2^{n(1-\alpha)} \frac{r}{\bar{\phi}_c^{-1}(t/r)}\right) \le c_8 \exp\left(-c_9 \frac{r}{\bar{\phi}_c^{-1}(t/r)}\right).$$

Here and in what follows, the constants will depend on c_* . On the other hand, according to VD and (1.10) again,

$$\begin{split} I_{2} &\leq c_{10} \exp\left(\frac{c_{11}r}{\bar{\phi}_{c}^{-1}(t/r)}\right) \sum_{n=0}^{\infty} \left(\frac{2^{n}r}{\rho_{n}}\right)^{d_{2}} \frac{\phi_{j}(r)}{\phi_{j}(\rho_{n})} \frac{t}{\phi_{j}(r)} \\ &\leq c_{12} \exp\left(\frac{c_{11}r}{\bar{\phi}_{c}^{-1}(t/r)}\right) \left(\frac{r}{\bar{\phi}_{c}^{-1}(t/r)}\right)^{d_{2}+\beta_{2,\phi_{j}}} \frac{t}{\phi_{j}(r)} \sum_{n=0}^{\infty} 2^{n(d_{2}-\alpha(d_{2}+\beta_{1,\phi_{j}}))} \\ &\leq c_{13} \exp\left(\frac{c_{14}r}{\bar{\phi}_{c}^{-1}(t/r)}\right) \frac{t}{\phi_{j}(r)}, \end{split}$$

where in the last inequality we used the condition $d_2/(d_2 + \beta_{1,\phi_j}) < \alpha < 1$ and the fact that

$$r^{d_2+\beta_{2,\phi_j}} \le c_{15}e^r, \quad r \ge c_{16} > 0.$$

We note that the argument up to here is independent of the definition of $r_*(t)$ and the choice of the constant C_0 . Now, according to Lemma 4.8 below, we can find constants $C_0, c_{17} > 0$ and a unique $r_*(t) \in (C_0 \phi_c^{-1}(t), \infty)$ such that

$$\exp\left(\frac{2c_{14}r}{\beta_{1,\phi_j}\bar{\phi}_c^{-1}(t/r)}\right) \le \frac{c_{17}r}{\phi_j^{-1}(t)}, \quad C_0\phi_c^{-1}(t) \le r \le r_*(t)$$
(4.6)

and

$$\exp\left(\frac{2c_{14}r}{\beta_{1,\phi_j}\bar{\phi}_c^{-1}(t/r)}\right) \ge \frac{c_{17}r}{\phi_j^{-1}(t)}, \quad r \ge r_*(t),$$

as well as

$$\exp\left(\frac{2c_{14}r_*(t)}{\beta_{1,\phi_j}\bar{\phi}_c^{-1}(t/r_*(t))}\right) = \frac{c_{17}r_*(t)}{\phi_j^{-1}(t)}.$$
(4.7)

•

(Here, without loss of generality we may and do assume that $2c_{14} \ge \beta_{1,\phi_j}$.) Then due to (1.10) again, $\left(\sum_{j=1, \phi_j} \beta_{1,\phi_j} / 2 \right) = \left(\sum_{j=1, \phi_j} \beta_{1,\phi_j} / 2 \right)$

$$I_{2} \le c_{18} \left(\frac{r}{\phi_{j}^{-1}(t)}\right)^{\beta_{1,\phi_{j}}/2} \left(\frac{t}{\phi_{j}(r)}\right) \le c_{19} \left(\frac{\phi_{j}^{-1}(t)}{r}\right)^{\beta_{1,\phi_{j}}/2}$$

Putting I_1 and I_2 together, we obtain (4.5).

Next we estimate $r_*(t)$ from above and below since they are needed in steps (iii) and (iv). We first consider the lower bound for $r_*(t)$. By (1.10), (1.12) and (1.13), we have

$$\exp\left(\frac{2c_{14}r}{\beta_{1,\phi_j}\bar{\phi}_c^{-1}(t/r)}\right) \le \exp\left(c_{20}\left(\frac{r}{\phi_c^{-1}(t)}\right)^{\beta_{2,\phi_c}/(\beta_{1,\phi_c}-1)}\right).$$
(4.8)

Hence, (4.6) holds if

$$\exp\left(c_{20}\left(\frac{r}{\phi_c^{-1}(t)}\right)^{\beta_{2,\phi_c}/(\beta_{1,\phi_c}-1)}\right) \le \frac{c_{17}r}{\phi_j^{-1}(t)}, \quad C_0\phi_c^{-1}(t) \le r \le r_*(t);$$

namely,

$$\log c_{17} + \log \frac{r}{\phi_c^{-1}(t)} + \log \frac{\phi_c^{-1}(t)}{\phi_j^{-1}(t)} \ge c_{20} \left(\frac{r}{\phi_c^{-1}(t)}\right)^{\beta_{2,\phi_c}/(\beta_{1,\phi_c}-1)}$$

holds for all $C_0 \phi_c^{-1}(t) \leq r \leq r_*(t)$. Hence, we have

$$r_*(t) \ge C_1 \phi_c^{-1}(t) \log^{(\beta_{1,\phi_c}-1)/\beta_{2,\phi_c}}(\phi_c^{-1}(t)/\phi_j^{-1}(t))$$

for some constant $C_1 > 0$ which is independent of t. For the upper bound of $r_*(t)$, similar to the argument for (4.8), we have

$$\exp\left(\frac{2c_{14}r}{\beta_{1,\phi_j}\bar{\phi}_c^{-1}(t/r)}\right) \ge \exp\left(c_{21}\left(\frac{r}{\phi_c^{-1}(t)}\right)^{\beta_{1,\phi_c}/(\beta_{2,\phi_c}-1)}\right).$$

Hence, we have

$$r_*(t) \le C_2 \phi_c^{-1}(t) \log^{(\beta_{2,\phi_c}-1)/\beta_{1,\phi_c}}(\phi_c^{-1}(t)/\phi_j^{-1}(t)),$$

where $C_2 > 0$ is also independent of t.

(iii) Suppose that $r \geq r^*(t) := C_3 \phi_c^{-1}(t) \log^N(\phi_c^{-1}(t)/\phi_j^{-1}(t))$, where $N \in \mathbb{N}$ is given in Lemma 4.7, and $C_3 > 0$ is determined later. According to Lemma 4.7, there exist constants a, c > 0 and $N \in \mathbb{N}$ such that for all $x, y \in M_0$ and $0 < t \leq 1$ with $d(x, y) \geq r^*(t)$,

$$\begin{split} p(t,x,y) &\leq \frac{ct}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{c}{V(x,\phi_c^{-1}(t))} \exp\left(-\frac{ad(x,y)^{1/N}}{\phi_c^{-1}(t)^{1/N}}\right) \\ &\leq \frac{ct}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{c_1}{V(x,d(x,y))} \left(\frac{d(x,y)}{\phi_c^{-1}(t)}\right)^{d_2} \exp\left(-\frac{ad(x,y)^{1/N}}{\phi_c^{-1}(t)^{1/N}}\right) \\ &\leq \frac{ct}{V(x,d(x,y))\phi_j(d(x,y))} + \frac{c_2}{V(x,d(x,y))} \exp\left(-\frac{a}{2}\frac{d(x,y)^{1/N}}{\phi_c^{-1}(t)^{1/N}}\right), \end{split}$$

where in the second inequality we used VD and the last inequality follows from the fact that

$$r^{d_2} \le c_3 e^{ar^{1/N}/2}, \quad r \ge c_4 > 0.$$

Without loss of generality, we may and do assume that $N \ge (\beta_{2,\phi_c} - 1)/\beta_{1,\phi_c}$. In particular, $r^*(t) \ge r_*(t)$ by the upper bond for $r_*(t)$ mentioned at the end of step (ii) and also by choosing $C_3 > 0$ large enough in the definition of $r^*(t)$ if necessary.

Next, suppose that there is a constant $C_3 > 0$ in the definition of $r^*(t)$ such that

$$\exp\left(\frac{a}{2}\frac{r^{1/N}}{\phi_c^{-1}(t)^{1/N}}\right) \ge \frac{c_5\phi_j(r)}{t}, \quad r \ge r^*(t)$$
(4.9)

holds for some $c_5 > 0$. Then, for all $x, y \in M_0$ and $0 < t \le 1$ with $d(x, y) \ge r^*(t)$, we have

$$p(t, x, y) \le \frac{c_6 t}{V(x, d(x, y))\phi_j(d(x, y))}$$

Hence, by (1.10) and Lemma 2.1, for all $x \in M_0$, $t \in (0, 1]$ and $r \ge r^*(t)$,

$$\int_{B(x,r)^c} p(t,x,y)\,\mu(dy) \le c_6 \int_{B(x,r)^c} \frac{t}{V(x,d(x,y))\phi_j(d(x,y))}\,\mu(dy) \le \frac{c_7t}{\phi_j(r)} \le c_8 \left(\frac{\phi_j^{-1}(t)}{r}\right)^{\beta_{1,\phi_j}},$$

proving (4.5).

Finally, we verify that (4.9) indeed holds by using the idea of the argument for the lower bound of $r_*(t)$ in the end of step (ii). By (1.10),

$$\frac{c_5\phi_j(r)}{t} \le c_9 \left(\frac{r}{\phi_j^{-1}(t)}\right)^{\beta_{2,\phi_j}}$$

Hence (4.9) is a consequence of the following inequality

$$\exp\left(\frac{a}{2}\frac{r^{1/N}}{\phi_c^{-1}(t)^{1/N}}\right) \ge c_9\left(\frac{r}{\phi_j^{-1}(t)}\right)^{\beta_{2,\phi_j}}, \quad r \ge r^*(t);$$

that is,

$$\frac{a}{2} \left(\frac{r}{\phi_c^{-1}(t)}\right)^{1/N} \ge \log c_9 + \beta_{2,\phi_j} \log \frac{r}{\phi_c^{-1}(t)} + \beta_{2,\phi_j} \log \frac{\phi_c^{-1}(t)}{\phi_j^{-1}(t)}, \quad r \ge r^*(t).$$

The above inequality clearly is true by a suitable choice of $C_3 > 0$ so that (4.9) holds true.

(iv) Let $r \in [r_*(t), r^*(t)]$. For any $x \in M_0$, $0 < t \le 1$ and $r \in [r_*(t), r^*(t)]$, we find by the conclusion in step (ii) that

$$\begin{split} \int_{B(x,r)^c} p(t,x,y) \, \mu(dy) &\leq \int_{B(x,r_*(t))^c} p(t,x,y) \, \mu(dy) \\ &\leq c_1 \left(\frac{\phi_j^{-1}(t)}{r_*(t)}\right)^{\beta_{1,\phi_j}/2} + c_1 \exp\left(-c_2 \frac{r_*(t)}{\bar{\phi}_c^{-1}(t/r_*(t))}\right) \\ &=: I_1 + I_2. \end{split}$$

It follows from $r_*(t) \ge C_1 \phi_c^{-1}(t) \log^{(\beta_{1,\phi_c}-1)/\beta_{2,\phi_c}}(\phi_c^{-1}(t)/\phi_j^{-1}(t))$ and $\phi_j^{-1}(t) \le \phi_c^{-1}(t)$ that

$$I_{1} \leq c_{3} \left(\frac{\phi_{j}^{-1}(t)}{\phi_{c}^{-1}(t) \log^{(\beta_{1,\phi_{c}}-1)/\beta_{2,\phi_{c}}}(\phi_{c}^{-1}(t)/\phi_{j}^{-1}(t))}} \right)^{\beta_{1,\phi_{j}}/2} \\ \leq c_{3} \left(\frac{\phi_{j}^{-1}(t)}{\phi_{c}^{-1}(t) \log^{N}(\phi_{c}^{-1}(t)/\phi_{j}^{-1}(t))}} \right)^{((\beta_{1,\phi_{c}}-1)\beta_{1,\phi_{j}})/(2N\beta_{2,\phi_{c}})} \leq c_{4} \left(\frac{\phi_{j}^{-1}(t)}{r} \right)^{((\beta_{1,\phi_{c}}-1)\beta_{1,\phi_{j}})/(2N\beta_{2,\phi_{c}})}$$

On the other hand, without loss of generality we may and do assume that $c_2 \in (0, 1)$ in the term I_2 . By (4.7) and the fact that we can assume in (4.7) that $2c_{14} \ge \beta_{1,\phi_j}$, there is a constant $\theta \in (0, 1)$ such that

$$I_{2} \leq c_{5} \left(\frac{\phi_{j}^{-1}(t)}{r_{*}(t)}\right)^{\theta} \leq c_{6} \left(\frac{\phi_{j}^{-1}(t)}{\phi_{c}^{-1}(t)\log^{(\beta_{1,\phi_{c}}-1)/\beta_{2,\phi_{c}}}(\phi_{c}^{-1}(t)/\phi_{j}^{-1}(t))}}\right)^{\theta} \leq c_{7} \left(\frac{\phi_{j}^{-1}(t)}{\phi_{c}^{-1}(t)\log^{N}(\phi_{c}^{-1}(t)/\phi_{j}^{-1}(t))}}\right)^{(\theta(\beta_{1,\phi_{c}}-1))/(N\beta_{2,\phi_{c}})} \leq c_{8} \left(\frac{\phi_{j}^{-1}(t)}{r}\right)^{(\theta(\beta_{1,\phi_{c}}-1))/(N\beta_{2,\phi_{c}})}$$

Combining all the estimates above, we get (4.5) with

$$\eta = \min\{((\beta_{1,\phi_c} - 1)\beta_{1,\phi_j})/(2N\beta_{2,\phi_c}), (\theta(\beta_{1,\phi_c} - 1))/(N\beta_{2,\phi_c})\}.$$

This completes the proof.

The following lemma has been used in the proof above.

Lemma 4.8. For any $C_* > 0$, there exist constants $C_0, C^* > 0$ such that

(i) for any $t \in (0,1]$ and $r \ge C_0 \phi_c^{-1}(t)$, the function

$$r \mapsto F_{1,t}(r) := \frac{\exp\left(C_* r / \bar{\phi}_c^{-1}(t/r)\right)}{r / \bar{\phi}_c^{-1}(t/r)}$$

is strictly increasing.

(ii) for any $t \in (0,1]$, there is a unique $r_*(t) \in (C_0 \phi_c^{-1}(t), \infty)$ such that $F_{1,t}(r_*(t)) = F_{2,t}(r_*(t))$,

$$F_{1,t}(r) < F_{2,t}(r), \quad r \in (C_0 \phi_c^{-1}(t), r_*(t))$$

and

$$F_{1,t}(r) > F_{2,t}(r), \quad r \in (r_*(t), \infty),$$

where

$$F_{2,t}(r) = \frac{C^* \bar{\phi}_c^{-1}(t/r)}{\phi_j^{-1}(t)}.$$

Proof. (i) We know from (1.10) and (1.13) that, if $r \ge C_0 \phi_c^{-1}(t)$ with C_0 large enough, then

$$\frac{r}{\bar{\phi}_c^{-1}(t/r)} \ge c_1 \left(\frac{\phi_c(r)}{t}\right)^{1/(\beta_{2,\phi_c}-1)} \ge c_2 \left(\frac{r}{\phi_c^{-1}(t)}\right)^{\beta_{1,\phi_c}/(\beta_{2,\phi_c}-1)} \ge c_2 C_0^{\beta_{1,\phi_c}/(\beta_{2,\phi_c}-1)},$$

where the constants c_1 and c_2 are independent of C_0 . Note further that the function $r \mapsto r/\bar{\phi}_c^{-1}(t/r)$ is strictly increasing due to the strictly increasing property of the function $\bar{\phi}_c(r)$ on \mathbb{R}_+ . Then the first required assertion follows from the fact that the function $s \mapsto s^{-1}e^{C_*s}$ is strictly increasing on $[c_2C_0^{\beta_{1,\phi_c}/(\beta_{2,\phi_c}-1)},\infty)$, by choosing C_0 (depending on C_*) large enough.

(ii) We fix $t \in (0, 1]$. As seen from (i) and its proof that the function $F_{1,t}(r)$ is strictly increasing on $[C_0\phi_c^{-1}(t),\infty)$ with $F_{1,t}(\infty) = \infty$. On the other hand, due to the strictly increasing property of the function $\bar{\phi}_c(r)$ on \mathbb{R}_+ and (1.13), we know that the function $F_{2,t}(r)$ is strictly decreasing on $[C_0\phi_c^{-1}(t),\infty)$ with $F_{2,t}(C_0\phi_c^{-1}(t)) = \frac{C^*\bar{\phi}_c^{-1}(t/(C_0\phi_c^{-1}(t)))}{\phi_j^{-1}(t)}$ and $F_{2,t}(\infty) = 0$.

Furthermore, according to (1.10) and (1.13) again, we can obtain that $F_{1,t}(C_0\phi_c^{-1}(t)) \leq c_3$ and

$$F_{2,t}(C_0\phi_c^{-1}(t)) = \frac{C^*\phi_c^{-1}(t/(C_0\phi_c^{-1}(t)))}{\phi_j^{-1}(t)} \ge C^*\bar{\phi}_c^{-1}(t/(C_0\phi_c^{-1}(t))) \ge C^*c_4,$$

where c_3, c_4 are independent of C^* .

Combining with both conclusions above and taking $C^* = 2c_3/c_4$, we then prove the second desired assertion.

Remark 4.9. Assume that VD, (1.10), UHKD(ϕ), $J_{\phi,\leq}$ and E_{ϕ} hold. By considering the cases of $d(x,y) \leq \phi^{-1}(t)$ and $d(x,y) \geq \phi^{-1}(t)$ separately and using similar argument as those for Lemma 4.7 for the second case, we have

$$\frac{1}{V(x,\phi^{-1}(t))} \exp\left(-\frac{ad(x,y)^{1/N}}{\phi^{-1}(t)^{1/N}}\right) \le c_1\left(\frac{1}{V(x,\phi^{-1}(t))} \wedge \frac{t}{V(x,d(x,y))\phi(d(x,y))}\right).$$

Thus it follows from Lemma 4.7 and UHKD(ϕ) that

$$p(t, x, y) \le c_2 \left(\frac{1}{V(x, \phi^{-1}(t))} \land \frac{t}{V(x, d(x, y))\phi(d(x, y))} \right).$$
(4.10)

Clearly, compared with UHK(ϕ_c, ϕ_j), (4.10) is far from the optimality. However, inequality (4.10) is useful in the derivation of characterizations of parabolic Harnack inequalities in the next section. For our later use, we say UHK_{weak}(ϕ) holds if the heat kernel satisfies the upper bound estimates (4.10).

Proof of Theorem 1.14. The (ii) \implies (i) part follows from Theorem 4.6 and Proposition 3.2(ii). By Propositions 3.1, 3.3 and 3.5, we get (i) \implies (iii). Clearly (iii) \implies (iv) by Proposition 2.5, while (iv) \implies (ii) follows from Propositions 4.2 and 4.3. This proves the theorem. \Box

5 Characterizations of two-sided heat kernel estimates

In this section, we will establish stable characterizations of two-sided heat kernel estimates. Since we have obtained characterizations for $\text{UHK}(\phi_c, \phi_j)$ in Theorem 1.14, we will mainly be concerned with lower bound estimates for heat kernel in this section. We first need some definitions.

Definition 5.1. (i) We say that the parabolic Hölder regularity (PHR(ϕ)) holds for the Markov process X if there exist constants c > 0, $\theta \in (0, 1]$ and $\varepsilon \in (0, 1)$ such that for every $x_0 \in M$, $t_0 \ge 0$, r > 0 and for every bounded measurable function u = u(t, x) that is caloric in $Q(t_0, x_0, \phi(r), r)$, there is a properly exceptional set $\mathcal{N}_u \supset \mathcal{N}$ so that

$$|u(s,x) - u(t,y)| \le c \left(\frac{\phi^{-1}(|s-t|) + d(x,y)}{r}\right)^{\theta} \operatorname{ess\,sup\,}_{[t_0,t_0 + \phi(r)] \times M} |u| \tag{5.1}$$

for every $s, t \in (t_0 + \phi(r) - \phi(\varepsilon r), t_0 + \phi(r))$ and $x, y \in B(x_0, \varepsilon r) \setminus \mathcal{N}_u$.

(ii) We say that the *elliptic Hölder regularity* (EHR) holds for the process X, if there exist constants c > 0, $\theta \in (0, 1]$ and $\varepsilon \in (0, 1)$ such that for every $x_0 \in M$, r > 0 and for every bounded measurable function u on M that is harmonic in $B(x_0, r)$, there is a properly exceptional set $\mathcal{N}_u \supset \mathcal{N}$ so that

$$|u(x) - u(y)| \le c \left(\frac{d(x,y)}{r}\right)^{\theta} \operatorname{ess\,sup}_{M}|u|$$
(5.2)

for any $x, y \in B(x_0, \varepsilon r) \setminus \mathcal{N}_u$.

Clearly PHR(ϕ) \implies EHR. Note that in the definition of PHR(ϕ) (resp. EHR) if the inequality (5.1) (resp. (5.2)) holds for some $\varepsilon \in (0, 1)$, then it holds for all $\varepsilon \in (0, 1)$ (with possibly different constant c). We take EHR for example. For every $x_0 \in M$ and r > 0, let u be a bounded function on M such that it is harmonic in $B(x_0, r)$. Then for any $\varepsilon' \in (0, 1)$ and $x \in B(x_0, \varepsilon'r) \setminus \mathcal{N}_u$, u is harmonic on $B(x, (1 - \varepsilon')r)$. Applying (5.2) for u on $B(x, (1 - \varepsilon')r)$), we find that for any $y \in B(x_0, \varepsilon'r) \setminus \mathcal{N}_u$ with $d(x, y) \leq (1 - \varepsilon')\varepsilon r$,

$$|u(x) - u(y)| \le c \left(\frac{d(x,y)}{r}\right)^{\theta} \operatorname{ess\,sup}_{z \in M} |u(z)|.$$

This implies that for any $x, y \in B(x_0, \varepsilon' r) \setminus \mathcal{N}_u$, (5.2) holds with $c' = c \vee \frac{2}{[(1-\varepsilon')\varepsilon]^{\theta}}$.

5.1 $\operatorname{PI}(\phi) + \operatorname{J}_{\phi_j} + \operatorname{CS}(\phi) \Longrightarrow \operatorname{HK}_{-}(\phi_c, \phi_j)$

Proposition 5.2. Let $B_r = B(x_0, r)$ for some $x_0 \in M$ and r > 0. Assume that $u \in \mathcal{F}_{loc}^{B_R}$ is a bounded and superharmonic function in a ball B_R such that $u \ge 0$ on B_R . If VD, (1.10), CS(ϕ) and $J_{\phi,<}$ hold, then for any l > 0 and $0 < 2r \le R$,

$$\begin{split} &\int_{B_r} d\Gamma_c(\log(u+l),\log(u+l)) + \int_{B_r \times B_r} \left[\log\left(\frac{u(x)+l}{u(y)+l}\right) \right]^2 \, J(dx,dy) \\ &\leq \frac{c_1 V(x_0,r)}{\phi(r)} \left(1 + \frac{\phi(r)}{\phi(R)} \frac{\operatorname{Tail}_{\phi}\left(u_-;x_0,R\right)}{l} \right), \end{split}$$

where $\operatorname{Tail}_{\phi}(u_{-}; x_{0}, R)$ is the nonlocal tail of u_{-} with respect to $\phi(r)$ in $B(x_{0}, R)$ defined as (2.9), and c_{1} is a constant independent of u, x_{0}, r, R and l.

Proof. For pure jump Dirichlet forms, a similar statement is given in [CKW2, Proposition 4.11]. In the present setting, we need to take into account on Dirichlet forms with both local and non-local terms. According to $CS(\phi)$, $J_{\phi,\leq}$ and Proposition 2.8, we can choose $\varphi \in \mathcal{F}^{B_{3r/2}}$ related to $Cap(B_r, B_{3r/2})$ so that

$$\mathcal{E}(\varphi,\varphi) \le 2\operatorname{Cap}(B_r, B_{3r/2}) \le \frac{c_1 V(x_0, r)}{\phi(r)}.$$
(5.3)

Let u be a bounded superharmonic function in a ball B_R . As $\frac{\varphi^2}{u+l} \in \mathcal{F}^{B_{3r/2}}$ for any l > 0,

$$\mathcal{E}\left(u, \frac{\varphi^2}{u+l}\right) \ge 0. \tag{5.4}$$

By the proof of [CKW2, Proposition 4.11],

$$\mathcal{E}^{(j)}\left(u,\frac{\varphi^2}{u+l}\right) \leq \mathcal{E}^{(j)}(\varphi,\varphi) + \frac{c_2 V(x_0,r)}{\phi(R)l} \operatorname{Tail}_{\phi}\left(u_-;x_0,R\right) - \int_{B_r \times B_r} \left[\log\left(\frac{u(x)+l}{u(y)+l}\right)\right]^2 J(dx,dy).$$

That is,

$$\int_{B_r \times B_r} \left[\log \left(\frac{u(x)+l}{u(y)+l} \right) \right]^2 J(dx, dy) \leq \mathcal{E}^{(j)}(\varphi, \varphi) + \frac{c_2 V(x_0, r)}{\phi(R)l} \operatorname{Tail}\left(u_-; x_0, R\right) - \mathcal{E}^{(j)}\left(u, \frac{\varphi^2}{u+l}\right).$$

On the other hand, by the Leibniz and chain rules and the Cauchy-Schwarz inequality (2.1),

$$\begin{split} &\int \varphi^2 \, d\Gamma_c(\log(u+l), \log(u+l)) \\ &= -\int \varphi^2 \, d\Gamma_c\left(u+l, \frac{1}{u+l}\right) = -\int \varphi^2 \, d\Gamma_c\left(u, \frac{1}{u+l}\right) \\ &= -\int d\Gamma_c\left(u, \frac{\varphi^2}{u+l}\right) + 2\int \frac{\varphi}{u+l} \, d\Gamma_c(u, \varphi) \\ &\leq -\int d\Gamma_c\left(u, \frac{\varphi^2}{u+l}\right) + 2\int \, d\Gamma_c(\varphi, \varphi) + \frac{1}{2}\int \frac{\varphi^2}{(u+l)^2} \, d\Gamma_c(u, u) \\ &= -\int d\Gamma_c\left(u, \frac{\varphi^2}{u+l}\right) + 2\int \, d\Gamma_c(\varphi, \varphi) + \frac{1}{2}\int \varphi^2 \, d\Gamma_c(\log(u+1), \log(u+1)). \end{split}$$

Namely,

$$\int \varphi^2 \, d\Gamma_c(\log(u+l), \log(u+l)) \le -2 \int d\Gamma_c\left(u, \frac{\varphi^2}{u+l}\right) + 4 \int \, d\Gamma_c(\varphi, \varphi).$$

Hence,

$$\int_{B_r} d\Gamma_c(\log(u+l), \log(u+l)) \le 4\mathcal{E}^{(c)}(\varphi, \varphi) - 2\mathcal{E}^{(c)}\left(u, \frac{\varphi^2}{u+l}\right).$$

Putting both estimates together, we conclude that

$$\begin{split} &\int_{B_r} d\Gamma_c(\log(u+l),\log(u+l)) + \int_{B_r \times B_r} \left[\log\left(\frac{u(x)+l}{u(y)+l}\right) \right]^2 J(dx,dy) \\ &\leq \int_{B_r} d\Gamma_c(\log(u+l),\log(u+l)) + 2 \int_{B_r \times B_r} \left[\log\left(\frac{u(x)+l}{u(y)+l}\right) \right]^2 J(dx,dy) \\ &\leq 2\mathcal{E}^{(j)}(\varphi,\varphi) + 4\mathcal{E}^{(c)}(\varphi,\varphi) + \frac{2c_2V(x_0,r)}{\phi(R)l} \mathrm{Tail}_{\phi}\left(u_-;x_0,R\right) - 2\mathcal{E}\left(u,\frac{\varphi^2}{u+l}\right) \\ &\leq 4\mathcal{E}(\varphi,\varphi) + \frac{2c_2V(x_0,r)}{\phi(R)l} \mathrm{Tail}_{\phi}\left(u_-;x_0,R\right) - 2\mathcal{E}\left(u,\frac{\varphi^2}{u+l}\right) \\ &\leq \frac{c_5V(x_0,r)}{\phi(r)} \left(1 + \frac{\phi(r)}{\phi(R)} \frac{\mathrm{Tail}_{\phi}\left(u_-;x_0,R\right)}{l}\right), \end{split}$$

where in the last inequality we used (5.3) and (5.4). The proof is complete.

With Proposition 5.2, we can follow the arguments for [CKW2, Corollary 4.12 and Propisition 4.13] to obtain the following.

Proposition 5.3. Assume VD, RVD and (1.10). Then

$$\operatorname{PI}(\phi) + \operatorname{J}_{\phi,<} + \operatorname{CS}(\phi) \Longrightarrow \operatorname{EHR}.$$

Proposition 5.4. If VD, RVD, (1.10), $PI(\phi)$, $J_{\phi,\leq}$ and $CS(\phi)$ hold, then we have $NDL(\phi)$.

Proof. It follows from Proposition 4.2 that under VD, RVD and (1.10),

$$\operatorname{PI}(\phi) + \operatorname{J}_{\phi,<} + \operatorname{CS}(\phi) \Longrightarrow \operatorname{E}_{\phi},$$

where we also used the fact that $PI(\phi)$ implies $FK(\phi)$ by Proposition 3.1. With this and Proposition 5.3, the desired assertion essentially follows from the proof of [CKW2, Proposition 4.9].

The following proposition establishes the $(v) \Longrightarrow (i)$ part in Theorem 1.13.

Proposition 5.5. Suppose VD, RVD, (1.10), $PI(\phi)$, J_{ϕ_j} and $CS(\phi)$ hold. Then we have $HK_{-}(\phi_c, \phi_j)$.

Proof. According to Proposition 5.4, we have $\text{NDL}(\phi)$. In particular, for any $x, y \in M_0$ and t > 0 with $d(x, y) \leq c_0 \phi^{-1}(t)$ for some constant $c_0 > 0$,

$$p(t, x, y) \ge \frac{c_1}{V(x, \phi^{-1}(t))}.$$

On the other hand, under our assumptions, we can get from the arguments in step (ii) for the proof of [CKW1, Proposition 5.4] that for all $x, y \in M_0$ and t > 0 with $d(x, y) \ge c_0 \phi^{-1}(t)$,

$$p(t, x, y) \ge \frac{c_2 t}{V(x, d(x, y))\phi_j(d(x, y))}$$

This establishes the heat kernel lower bound (1.31) for $HK_{-}(\phi_c, \phi_j)$. The upper bound of $HK_{-}(\phi_c, \phi_j)$ follows from Proposition 3.1 and the equivalence between (i) and (iv) of Theorem 1.14.

5.2 From $HK_{-}(\phi_c, \phi_j)$ to $HK(\phi_c, \phi_j)$

To consider $\text{HK}(\phi_c, \phi_j)$, we assume in addition that (M, d, μ) is connected and satisfies the chain condition (see the end of Remark 1.12 (i)). We emphasize that the results in the previous sections hold true without this additional assumption on the state space (M, d, μ) .

Proposition 5.6. Suppose that (M, d, μ) is connected and satisfies the chain condition. Under VD and (1.10), HK₋(ϕ_c, ϕ_j) implies HK(ϕ_c, ϕ_j). In particular, if VD, RVD, (1.10), PI(ϕ), J $_{\phi_j}$ and CS(ϕ) hold, then so does HK(ϕ_c, ϕ_j).

Proof. By $\text{HK}_{-}(\phi_c, \phi_j)$ and (1.27), we only need to verify the case that $t \in (0, 1]$ and $x, y \in M_0$ with $d(x, y) \ge c_0 \phi_c^{-1}(t)$ for some constant $c_0 > 0$. The proof is based on the standard chaining argument, e.g. see the proof of [BGK, Proposition 5.2(i)].

First we assume HK₋(ϕ_c, ϕ_j) holds. Then, by Remark 1.12(ii), NL(ϕ) holds. Furthermore, in view of Remark 1.12(iii), it suffices to consider the case that $t \in [0, 1]$. Now, fix $t \in (0, 1]$ and $x, y \in M_0$. Let r = d(x, y). Since the space (M, d, μ) satisfies the chain condition, there exists a constant C > 0 such that, for any $x, y \in M$ and for any $n \in \mathbb{N}$, there exists a sequence $\{x_i\}_{i=0}^n \subset M$ such that $x_0 = x, x_n = y$ and $d(x_i, x_{i+1}) \leq Cd(x, y)/n$ for all $i = 0, 1, \dots, n-1$. In the following, we set n = m := m(t, r), where m(t, r) is defined by (1.24). Define $r_n = Cr/(3n)$. In particular, by (1.25), (1.12) and (1.10), $r_n \approx \phi_c^{-1}(t/n)$. By NL(ϕ), for all $z_i \in B(x_i, r_n)$ and $0 \leq i \leq n+1$,

$$p(t/n, z_i, z_{i+1}) \ge \frac{c_1}{V(z_i, \phi_c^{-1}(t/n))}.$$

Hence,

$$\begin{split} p(t,x,y) \\ &\geq \int_{B(z_1,r_n)} p(t/n,x,z_1) \, \mu(dz_1) \int_{B(z_2,r_n)} p(t/n,z_1,z_2) \, \mu(dz_2) \cdots \int_{B(z_{n-1},r_n)} p(t/n,z_{n-1},y) \, \mu(dz_{n-1}) \\ &\geq \frac{c_2}{V(x,\phi_c^{-1}(t/n))} c_3^n \geq \frac{c_4}{V(x,\phi_c^{-1}(t))} n^{d_1/\beta_{2,\phi_c}} c_3^n \geq \frac{c_5}{V(x,\phi_c^{-1}(t))} c_6^n, \end{split}$$

where the constants $c_3, c_6 \in (0, 1)$, and in the third inequality we used VD and (1.10). That is, we arrive at a lower bound of p(t, x, y) with the form given in (1.23), thanks to VD and (1.10)

again. This gives $HK(\phi_c, \phi_j)$ and hence the first assertion of this theorem. The last assertion follows from the first one, Proposition 5.4 and the fact that $NDL(\phi)$ implies $NL(\phi)$.

We need the following simple lemma for the proof of Theorem 1.13, which holds without the connectedness and the chain condition on the state space (M, d, μ) .

Lemma 5.7. Under VD and (1.10), UHK(ϕ_c, ϕ_j) and NL(ϕ) together imply NDL(ϕ).

Proof. We will use the ideas of the argument in [BGK, Subsection 4.1] and the proof of [BBK2, Lemma 3.2]. By carefully checking these proofs, to obtain the required assertion we only need to verify that for any $x \in M_0$, t > 0 and r > 0 with $r \simeq \phi^{-1}(t)$,

$$\sup_{0 \le s \le t} \sup_{y \in B(x,r), z \in B(x,2r)^c} p(s,y,z) \le \frac{c_1}{V(x,\phi^{-1}(t))}.$$

According to VD, (1.10) and UHK (ϕ_c, ϕ_j) , for any $x \in M_0$, $t \ge 1$ and r > 0 with $r \asymp \phi^{-1}(t) = \phi_j^{-1}(t)$,

$$\sup_{0 \le s \le t} \sup_{y \in B(x,r), z \in B(x,2r)^c} p(s,y,z) \le \sup_{y \in B(x,r)} \frac{c_2 t}{V(y,r)\phi_j(r)} \le \frac{c_3}{V(x,r)} \le \frac{c_4}{V(x,\phi^{-1}(t))}$$

On the other hand, also by VD, (1.10) and UHK (ϕ_c, ϕ_j) , for any $x \in M_0$, $t \in (0, 1]$ and r > 0with $r \simeq \phi^{-1}(t) = \phi_c^{-1}(t)$,

$$\begin{split} \sup_{0 \le s \le t} \sup_{y \in B(x,r), z \in B(x,2r)^c} p(s,y,z) \\ &\le \sup_{y \in B(x,r)} \frac{c_5 t}{V(y,r)\phi_j(r)} + \sup_{0 \le s \le t} \sup_{y \in B(x,r)} \frac{c_5}{V(y,\phi_c^{-1}(s))} \exp\left(-c_6 \frac{r}{\phi_c(s/r)}\right) \\ &\le \frac{c_7 t}{V(x,r)\phi_c(r)} + \frac{c_7}{V(x,\phi_c^{-1}(t))} \sup_{0 \le s \le c_*\phi_c(r)} \left(\frac{r}{\phi_c^{-1}(s)}\right)^{d_2} \exp\left(-c_6 \frac{r}{\phi_c^{-1}(s/r)}\right) \\ &\le \frac{c_7 t}{V(x,r)\phi_c(r)} + \frac{c_8}{V(x,\phi_c^{-1}(t))} \sup_{0 \le s \le c_*\phi_c(r)} \left(\frac{\phi_c(r)}{s}\right)^{d_2/\beta_{1,\phi_c}} \exp\left(-c_9 \left(\frac{\phi_c(r)}{s}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\le \frac{c_7 t}{V(x,r)\phi_c(r)} + \frac{c_{10}}{V(x,\phi_c^{-1}(t))} \sup_{0 \le s \le c_*\phi_c(r)} \exp\left(-c_{11} \left(\frac{\phi_c(r)}{s}\right)^{1/(\beta_{2,\phi_c}-1)}\right) \\ &\le \frac{c_{12}}{V(x,\phi_c^{-1}(t))}, \end{split}$$

where in the second inequality we used the fact that $\phi_c(r) \leq c_0 \phi_j(r)$ for all $r \in (0, r_0]$ and some $r_0 > 0$, in the third inequality we applied (1.12) and (1.13), and the fourth inequality follows from the following elementary inequality:

$$r^{d_2/\beta_{1,\phi_c}} \le c_{13} \exp\left(\frac{c_9}{2}r^{1/(\beta_{2,\phi_c}-1)}\right), \quad r \ge c_{14} > 0.$$

Combining both conclusions above, we get the desired conclusion.

We are now in a position to give the

Proof of Theorem 1.13. It is obvious that (i) \Longrightarrow (ii), thanks to Proposition 3.3. (ii) \Longrightarrow (iii) follows from Lemma 5.7. By Proposition 4.1, under VD and (1.10), NDL(ϕ) implies PI(ϕ), and NDL(ϕ) also implies E_{ϕ} under the additional assumption RVD. With these at hand, we have (iii) \Longrightarrow (iv) by the (ii) \Longrightarrow (iii) part of Theorem 1.14 and Proposition 3.5. (iv) \Longrightarrow (v) has been proved in Proposition 2.5. Furthermore, by Proposition 5.5, (v) \Longrightarrow (i). When the space (M, d, μ) is connected and satisfies the chain condition, (v) \Longrightarrow (vi) has been proven in Proposition 5.6. Clearly, (vi) \Longrightarrow (i). This completes the proof of the theorem.

6 Characterizations of parabolic Harnack inequalities

The goal of this section is to present three different characterizations of parabolic Harnack inequalities, see Theorems 6.3, 6.4 and 6.5.

By the arguments in [CKW2, Section 3.1], we have the following consequences of $PHI(\phi)$. Note that, though [CKW2] is concerned with pure jump non-local Dirichlet forms, the arguments in [CKW2, Section 3.1] works for general symmetric Dirichlet forms (\mathcal{E}, \mathcal{F}) under the present setting.

Proposition 6.1. Assume that VD, (1.10) and PHI(ϕ) hold. Then UHKD(ϕ) and NDL(ϕ), as well as UJS, hold true. Consequently, PI(ϕ) and $E_{\phi,\geq}$ hold, and $X := \{X_t\}_{t\geq 0}$ is conservative. If furthermore RVD is satisfied, then we also have FK(ϕ) and $E_{\phi,\leq}$ (and so E_{ϕ}).

Proof. See [CKW2, Propositions 3.1, 3.2, 3.5 and 3.6] for the proof.

We point out that $PHI(\phi)$ alone can not guarantee $J_{\phi_j,\leq}$. Indeed, following the argument of [CKW2, Corollary 3.4], under VD, (1.10), UJS and NDL(ϕ), since $\phi(r) \leq \phi_j(r)$ for all r > 0, we can only obtain $J_{\phi,\leq}$, which is weaker than $J_{\phi_j,\leq}$. See Example 7.1 for a concrete counterexample. In spite of this, we still have the following statement.

Proposition 6.2. Assume that VD, (1.10), NDL(ϕ), $E_{\phi,\leq}$ and $J_{\phi,\leq}$ hold. For every $\delta \in (0,1)$, there exist positive constants C > 0 and $\gamma \in (0,1]$, where γ is independent of δ , so that for any bounded caloric function u in $Q(t_0, x_0, \phi(r), r)$, there is a properly exceptional set $\mathcal{N}_u \supset \mathcal{N}$ such that

$$|u(s,x) - u(t,y)| \le C \left(\frac{\phi^{-1}(|s-t|) + d(x,y)}{r}\right)^{\gamma} \operatorname{ess\,sup}_{[t_0,t_0+\phi(r)] \times M} |u|$$

for every $s, t \in (t_0 + \phi(r) - \phi(\delta r), t_0 + \phi(r))$ and $x, y \in B(x_0, \delta r) \setminus \mathcal{N}_u$. In other words, under VD and (1.10), NDL(ϕ) + E $_{\phi,\leq}$ + J $_{\phi,\leq}$ imply PHR(ϕ) and EHR. In particular, PHI(ϕ) implies PHR(ϕ) and EHR.

Proof. The proof is the same as that of [CKW2, Proposition 3.8], so it is omitted. \Box

We next present characterizations of $PHI(\phi)$. Recall that in Remark 4.9 upper bound estimate (4.10) of the heat kernel p(t, x, y) is named by $UHK_{weak}(\phi)$.

Theorem 6.3. Assume that μ and ϕ satisfy VD, RVD and (1.10) respectively. Then the following holds

$$PHI(\phi) \iff UHK_{weak}(\phi) + NDL(\phi) + UJS.$$

Proof. According to Proposition 6.1, we have the assertion that

$$PHI(\phi) \Longrightarrow UHKD(\phi) + NDL(\phi) + UJS + E_{\phi} + J_{\phi,\leq},$$

where we note that RVD is only used to prove $PHI(\phi) \Longrightarrow E_{\phi,\leq}$. Then by Remark 4.9, we get

$$PHI(\phi) \Longrightarrow UHK_{weak}(\phi) + NDL(\phi) + UJS.$$

For $\text{UHK}_{weak}(\phi) + \text{NDL}(\phi) + \text{UJS} \implies \text{PHI}(\phi)$, we can follow most of the arguments in [CKW2, Subsection 4.1]. One different point is that in the proof of [CKW2, Lemma 4.1] (see the arXiv version of the paper [CKW2]), we need to verify that under VD and (1.10), $\text{UHK}_{weak}(\phi)$ implies the following: for any $x, y \in M_0$ and t > 0 with $d(x, y) \approx \phi^{-1}(t)$,

$$p(t, x, y) \le \frac{c_1}{V(x, \phi^{-1}(t))}.$$

Yet, this inequality is a direct consequence of (4.10). The proof is complete.

The next characterization of $PHI(\phi)$ involves the property of exit times E_{ϕ} .

Theorem 6.4. Assume that μ and ϕ satisfy VD, RVD and (1.10) respectively. Then the following hold

$$PHI(\phi) \iff PHR(\phi) + E_{\phi} + UJS + J_{\phi,\leq}$$
$$\iff EHR + E_{\phi} + UJS + J_{\phi,\leq}.$$

Proof. According to Propositions 6.2 and 6.1, we have

$$PHI(\phi) \Longrightarrow PHR(\phi) + E_{\phi} + UJS + J_{\phi,\leq} \Longrightarrow EHR + E_{\phi} + UJS + J_{\phi,\leq}.$$

For EHR + E_{ϕ} + UJS + $J_{\phi,\leq} \Longrightarrow$ PHI(ϕ), we mainly follow the arguments in [CKW2, Subsection 4.2]. In particular, the proof of [CKW2, Proposition 4.9] yields that EHR together with E_{ϕ} imply NDL(ϕ). Then according to the argument of [CKW2, Proposition 3.5], FK(ϕ) holds, thanks to RVD. By Proposition 4.3, under FK(ϕ), E_{ϕ} and $J_{\phi,\leq}$, UHKD(ϕ) holds true. This along with Remark 4.9 yields that UHK_{weak}(ϕ) holds. Combining all these with Theorem 6.3, we have

$$\operatorname{EHR} + \operatorname{E}_{\phi} + \operatorname{UJS} + \operatorname{J}_{\phi,\leq} \Longrightarrow \operatorname{PHI}(\phi).$$

The proof is complete.

Finally, we turn to the stable analytic characterization of $PHI(\phi)$.

Theorem 6.5. Assume that μ and ϕ satisfy VD, RVD and (1.10) respectively. Then the following hold

$$PHI(\phi) \iff PI(\phi) + J_{\phi,\leq} + Gcap(\phi) + UJS$$
$$\iff PI(\phi) + J_{\phi,\leq} + CS(\phi) + UJS.$$

Proof. By Proposition 4.2, Theorem 6.4 and Proposition 5.3,

$$\mathrm{PI}(\phi) + \mathrm{J}_{\phi,<} + \mathrm{CS}(\phi) + \mathrm{UJS} \Longrightarrow \mathrm{PHI}(\phi),$$

where we used Proposition 3.1 that $PI(\phi)$ implies $FK(\phi)$ under the additional assumption RVD.

It follows from Proposition 2.5 that

$$PI(\phi) + J_{\phi,<} + Gcap(\phi) + UJS \Longrightarrow PI(\phi) + J_{\phi,<} + CS(\phi) + UJS.$$

Finally, according to Proposition 6.1 and Theorem 6.3, we have

$$PHI(\phi) \Longrightarrow NDL(\phi) + UJS + E_{\phi} + J_{\phi,\leq} + UHK_{weak}(\phi).$$

This in particular implies that the Dirichlet form $(\mathcal{E}, \mathcal{F})$ is conservative by Proposition 3.2(ii). Furthermore, by the argument of [CKW2, Proposition 3.5], NDL $(\phi) \Longrightarrow$ PI (ϕ) . We note that, from the proof of Lemma 3.4, we can see that UHK_{weak} (ϕ) along with the fact $(\mathcal{E}, \mathcal{F})$ is conservative implies $\text{EP}_{\phi,\leq}$, which in turn gives us $\text{Gcap}(\phi)$ by Proposition 2.4. Thus, we prove that

$$PHI(\phi) \Longrightarrow PI(\phi) + J_{\phi,\leq} + Gcap(\phi) + UJS.$$

The proof is complete.

Proof of Theorem 1.17. As noted earlier, the equivalence between (i) and (ii) of Theorem 1.17 has been proved in Theorem 6.3, while the equivalence between (i), (iii) and (iv) has been established in Theorem 6.4. The equivalence between (i), (v) and (vi) follows from Theorem 6.3, Theorem 6.4 and Theorem 6.5. The last assertion of Theorem 1.17 follows from the equivalence between (i) and (v) of Theorem 1.17 and from Theorem 1.13.

7 Examples/Applications

In this section, we give some examples/applications of our results.

Example 7.1. (PHI(ϕ) alone does not imply $J_{\phi_i,<}$) Let $M = \mathbb{R}^d$, and

$$J(x,y) \asymp \begin{cases} \frac{1}{|x-y|^{d+\alpha}} & |x-y| \le 1;\\ \frac{1}{|x-y|^{d+\beta}} & |x-y| \ge 1; \end{cases}$$

where $\alpha, \beta \in (0, 2)$. We consider the following regular Dirichlet form

$$\mathcal{E}(f,g) = \int_{\mathbb{R}^d} \nabla f(x) \cdot A(x) \nabla g(x) \, dx + \iint_{\mathbb{R}^d \times \mathbb{R}^d} (f(x) - f(y))(g(x) - g(y)) J(x,y) \, dx \, dy$$

and $\mathcal{F} = \overline{C_1(\mathbb{R}^d)}^{\mathcal{E}_1}$, where A(x) is a measurable $d \times d$ matrix-valued function on \mathbb{R}^d that is uniformly elliptic and bounded. It has been proven in [CK3, Theorem 1.4] that $\mathrm{HK}(\phi_c, \phi_j)$ holds with $\phi_c(r) = r^2$ and $\phi_j(r) = r^{\alpha} \mathbf{1}_{\{r \leq 1\}} + r^{\beta} \mathbf{1}_{\{r \geq 1\}}$. Hence by (1.36), PHI(ϕ) holds with

$$\phi(r) = \phi_c(r) \land \phi_i(r) = r^\beta \land r^2.$$

Since $PHI(\phi)$ holds regardless of the choice of $\alpha \in (0, 2)$, $PHI(\phi)$ alone can not imply the upper bound of the jumping kernel.

Example 1.1 (continued) Here we provide proof of Example 1.1.

First, consider a reflected Brownian motion $\{B_t\}_{t\geq 0}$ on U. It is known that its heat kernel enjoys Aronson-type Gaussian bounds, see for instance [GSC, Theorem 2.31]. (In fact, as discussed in [GSC, Theorem 2.31], similar results hold for reflected Brownian motions on inner uniform domains in Harnack-type Dirichlet spaces, so the results in this example can be extended to that framework.) In particular, according to [GSC, Theorem 2.31], we know that the following Poincaré inequality for the strongly local part of $(\mathcal{E}, W^{1,2}(U))$ given by (1.6), i.e., there exist constants $C_1 > 0$ and $\kappa_1 \geq 1$ such that for any ball $B_r := B(x, r)$ with $x \in U$ and r > 0 and for any $f \in W^{1,2}(U) \cap B_b(U)$,

$$\int_{B_r} (f - \overline{f}_{B_r})^2 \, d\mu \le C_1 r^2 \left(\int_{B_{\kappa_1 r}} \nabla f(z) \cdot A(z) \nabla f(z) \, dz \right).$$

On the other hand, for $(\mathcal{E}, W^{1,2}(U))$ given by (1.6), it is obvious that condition J_{ϕ_j} holds with $\phi_j(r) = r^{\alpha}$, which in turn yields that there exist constants $C_2 > 0$ and $\kappa_2 \ge 1$ such that for any ball $B_r = B(x, r)$ with $x \in U$ and r > 0 and for any $f \in W^{1,2}(U) \cap B_b(U)$,

$$\int_{B_r} (f - \overline{f}_{B_r})^2 \, d\mu \le C_2 r^{\alpha} \left(\int_{B_{\kappa_2 r}} \int_{B_{\kappa_2 r}} \frac{(f(y) - f(z))^2}{d(y, z)^{d + \alpha}} c(y, z) \, dy \, dz \right)$$

Hence, we have, for any ball $B_r = B(x, r)$ with $x \in U$ and r > 0 and for any $f \in W^{1,2}(U) \cap B_b(U)$,

$$\begin{split} &\int_{B_r} (f - \overline{f}_{B_r})^2 \, d\mu \\ &\leq C_0 (r^2 \wedge r^\alpha) \left(\int_{B_{\kappa_0 r}} \nabla f(z) \cdot A(z) \nabla f(z) \, dz + \int_{B_{\kappa_0 r}} \int_{B_{\kappa_0 r}} \frac{(f(y) - f(z))^2}{d(y, z)^{d + \alpha}} c(y, z) \, dy \, dz \right) \end{split}$$

with $C_0 = C_1 \vee C_2$ and $\kappa_0 = \kappa_1 \vee \kappa_2$. That is, $\operatorname{PI}(\phi)$ holds with $\phi(r) = r^2 \wedge r^{\alpha}$. Note further that, in the present setting $\operatorname{CS}(\phi)$ holds trivially, as mentioned in Remark 1.7 (iii). Therefore, it immediately follows from our stability theorem (Theorem 1.13) that the heat kernel for the Dirichlet form (1.6) enjoys the estimates $\operatorname{HK}(\phi_c, \phi_j)$, hence $\operatorname{PHI}(\phi)$ as well.

An alternative way to prove Example 1.1 is to use subordination of Brownian motion and use the transferring method to be discussed below.

The stability results in Theorems 1.13, 1.14 and 1.17 allow us to obtain heat kernel estimates and parabolic Harnack inequalities for a large class of symmetric diffusions with jumps using "transferring method"; that is, by first establishing heat kernel estimates and parabolic Harnack inequalities for a particular diffusion with jumps with nice jumping kernel J(x, y), we then use Theorems 1.13, 1.14 and 1.17 to obtain heat kernel estimates and parabolic Harnack inequalities for other symmetric jump processes whose strongly local parts along the diagonal are comparable to that of the original process and whose jumping kernels are comparable to J(x, y). Examples for the pure jump case have been given in [CKW1, Section 6.1] and [CKW2, Section 5].

In the following, we illustrate this method for symmetric Dirichlet forms (1.1) on a *d*-sets *M* on which there exists a diffusion whose heat kernel enjoys (sub-)Gaussian estimates as in (7.1).

Example 7.2. (Diffusion with jumps on *d*-set.) Let (M, d, μ) be an Alfhors *d*-regular set. Suppose that there is a μ -symmetric diffusion $\{Z_t; t \ge 0; \mathbb{P}_x, x \in M\}$ on M such that it

has a transition density function q(t, x, y) with respect to the measure μ that has the following two-sided estimates:

$$q(t,x,y) \asymp t^{-d/\beta} \exp\left(-\left(\frac{d(x,y)^{\beta}}{t}\right)^{1/(\beta-1)}\right), \quad t > 0, x, y \in M$$
(7.1)

for some $\beta \geq 2$. Denote by $(\bar{\mathcal{E}}, \bar{\mathcal{F}})$ the corresponding Dirichlet form. A prototype is a Brownian motion on the *D*-dimensional unbounded Sierpiński gasket; see for instance [BP]. In this case, $d = \log(D+1)/\log 2$ is the Hausdorff dimension of the gasket, and $\beta = \log(D+3)/\log 2$ is called the walk dimension in (7.1).

Take any $\alpha \in (0,\beta)$, and a symmetric strongly local regular Dirichlet form $(\mathcal{E}^{(c)}, \bar{\mathcal{F}})$ in $L^2(M;\mu)$ with the property that $\mathcal{E}^{(c)}(f,f) \simeq \bar{\mathcal{E}}(f,f)$ for all $f \in \bar{\mathcal{F}}$. Consider the following regular Dirichlet form $(\mathcal{E}, \bar{\mathcal{F}})$ in $L^2(M;\mu)$ defined by

$$\mathcal{E}(u,v) = \mathcal{E}^{(c)}(u,v) + \int_M \int_M (u(x) - u(y))(v(x) - v(y)) \frac{c(x,y)}{d(x,y)^{d+\alpha}} \,\mu(dx) \,\mu(dy), \tag{7.2}$$

where $c(\cdot, \cdot)$ is a symmetric measurable function on $M \times M$ that is bounded between two positive constants. Define $\phi_c(r) = r^{\beta}$, $\phi_j(r) = r^{\alpha}$ and $\phi(r) = \phi_c(r) \wedge \phi_j(r) = r^{\beta} \wedge r^{\alpha}$. We claim that $(\mathcal{E}, \bar{\mathcal{F}})$ enjoys HK (ϕ_c, ϕ_j) and PHI (ϕ) .

Below we prove this using subordination and the transferring method. First, let $\{\xi_t\}_{t\geq 0}$ be a γ -stable subordinator with non-zero drift, independent of $\{Z_t\}_{t\geq 0}$, such that its Laplace exponent $\bar{\phi}$ is given by

$$\mathbb{E}[\exp(-\lambda\xi_t)] = \exp(-t\bar{\phi}(\lambda)), \quad \lambda, t > 0$$
(7.3)

with $\bar{\phi}(\lambda) = \lambda + \lambda^{\gamma}$ for $\gamma := \alpha/\beta \in (0, 1)$. The process $\{X_t\}_{t\geq 0}$ defined by $X_t = Z_{\xi_t}$ for any $t \geq 0$ is called a γ -stable subordinated process with drift. Let $\{\eta_t(u) : t > 0, u \geq 0\}$ be the transition density of $\{\xi_t\}_{t\geq 0}$, and $\nu(z)$ be the Lévy density of $\{\xi_t\}_{t\geq 0}$. Let p(t, x, y) be the heat kernel of $\{X_t\}_{t\geq 0}$, and J(x, y) be the jumping density of $\{X_t\}_{t\geq 0}$. Then, it is known that

$$p(t, x, y) = \int_0^\infty q(u, x, y)\eta_t(u) \, du, \quad t > 0, x, y \in M,$$
$$J(x, y) = \int_0^\infty q(u, x, y)\nu(u) \, du, \quad x, y \in M.$$

By [CKW1, Section 6.1], we know that

$$J(x,y) \simeq \frac{1}{d(x,y)^{d+\beta\gamma}} = \frac{1}{d(x,y)^{d+\alpha}}, \quad x,y \in M.$$

Namely, J_{ϕ_i} holds with $\phi_i(r) = r^{\alpha}$.

On the other hand, by the definition of $\{\xi_t\}_{t\geq 0}$, $\eta_t(u) = \eta_t^{(\gamma)}(u-t)$, where $\eta_t^{(\gamma)}(u)$ corresponds to the transition density of the standard γ -stable subordinator (without drift). Then,

$$\begin{split} p(t,x,y) &= \int_0^\infty q(u,x,y) \eta_t^{(\gamma)}(u-t) \, du = \int_t^\infty q(u,x,y) \eta_t^{(\gamma)}(u-t) \, du \\ &= \int_0^\infty q(u+t,x,y) \eta_t^{(\gamma)}(u) \, du = \int_M q(t,x,z) \int_0^\infty q(u,z,y) \eta_t^{(\gamma)}(u) \, du \, \mu(dz) \\ &=: \int_M q(t,x,z) q^{(\gamma)}(t,z,y) \, \mu(dz), \end{split}$$

where $\{q^{(\gamma)}(t, x, y) : t > 0, x, y \in M\}$ is the heat kernel corresponding to the standard γ -stable subordination of the process Z. In particular, according to [CKW1, Section 6.1],

$$q^{(\gamma)}(t,x,y) \simeq t^{-d/\alpha} \wedge \frac{t}{d(x,y)^{d+\alpha}}, \quad t > 0, x, y \in M.$$

Furthermore, by standard calculations (see the proof of [SV, Theorem 2.13] for the case that $\beta = 2, \ \gamma = \alpha/2$ and $M = \mathbb{R}^d$), one can check that p(t, x, y) enjoys the form of (1.30) with $V(x, r) \simeq r^d, \ \phi_c(r) = r^\beta$ and $\phi_j(r) = r^\alpha$. This is, the heat kernel for $\{X_t\}_{t\geq 0}$ enjoys $\operatorname{HK}(\phi_c, \phi_j)$ (hence $\operatorname{PHI}(\phi)$ as well) with $\phi(r) = r^\beta \wedge r^\alpha$.

Let $Y = \{Y_t, t \ge 0; \mathbb{P}_x^Y, x \in M\}$ be the Hunt process associated with the regular Dirichlet form $(\mathcal{E}, \bar{\mathcal{F}})$ in $L^2(M; \mu)$ given by (7.2). Clearly, J_{ϕ_j} holds and so does $\operatorname{PI}(\phi)$ for $(\mathcal{E}, \bar{\mathcal{F}})$ by the same argument as in Example 1.1. Note that if two local Dirichlet forms are comparable, then their energy measures are also comparable (see for instance [FOT, Section 3.2]). Hence we see that $\operatorname{CS}(\phi)$ holds for the process Y because it holds for $\{X_t\}_{t\ge 0}$. Now thanks to Theorem 1.13, we obtain $\operatorname{HK}(\phi_c, \phi_j)$, and consequently $\operatorname{PHI}(\phi)$ for the Hunt process Y, or equivalently, for the Dirichlet form $(\mathcal{E}, \bar{\mathcal{F}})$ in $L^2(M; \mu)$.

It is desirable to prove Example 7.2 directly (i.e. without using the subordination) from the stability results of the diffusion and the jump process as we did in Example 1.1. However, it is highly non-trivial to verify $CS(\phi)$ when $\beta > 2$. Indeed, in that case the cut-off functions for the diffusion and the jump process may be different and we cannot simply sum up two forms.

We end this section a remark on two possible extensions of Example 7.2.

Remark 7.3. (i) We can start from more general symmetric diffusions on general measure metric spaces. For example, let (M, d, μ) be a metric measure space as in the setting of this paper that is connected and also satisfies VD and the chain condition. Assume that there is a μ -symmetric conservative diffusion process $\{Z_t\}$ whose heat kernel enjoys (1.22). This includes symmetric diffusions on certain fractal-like manifolds; see [CKW1, Section 6.1].

(ii) We can also consider more general subordinator. For instance, let $\{\xi_t\}_{t\geq 0}$ be a subordinator with non-zero drift such that its Laplace exponent $\bar{\phi}$ defined by (7.3) has the following form

$$\bar{\phi}(\lambda) = b\lambda + \phi_0(\lambda),$$

where b > 0, and $\phi_0(r)$ satisfies (1.15) with $\beta_{1,\phi_0}, \beta_{2,\phi_0} \in (0,1)$, and the associated Lévy measure $\nu(dz)$ of $\phi_0(r)$ has a density function $\nu(z)$ with respect to the Lebesgue measure such that the function $t \mapsto t\nu(t)$ is non-increasing on $(0, \infty)$. Under these assumptions, two-sided estimates for the transition density of the subordinator corresponding to $\phi_0(r)$ recently have been obtained in [CKKW1, Theorem 4.4]. Thus, with aid of [CKKW1, Theorem 4.4], the argument of Example 7.2 could be workable for this larger class of subordinators.

Acknowledgements. The research of Zhen-Qing Chen is partially supported by Simons Foundation Grant 520542, a Victor Klee Faculty Fellowship at UW, and NNSFC grant 11731009. The research of Takashi Kumagai is supported by JSPS KAKENHI Grant Number JP17H01093 and by the Alexander von Humboldt Foundation. The research of Jian Wang is supported by the National Natural Science Foundation of China (No. 11831014), the Program for Probability and Statistics: Theory and Application (No. IRTL1704) and the Program for Innovative Research Team in Science and Technology in Fujian Province University (IRTSTFJ).

References

- [AB] S. Andres and M.T. Barlow. Energy inequalities for cutoff-functions and some applications. J. Reine Angew. Math. 699 (2015), 183–215.
- [B] M.T. Barlow. Diffusions on fractals. In: Lectures on Probability Theory and Statistics, Ecole d'Éte de Probabilités de Saint-Flour XXV - 1995, 1–121. Lect. Notes Math. 1690, Springer 1998.
- [BB1] M.T. Barlow and R.F. Bass. Brownian motion and harmonic analysis on Sierpiński carpets. Canad. J. Math. 51 (1999), 673–744.
- [BB2] M.T. Barlow and R.F. Bass. Stability of parabolic Harnack inequalities. Trans. Amer. Math. Soc. 356 (2003), 1501–1533.
- [BBCK] M.T. Barlow, R.F. Bass, Z.-Q. Chen and M. Kassmann. Non-local Dirichlet forms and symmetric jump processes. Trans. Amer. Math. Soc. 361 (2009), 1963–1999.
- [BBK1] M.T. Barlow, R.F. Bass and T. Kumagai. Stability of parabolic Harnack inequalities on metric measure spaces. J. Math. Soc. Japan 58 (2006), 485–519.
- [BBK2] M.T. Barlow, R. F. Bass and T. Kumagai. Parabolic Harnack inequality and heat kernel estimates for random walks with long range jumps. *Math. Z.* 261 (2009), 297–320.
- [BGK] M.T. Barlow, A. Grigor'yan and T. Kumagai. On the equivalence of parabolic Harnack inequalities and heat kernel estimates. J. Math. Soc. Japan 64 (2012), 1091–1146.
- [BP] M.T. Barlow and E.A. Perkins. Brownian motion on the Sierpiński gasket. Probab. Theory Relat. Fields 79 (1988), 543–623.
- [BL] R.F. Bass and D. A. Levin. Transition probabilities for symmetric jump processes. Trans. Amer. Math. Soc. 354 (2002), 2933–2953.
- [BKKL] J. Bae, J. Kang, P. Kim and J. Lee. Heat kernel estimates for symmetric jump processes with mixed polynomial growths. To appear in *Ann. Probab.*, available at arXiv:1804.06918.
- [BGT] N.H. Bingham, C.M. Goldie and J.L. Teugels. Regular Variation. Cambridge University Press, Cambridge, 1987.
- [BM] M. Biroli and U. A. Mosco. Saint-Venant type principle for Dirichlet forms on discontinuous media. Ann. Mat. Pura Appl. 169 (1995), 125–181.
- [CKS] E.A. Carlen, S. Kusuoka and D.W. Stroock. Upper bounds for symmetric Markov transition functions. Ann. Inst. Henri Poincaré-Probab. Stat. 23 (1987), 245–287.
- [C] Z.-Q. Chen. On notions of harmonicity. Proc. Amer. Math. Soc. 137 (2009), 3497–3510.
- [CF] Z.-Q. Chen and M. Fukushima. Symmetric Markov Processes, Time Change, and Boundary Theory. Princeton Univ. Press, Princeton and Oxford, 2012.
- [CKK] Z.-Q. Chen, P. Kim and T. Kumagai. Weighted Poincaré inequality and heat kernel estimates for finite range jump processes. *Math. Ann.* 342 (2008), 833–883.
- [CKK2] Z.-Q. Chen, P. Kim and T. Kumagai. On heat kernel estimates and parabolic Harnack inequality for jump processes on metric measure spaces. Acta Math. Sin. (Engl. Ser.) 25 (2009), 1067–1086.
- [CKKW1] Z.-Q. Chen, P. Kim, T. Kumagai and J. Wang. Time fractional poisson equations: representations and estimates. Available at arXiv:1812.04902.
- [CKKW2] Z.-Q. Chen, P. Kim, T. Kumagai and J. Wang. Heat kernel estimates for reflected diffusions with jumps on metric measure spaces. In preparation.
- [CK1] Z.-Q. Chen and T. Kumagai. Heat kernel estimates for stable-like processes on d-sets. Stochastic Process Appl. 108 (2003), 27–62.

- [CK2] Z.-Q. Chen and T. Kumagai. Heat kernel estimates for jump processes of mixed types on metric measure spaces. *Probab. Theory Relat. Fields* 140 (2008), 277–317.
- [CK3] Z.-Q. Chen and T. Kumagai. A priori Hölder estimate, parabolic Harnack principle and heat kernel estimates for diffusions with jumps. *Revista Mat. Iberoamericana* 26 (2010), 551–589.
- [CKW1] Z.-Q. Chen, T. Kumagai and J. Wang. Stability of heat kernel estimates for symmetric non-local Dirichlet forms. To appear in *Memoirs Amer. Math. Soc.*, available at arXiv:1604.04035.
- [CKW2] Z.-Q. Chen, T. Kumagai and J. Wang. Stability of parabolic Harnack inequalities for symmetric non-local Dirichlet forms. To appear in J. European Math. Soc., available at arXiv:1609.07594.
- [CKW3] Z.-Q. Chen, T. Kumagai and J. Wang. Elliptic Harnack inequalities for symmetric non-local Dirichlet forms. J. Math. Pures Appl. 125 (2019), 1–42.
- [ChK] Z.-Q. Chen and K. Kuwae. On subhamonicity for symmetric Markov processes. J. Math. Soc. Japan 64 (2012), 1181–1209.
- [De] T. Delmotte. Parabolic Harnack inequality and estimates of Markov chains on graphs. Revista Mat. Iberoamericana 15 (1999), 181–232.
- [FOT] M. Fukushima, Y. Oshima and M. Takeda. Dirichlet Forms and Symmetric Markov Processes. de Gruyter, Berlin, 2nd rev. and ext. ed., 2011.
- [Gr1] A. Grigor'yan. The heat equation on noncompact Riemannian manifolds. (in Russian) Matem. Sbornik. 182 (1991), 55–87. (English transl.) Math. USSR Sbornik 72 (1992), 47–77.
- [Gr2] A. Grigor'yan. Heat Kernel and Analysis on Manifolds. Amer. Math. Soc., Providence, RI, International Press, Boston, MA, 2009.
- [GH] A. Grigor'yan and J. Hu. Upper bounds of heat kernels on doubling spaces. *Mosco Math. J.* 14 (2014), 505–563.
- [GHH] A. Grigor'yan, E. Hu and J. Hu. Two-sided estimates of heat kernels of jump type Dirichlet forms. Adv. Math. 330 (2018), 433–515.
- [GHL] A. Grigor'yan, J. Hu and K.-S. Lau. Generalized capacity, Harnack inequality and heat kernels on metric spaces. J. Math. Soc. Japan 67 (2015), 1485–1549.
- [GT] A. Grigor'yan and A. Telcs. Two-sided estimates of heat kernels on metric measure spaces. Ann. Probab. 40 (2012), 1212–1284.
- [GSC] P. Gyrya and L. Saloff-Coste. Neumann and Dirichlet heat kernels in inner uniform domains. Astérisque **336**, 2011.
- [HK] B.M. Hambly and T. Kumagai. Transition density estimates for diffusion processes on post critically finite self-similar fractals. Proc. London Math. Soc. 78 (1999), 431–458.
- [MS] M. Murugan and L. Saloff-Coste. Heat kernel estimates for anomalous heavy-tailed random walks. To appear in Ann. Inst. Henri Poincaré-Probab. Stat., available at arXiv:1512.02361.
- [Sa1] L. Saloff-Coste. A note on Poincaré, Sobolev, and Harnack inequalities. Inter. Math. Res. Notices 2 (1992), 27–38.
- [Sa2] L. Saloff-Coste. Aspects of Sobolev-type Inequalities. Cambridge Univ. Press, Cambridge, 2002.
- [SV] R. Song and Z. Vondracek. Parabolic Harnack inequality for the mixture of Brownian motion and stable process. *Tohoku Math. J.* 59 (2007), 1–19.
- [St1] K.-T. Sturm. Analysis on local Dirichlet spaces II. Gaussian upper bounds for the fundamental solutions of parabolic Harnack equations. Osaka J. Math. 32 (1995), 275–312.
- [St2] K.-T. Sturm. Analysis on local Dirichlet spaces III. The parabolic Harnack inequality. J. Math. Pures Appl. 75 (1996), 273–297.

Zhen-Qing Chen

Department of Mathematics, University of Washington, Seattle, WA 98195, USA and School of Mathematics and Statistics, Beijing Institute of Technology, China E-mail: zqchen@uw.edu

Takashi Kumagai:

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan Email: kumagai@kurims.kyoto-u.ac.jp

Jian Wang:

College of Mathematics and Informatics & Fujian Key Laboratory of Mathematical Analysis and Applications (FJKLMAA), Fujian Normal University, 350007, Fuzhou, P.R. China. Email: jianwang@fjnu.edu.cn