arXiv:2001.07599v2 [math.AP] 21 Feb 2020

INVERSE PROBLEMS FOR REAL PRINCIPAL TYPE OPERATORS

LAURI OKSANEN, MIKKO SALO, PLAMEN STEFANOV, AND GUNTHER UHLMANN

ABSTRACT. We consider inverse boundary value problems for general real principal type differential operators. The first results state that the Cauchy data set uniquely determines the scattering relation of the operator and bicharacteristic ray transforms of lower order coefficients. We also give two different boundary determination methods for general operators, and prove global uniqueness results for determining coefficients in nonlinear real principal type equations. The article presents a unified approach for treating inverse boundary problems for transport and wave equations, and highlights the role of propagation of singularities in the solution of related inverse problems.

CONTENTS

1.	Introduction and main results	1
2.	Preliminaries	14
3.	Quasimode construction	20
4.	Determining the scattering relation and ray transforms	34
5.	Boundary determination	41
6.	Semilinear equations	52
Ap	ppendix A. Additional proofs	58
References		62

1. INTRODUCTION AND MAIN RESULTS

1.1. Motivation. This article studies inverse boundary value problems for linear and nonlinear differential operators with variable coefficients. In many works in the theory of inverse boundary value problems, it has been customary to focus on specific problems (e.g. the Calderón problem for Laplace type equations, the inverse boundary problem for the wave equation that we will call *Gel'fand problem* due to the roughly equivalent formulation posed in [Ge54], boundary and scattering rigidity problems for transport equations, or geodesic X-ray transforms). However, there have been unexpected connections between problems having different character, such as

- the explicit appearance of geodesic X-ray transforms in the Calderón problem [DKSU09];
- a reduction from the Calderón problem to the Gel'fand problem involving the boundary control method for the wave equation [DKLS16]; and
- a direct relation between the boundary rigidity problem and the Calderón problem [PU05].

In this article we will work in the spirit of a general point of view to linear partial differential equations, as advocated in [Hö85]. The objective is to find structural conditions that allow us to consider large classes of operators, and to identify fundamental principles that make it possible to solve related inverse problems. In addition to obtaining results for general equations, we hope that the new point of view will yield a better understanding of the methods that are currently available, the connections between different problems, and the extent to which it is possible to push the methods. Of course this is a large program and we view the present work as a starting point.

Here, we will consider operators of real principal type and the consequences of propagation of singularities for related inverse problems. This will include inverse problems for transport equations (boundary and scattering rigidity) and wave equations (Gel'fand problem) as special cases. As an example, if one knows the outgoing Dirichlet-to-Neumann map for the operator P appearing in the wave equation on a Riemannian manifold (M, g) with time-independent magnetic potential A and electric potential V, then up to suitable gauge transformations it is possible to recover

- the Taylor series of the coefficients at the boundary, up to the natural gauge,
- the scattering relation for the metric g,
- the geodesic X-ray transform of the magnetic potential A up to $2\pi\mathbb{Z}$, and
- the geodesic X-ray transform of the electric potential V.

We refer to [SY18], where these results are obtained for Lorentzian wave equations, including classical ones with time-dependent coefficients, and to references therein for earlier results. This reduces the problem of finding the coefficients of P to the well studied boundary/lens rigidity one and to the problem of inverting the geodesic X-ray transform on functions or more generally on tensor fields.

In this work we prove counterparts of the above results for any real principal type differential operator P. The scattering relation for the metric (i.e. for geodesic flow) will be replaced by the scattering relation for the null bicharacteristic flow for P. The geodesic X-ray transform will in turn be replaced by the relevant null bicharacteristic ray transform. In order to recover the actual coefficients it remains to solve the related scattering rigidity and bicharacteristic ray transform problems, if possible. These issues are left to forthcoming works. However, we will also prove boundary determination results for real principal type operators (these can be used to recover real-analytic coefficients), and interior determination results for related nonlinear operators.

1.2. **Real principal type operators.** Before stating the main results, we will need to give some facts concerning real principal type operators. First we recall the definition of these operators from [Hö85, Definition 26.1.8 in Section 26.1].

Definition. Let X be an open C^{∞} manifold. A properly supported pseudodifferential operator $P \in \Psi^m(X)$ is of *real principal type* if it has real homogeneous principal symbol p_m of order m, and if no complete null bicharacteristic of P stays over a compact subset of X.

Examples. Any real vector field with no trapped integral curves is of real principal type. The wave operator $\partial_t^2 - \Delta_{g_0}$ in a space-time cylinder $\mathbb{R} \times M_0$, with (M_0, g_0) a Riemannian manifold, is of real principal type, and so are more general Lorentzian wave operators and higher order hyperbolic operators if the nontrapping condition is satisfied. Tricomi type operators $x_2 \partial_{x_1}^2 + \partial_{x_2}^2$ are real principal type, but Keldysh type operators $x_1 \partial_{x_1}^2 + \partial_{x_2}^2$ are not. Elliptic operators with real principal symbol are real principal type, but they have no null bicharacteristics so for such operators most of our results are void. Our results do apply to operators with elliptic factors such as $(\partial_t^2 - \Delta_{g_0})(\partial_t^2 + \Delta_{g_0})$. Of course, lower order terms with complex coefficients can be included (heuristically, principal type operators are those whose main behaviour is described by the principal symbol alone). We remark that different definitions of real principal type exist in the literature (e.g. the stronger local condition $d_{\xi p} \neq 0$ appears, ensuring that null bicharacteristic curves have no cusps). Moreover, operators such as the Schrödinger operator $i\partial_t - \Delta$ or the plate operator $\partial_t^2 + \Delta^2$ are not real principal type, but using a suitable (anisotropic) weighting for the time derivative makes it possible to treat such operators in a similar way as real principal type operators, although new phenomena appear (see e.g. [Ta04]). One could also consider systems of real principal type, and there are several interesting physical examples (e.g. Maxwell and elastic wave equations).

A major feature of operators with real principal symbol is the fact that singularities propagate along *null bicharacteristics* (i.e. integral curves of the Hamilton vector field H_{p_m} in the characteristic set $\operatorname{Char}(P) := \{(x,\xi) \in T^*X \setminus 0; p_m(x,\xi) = 0\}$): if u solves Pu = 0 in X, then the wave front set of u is contained in $\operatorname{Char}(P)$ and it is invariant under the bicharacteristic flow there. In the definition of real principal type operators, the condition that no complete null bicharacteristic flow. It ensures that the equation Pu = f can be solved in any compact set if f satisfies finitely many linear constraints. These facts from [DH72] and [Hö85, Section 26.1] are recalled in Section 2.

We wish to study inverse boundary value problems for real principal type operators. In order to avoid technicalities related to the presence of a boundary, we will mostly focus on *differential operators* on *compact manifolds with smooth boundary*. We will next examine the real principal type condition and introduce several related definitions in the boundary case.

Let M be a compact manifold with smooth boundary and let P be a differential operator of order $m \geq 1$ having smooth coefficients on M. Assume that P has real principal symbol $\sigma_{\rm pr}[P] = p_m$. Then for any $(x,\xi) \in \operatorname{Char}(P)$, there is a maximal interval $[-\tau_-(x,\xi), \tau_+(x,\xi)]$, where $\tau_{\pm}(x,\xi) \in [0,\infty]$, such that the null bicharacteristic curve

$$\gamma = \gamma_{x,\xi} : \left[-\tau_{-}(x,\xi), \tau_{+}(x,\xi) \right] \to T^*M \setminus 0$$

with $\gamma_{x,\xi}(0) = (x,\xi)$ cannot be extended to a strictly larger closed interval as a smooth bicharacteristic curve in $T^*M \setminus 0$. A bicharacteristic curve is called *maximal* if it is defined in its maximal interval. The null bicharacteristic flow for P is called *nontrapping* if $\tau_{\pm}(x,\xi) < \infty$ for every $(x,\xi) \in \operatorname{Char}(P)$. The operator P is said to be of *real principal type* on a compact manifold Mwith boundary if it has real principal symbol, and if the null bicharacteristic flow is nontrapping. We remark that the above definition is compatible with that of real principal type operators on an open manifold, in the sense that any real principal type differential operator P on a compact manifold M with boundary can be extended smoothly as a real principal type differential operator in some open manifold X containing M (see [Hö85, proof of Theorem 26.1.7]).

We now define the (null bicharacteristic) scattering relation for P. This will be a map that takes the initial point of a maximal null bicharacteristic to its end point, and vice versa. The definition is chosen so that it allows the bicharacteristics to have arbitrary tangential intersections with the boundary.

Definition. Let M be a compact manifold with boundary, and let P be a real principal type differential operator in M. The (null) *incoming* and *outgoing* boundaries are defined as

$$\partial^{\pm}_{\text{null}}(T^*M) := \{(x,\xi) \in \text{Char}(P) \ ; \ \tau_{\mp}(x,\xi) = 0\}$$

We set $\partial'_{\text{null}}(T^*M) := \partial^+_{\text{null}}(T^*M) \cup \partial^-_{\text{null}}(T^*M)$. The scattering relation for P is the map

$$\alpha_P : \partial'_{\text{null}}(T^*M) \to \partial'_{\text{null}}(T^*M),$$
$$\alpha_P(x,\xi) := \begin{cases} \gamma_{x,\xi}(\tau_+(x,\xi)), & (x,\xi) \in \partial^+_{\text{null}}(T^*M), \\ \gamma_{x,\xi}(-\tau_-(x,\xi)), & (x,\xi) \in \partial^-_{\text{null}}(T^*M). \end{cases}$$

It is immediate that $\partial'_{\text{null}}(T^*M) \subset \partial(T^*M) \cap \text{Char}(P)$, and that the scattering relation has the following properties:

$$\alpha_P(\partial_{\text{null}}^{\pm}(T^*M)) = \partial_{\text{null}}^{\mp}(T^*M),$$
$$\alpha_P \circ \alpha_P = \text{Id.}$$

The next step is to consider boundary measurements for P. For second order operators it is customary to consider a Dirichlet-to-Neumann map, which maps the Dirichlet boundary value (on a suitable part of the boundary) of a solution of Pu = 0 in M (the outgoing one for wave equations) to the Neumann boundary value. For a general real principal type operator P of order $m \ge 1$, it is not clear if there is a suitable well-posed boundary value problem or a corresponding boundary map. However, one can always define boundary measurements in terms of the Cauchy data set.

For the next definition and for later purposes, we fix an auxiliary smooth Riemannian metric g on M and let ∇ be the covariant derivative with respect to g. Then $\nabla^k u$ is the kth total covariant derivative of u, and the Cauchy data set below encodes the derivatives of solutions of Pu = 0 in M up to order m - 1 on the boundary.

Definition. The Cauchy data set of P is defined as

$$C_P = \{ (u|_{\partial M}, \nabla u|_{\partial M}, \dots, \nabla^{m-1} u|_{\partial M}) ; u \in H^m(M) \text{ solves } Pu = 0 \text{ in } M \}$$

We say that two differential operators P_1 and P_2 agree to infinite order on ∂M , if $(P_1 - P_2)w$ vanishes to infinite order on ∂M for any $w \in C^{\infty}(M)$.

Note that even though C_P depends on the choice of g, the property $C_P = C_{\tilde{P}}$ for two operators is independent of g.

1.3. Determining the scattering relation and bicharacteristic ray transforms. Let P be a differential operator of order m on M and let μ be a nonvanishing half density on M. We obtain a differential operator P^{μ} acting on half densities by the following definition

(1.1)
$$P^{\mu} = \mu P \mu^{-1},$$

where the operators $\mu : u \mapsto u\mu$ and $\mu^{-1} : u\mu \mapsto u$ give an isomorphism between functions and half densities. Writing $\sum_{j=0}^{m} p_{j}^{\mu}$ for the polyhomogeneous full symbol of P^{μ} in a local coordinate system, the subprincipal symbol

(1.2)
$$\sigma_{\rm sub}[P^{\mu}] = p_{m-1}^{\mu} + \frac{i}{2} \sum_{j=1}^{n} \partial_{x_j \xi_j} p_m^{\mu}$$

is an invariantly defined function on $T^*M \setminus 0$, see e.g. [Hö85, Theorem 18.1.33].

We can now state the first main theorem, showing that the boundary measurements for any real principal type differential operator P uniquely determine the scattering relation which depends on the principal symbol p_m . Moreover, if p_m is known then the integrals of the subprincipal symbol $\sigma_{sub}[P^{\mu}]$ over maximal null bicharacteristics are uniquely determined modulo $2\pi\mathbb{Z}$.

The result assumes that the coefficients of P have already been determined up to infinite order on ∂M (after suitable gauge transformations); this is typically done using boundary determination results, which are discussed later.

Theorem 1.1. Let M be a compact manifold with smooth boundary, and let P_1, P_2 be real principal type differential operators of order $m \ge 1$ on M. If

$$C_{P_1} = C_{P_2}$$

and if $P_1 = P_2$ to infinite order on ∂M , then

$$\partial'_{\text{null},P_1}(T^*M) = \partial'_{\text{null},P_2}(T^*M),$$
$$\alpha_{P_1} = \alpha_{P_2}.$$

Moreover, if the principal symbols of P_1 and P_2 coincide, then for any nonvanishing half density μ on M one has

(1.3)
$$\exp\left[i\int_0^T \sigma_{\rm sub}[P_1^{\mu}](\gamma(t))\,dt\right] = \exp\left[i\int_0^T \sigma_{\rm sub}[P_2^{\mu}](\gamma(t))\,dt\right]$$

whenever $\gamma: [0,T] \to T^*M$ is a maximal null bicharacteristic curve for P_1 .

Remark. The equation (1.3) can be formulated equivalently without half densities as

$$\exp\left[i\int_0^T (p_{m-1,1} - p_{m-1,2})(\gamma(t))\,dt\right] = 0,$$

where $\sum_{j=0}^{m} p_{j,1}$ and $\sum_{j=0}^{m} p_{j,2}$ are polyhomogeneous full symbols of P_1 and P_2 , acting on functions, in a local coordinate system. The above expression is coordinate invariant by Lemma 3.2 under the assumptions above, and the equivalence follows from (3.2) below.

The conclusion (1.3) is of course equivalent to

$$\int_0^T \sigma_{\rm sub}[P_1^{\mu}](\gamma(t)) \, dt = \int_0^T \sigma_{\rm sub}[P_2^{\mu}](\gamma(t)) \, dt \quad \text{modulo } 2\pi\mathbb{Z}.$$

This nonuniqueness modulo $2\pi\mathbb{Z}$ is related to the *Aharonov-Bohm effect*, which appears when one tries to determine subprincipal terms on domains with nontrivial topology (see Lemma 2.6).

The next result shows that it is also possible to determine bicharacteristic ray transforms of the coefficients of order $\leq m-2$, and that no nonuniqueness modulo $2\pi\mathbb{Z}$ appears in this case.

Theorem 1.2. Let M be a compact manifold with smooth boundary, let P be a real principal type differential operator of order $m \ge 2$ on M, and let Q_1 and Q_2 be differential operators of order $\le m - 2$ on M with $Q_1 = Q_2$ to infinite order on ∂M . If

$$C_{P+Q_1} = C_{P+Q_2},$$

then

(1.4)
$$\int_{0}^{T} \sigma_{\rm pr}[Q_1](\gamma(t)) \, dt = \int_{0}^{T} \sigma_{\rm pr}[Q_2](\gamma(t)) \, dt$$

whenever $\gamma: [0,T] \to T^*M$ is a maximal null bicharacteristic curve for P.

Remark. Note that $\sigma_{pr}[Q_j](x,\xi) = q_j^{j_1\dots j_k}(x)\xi_{j_1}\dots\xi_{j_k}$ is a polynomial in ξ in local coordinates. Hence (1.4) may be written as

$$\int_0^T q_1^{j_1\dots j_k}(x(t))\xi_{j_1}(t)\dots\xi_{j_k}(t)\,dt = \int_0^T q_2^{j_1\dots j_k}(x(t))\xi_{j_1}(t)\dots\xi_{j_k}(t)\,dt$$

where $\gamma(t) = (x(t), \xi(t))$.

Note that the above theorems are rather general, in the sense that they are valid for

- differential operators of any order, with real principal symbol and satisfying the nontrapping condition (in particular, no wellposedness assumptions are required); and
- any maximal null bicharacteristic, possibly with cusp points (such as for Tricomi operators) or tangential intersections with the boundary.

The proofs of the above theorems extend in a straightforward way to the case where one only has access to measurements on a subset $\Gamma \subset \partial M$. Then one obtains information along null bicharacteristics that do not meet ∂M away from Γ . We refer to Section 4 for precise statements.

Another related case is that of hyperbolic operators, for instance the wave operator in a spacetime cylinder $M_0 \times (0,T)$, where the boundary measurements are typically given on the lateral boundary $\partial M_0 \times (0,T)$ for outgoing solutions that vanish near t = 0. This case can also be included with small modifications, and it is a special case of the following result.

Consider a differential operator P of order m on a smooth open manifold X equipped with a smooth function ϕ . We say that P is *strictly hyperbolic* with respect to the level surfaces of ϕ (see [Hö85, Definition 23.2.3]) if, writing $p_m = \sigma_{\rm pr}[P]$, one has $p_m(x, d\phi(x)) \neq 0$ for $x \in X$ and if the polynomial $\tau \mapsto p_m(x, \xi + \tau d\phi(x)) = 0$ has m distinct real roots for all $(x, \xi) \in T^*X$ such that ξ is linearly independent from $d\phi(x)$. Without loss of generality we may assume that p_m is real valued (see [Hö85, Lemma 8.7.3]).

Let $M_1 \subset X$ be a compact domain with smooth boundary and let T > 0. We consider the cylinder

(1.5)
$$M = \{ x \in M_1 ; 0 \le \phi(x) \le T \},\$$

and write $\Gamma = \partial M \cap \partial M_1$, $\Gamma_- = \partial M \cap \{x \in M_1; \phi(x) = 0\}$ and $\Gamma_+ = \partial M \cap \{x \in M_1; \phi(x) = T\}$. If $\Sigma \subset \partial M$, we use the shorthand notation

$$\operatorname{tr}_{\Sigma}^{m-1} u = (u|_{\Sigma}, \nabla u|_{\Sigma}, \dots, \nabla^{m-1} u|_{\Sigma}).$$

Finally we define the lateral Cauchy data set

$$C_P^{\text{lat}} = \{ \operatorname{tr}_{\Gamma}^{m-1} u \, ; \, u \in H^m(M) \text{ solves } Pu = 0 \text{ in } M \text{ and } \operatorname{tr}_{\Gamma_-}^{m-1} u = 0 \}.$$

We have the following analogue of Theorems 1.1 and 1.2.

Theorem 1.3. Let M be as in (1.5), and let P_1, P_2 be differential operators of order $m \ge 1$, strictly hyperbolic with respect to the level surfaces of ϕ . Suppose that $P_1 = P_2$ to infinite order on Γ .

If $C_{P_1}^{\text{lat}} = C_{P_2}^{\text{lat}}$, then

$$\alpha_{P_1}(x,\xi) = \alpha_{P_2}(x,\xi)$$

for any $(x,\xi) \in \partial'_{\operatorname{null},P_1}(T^*M)$ such that both for j = 1,2, the maximal null P_j -bicharacteristic curve through (x,ξ) does not meet $\partial(T^*M)$ away from Γ .

Moreover, if the principal symbols of P_1 and P_2 coincide, then for any nonvanishing half density μ on M one has

(1.6)
$$\exp\left[i\int_0^T \sigma_{\rm sub}[P_1^{\mu}](\gamma(t))\,dt\right] = \exp\left[i\int_0^T \sigma_{\rm sub}[P_2^{\mu}](\gamma(t))\,dt\right]$$

whenever $\gamma : [0,T] \to T^*M$ is a maximal null bicharacteristic curve for P_1 whose spatial projection does not meet ∂M away from Γ .

Finally, if
$$m \ge 2$$
 and if $P_j = P + Q_j$ where Q_j has order $\le m - 2$ for $j = 1, 2$, then

$$\int_0^T \sigma_{\rm pr}[Q_1](\gamma(t)) \, dt = \int_0^T \sigma_{\rm pr}[Q_2](\gamma(t)) \, dt$$

whenever $\gamma : [0,T] \to T^*M$ is a maximal null bicharacteristic curve for P whose spatial projection does not meet ∂M away from Γ . 1.4. Boundary determination results. We proceed to the statement of boundary determination results. Given $(x,\xi) \in T^*(\partial M)$, the boundary determination results at x will depend on properties of the characteristic polynomial in the normal direction, i.e. on the polynomial

$$\tau \mapsto p_m(x,\xi + \tau\nu),$$

where ν is the *inward pointing unit conormal* to ∂M with respect to the auxiliary metric g. This is a polynomial of order m with real coefficients, and hence has m complex roots counted with multiplicity so that the non-real roots occur in complex conjugate pairs. We will employ two methods for determining boundary values at x:

- (Elliptic region) If $\tau \mapsto p_m(x, \xi + \tau \nu)$ has a simple non-real root, we use exponentially decaying solutions that concentrate near x to give an analogue of boundary determination results for second order elliptic equations.
- (Hyperbolic region) If $\tau \mapsto p_m(x, \xi + \tau \nu)$ has at least two distinct real roots, we use solutions concentrating near two null bicharacteristics through x and obtain an analogue of boundary determination results for the wave equation.

In fact the regions could be mixed, and we will use a combination of both methods. The terminology above is analogous to the case of second order operators [Hö85, Section 24.2]: if P is of second order with real principal symbol p_2 and the boundary is noncharacteristic, then the elliptic region (resp. hyperbolic region) is the set of those points in $(x,\xi) \in T^*(\partial M)$ such that the map $\tau \mapsto p_2(x,\xi+\tau\nu)$ has no real zeros (resp. has two distinct real zeros).

The next result proves that the Cauchy data set determines the Taylor series of the principal symbol of P at x, modulo a natural gauge invariance given by $C_P = C_{cP}$ where $c \in C^{\infty}(M, \mathbb{R})$ is nonvanishing. The result assumes that there is some $\xi \in T_x^*(\partial M)$ such that $\tau \mapsto p_m(x, \xi + \tau \nu)$ has only simple roots, with an additional geometric condition for the real roots.

Definition. We say that two smooth curves $\gamma_j = (x_j, \xi_j) : [0, T_j] \to T^*M$ intersect nicely at $x \in \partial M$ if $x_1(0) = x_2(0) = x$ with $\dot{x}_1(0)$ and $\dot{x}_2(0)$ linearly independent, if the curves $x_j(t)$ never intersect at the boundary again, and if $\xi_1(t) \neq \xi_2(s)$ when $x_1(t) = x_2(s)$.

Example. Let $P = \partial_t^2 - \Delta_{g_0}$ be the wave operator in the space-time cylinder $M = M_0 \times (0, T)$ where (M_0, g_0) is a compact Riemannian manifold with smooth boundary, and let $(x_0, t_0) \in \partial M_0 \times (0, T)$. Consider two null bicharacteristics γ_1 and γ_2 through (x_0, t_0) so that γ_1 goes forward in time and γ_2 backward in time. Then γ_1 and γ_2 intersect nicely at (x_0, t_0) . A similar picture holds for strictly hyperbolic operators.

Theorem 1.4. Let M be a compact manifold with smooth boundary and let P_1, P_2 be real principal type differential operators of order $m \ge 2$. Assume that

$$C_{P_1} = C_{P_2}.$$

Suppose that $x_0 \in \partial M$ is noncharacteristic for P_j , and for some $\xi_0 \in T^*_{x_0}(\partial M)$ the maps $\tau \mapsto p_{m,j}(x_0,\xi_0+\tau\nu)$ have only simple roots. Moreover, assume one of the following conditions:

- (1) some root is non-real, and no bicharacteristic corresponding to a real root returns to x_0 ; or
- (2) all roots are real, and whenever $p_{m,1}(x_0, \xi_0 + \tau_1 \nu) = p_{m,2}(x_0, \xi_0 + \tau_2 \nu) = 0$ for $\tau_1 \neq \tau_2$ the bicharacteristics through $(x_0, \xi_0 + \tau_1 \nu)$ and $(x_0, \xi_0 + \tau_2 \nu)$ intersect nicely at x_0 .

Then there is a nonvanishing function $c \in C^{\infty}(M)$ so that in boundary normal coordinates (x', x_n) (with respect to the background Riemannian metric) one has

(1.7)
$$\partial_{x_n}^j (p_{m,2}(x,\eta) - c(x)p_{m,1}(x,\eta))|_{\substack{x=x_0 \\ 7}} = 0, \qquad j \ge 0, \ \eta \in T^*_{x_0}M.$$

The next result considers boundary determination of Taylor series of lower order terms. It turns out that to determine coefficients of order r, one needs that $\tau \mapsto p_m(x_0, \xi_0 + \tau \nu)$ has at least r + 1simple roots, with an additional geometric condition for the real roots.

Theorem 1.5. Let M be compact with smooth boundary, let P be a real principal type differential operator of order $m \ge 2$, and let Q_1, Q_2 be differential operators of order $r \le m - 1$. Assume that

$$C_{P+Q_1} = C_{P+Q_2}.$$

Suppose that for some $(x_0, \xi_0) \in T^*(\partial M)$ the map $\tau \mapsto p_m(x_0, \xi_0 + \tau \nu)$ has at least s simple roots τ_1, \ldots, τ_s with nonnegative imaginary parts, and assume one of the following two conditions:

- (1) $s \ge r+1$, one of the τ_j is non-real, and no bicharacteristic corresponding to a real τ_j returns to x_0 ; or
- (2) $s \ge \max(r+1,2)$, each τ_j is real, and for $j \ne k$ the bicharacteristics through $(x_0,\xi_0+\tau_j\nu)$ and $(x_0,\xi_0+\tau_k\nu)$ intersect nicely at x_0 .

Then in boundary normal coordinates

$$\partial_{x_n}^j \sigma_{\mathrm{pr}}[Q_1](x_0,\eta) = \partial_{x_n}^j \sigma_{\mathrm{pr}}[Q_2](x_0,\eta), \qquad j \ge 0, \ \eta \in T_{x_0}^* M.$$

Moreover, if $Q_1 - Q_2$ has real principal symbol, the result is valid without the assumption that the roots τ_i have nonnegative imaginary parts.

The results above give an immediate consequence for determining zero order terms.

Theorem 1.6. Let M be a compact manifold with smooth boundary, and let P be a real principal type differential operator of order $m \ge 2$. Let $V_1, V_2 \in C^{\infty}(M)$, and assume that

$$C_{P+V_1} = C_{P+V_2}.$$

Then the Taylor series of V_1 and V_2 agree at any point $x \in \partial M$ so that for some $\xi \in T^*_x(\partial M)$, the map $\tau \mapsto p_m(x, \xi + \tau \nu)$ either has a simple non-real root, or two distinct real roots τ_1 and τ_2 so that the corresponding bicharacteristics intersect nicely at x.

In particular, if M, V_1 and V_2 are real-analytic and one such point x exists, then $V_1 = V_2$ in M.

The approach for proving the above results yields the following, perhaps surprising observations:

- It is possible to perform boundary determination for general real principal type operators, including wave operators, just as for the Laplace operator by working in the elliptic region. The method also works for elliptic operators of any order if sufficiently many roots are simple.
- Boundary determination in the elliptic region is local in character (one does not need any global assumption on behaviour of bicharacteristics).
- Boundary determination in the hyperbolic region may be local or global in character. If two null bicharacteristics only meet at their initial point (e.g. they go in opposite time directions), one recovers the Taylor series at that point. If they meet at two distinct boundary points, then one recovers a sum of contributions from these two points.
- Boundary determination for the wave operator is possible also at the bottom of a spacetime cylinder, and this argument is global in character. Measuring local Cauchy data at the bottom is not enough to determine the Taylor series of coefficients, since the Cauchy problem is well-posed (i.e. Cauchy data at the bottom does not carry any information about the coefficients). However, it is possible to use two null bicharacteristics intersecting at a point in the bottom, and information measured at the lateral boundary can be used to determine the Taylor series of coefficients at the bottom.

1.5. Nonlinear equations. We will next state our results for nonlinear equations whose principal part is a linear real principal type operator. In this case, instead of obtaining conditional results that reduce the problem of recovering coefficients to inverting a scattering relation or ray transform, we can actually determine the unknown coefficients at all points satisfying a geometric condition. Here we adapt the methods initiated in the case of Einstein equations and nonlinear wave equations [KLU18, KLOU14, LLLS19, FO19], which are based on higher order linearizations and use the fact that the presence of nonlinear interactions may make the inverse problem easier to solve.

The model equation that we consider here is

$$Pu + a(x, u) = 0$$
 in M

where P is a linear real principal type differential operator of order m and a(x, u) is a nonlinearity satisfying the following conditions: for a fixed integer $s > \max(m, n/2)$,

(1.8) a(x,z) is analytic in z near 0 as a $H^{s}(M)$ -valued function,

(1.9)
$$a(x,0) = \partial_z a(x,0) = \partial_z^2 a(x,0) = 0.$$

The condition $\partial_z^2 a(x,0) = 0$ simplifies the proof, and possibly could be removed with extra work. The condition $\partial_z a(x,0) = 0$ however is important, since it ensures that the equation is not linear (if one removes this assumption then one could only determine the genuinely nonlinear parts, i.e. $\partial_z^j a(x,0)$ for $j \ge 2$).

We consider boundary measurements given in terms of the Cauchy data set with $\delta > 0$ small,

$$C_{a,\delta} = \{ (u|_{\partial M}, \nabla u|_{\partial M}, \dots, \nabla^{m-1} u|_{\partial M}) ; u \in H^{s}(M), Pu + a(x,u) = 0, \text{ and } \|u\|_{H^{s}(M)} \le \delta \}.$$

Here it is instrumental to work in the regime of small solutions, since the argument relies on higher order linearizations around the zero solution. We will also need to make a weak uniqueness assumption for the linearized equation and its adjoint. The space

$$N(P) = \{ u \in H_0^m(M) ; Pu = 0 \text{ in } M \}$$

is always finite dimensional (see Proposition 2.2). We will assume that this space is trivial, which roughly means that the linearized equation does not admit nontrivial compactly supported solutions.

Theorem 1.7. Let M be a compact manifold with smooth boundary, and let P be a real principal type differential operator on M. Let a, \tilde{a} satisfy (1.8)–(1.9), and assume that a and \tilde{a} agree to high order on ∂M in the sense that

$$a(\cdot, z) - \tilde{a}(\cdot, z) \in H^s_0(M)$$
 for z near 0.

Assume also that

$$N(P) = N(P^*) = \{0\}.$$

If for any sufficiently small $\delta > 0$ there is $\delta_1 < \delta$ so that

$$C_{a,\delta_1} \subset C_{\tilde{a},\delta},$$

then

$$a(x_0, z) = \tilde{a}(x_0, z)$$
 for any $x_0 \in B$ and any z near $0, z$

where

$$B = \{x_0 \in M^{\text{int}}; \text{ there are two maximal null bicharacteristics whose spatial} \\ \text{projections } x_j, \ j = 1, 2, \ \text{only intersect once at } x_0 \ \text{and} \\ \text{the tangent vectors of } x_j \ \text{are linearly independent at } x_0 \}.$$

Example. Let M be a compact subdomain of the space-time cylinder $M_0 \times (0, T)$ where (M_0, g_0) is a compact Riemannian manifold with boundary, and let $P = \partial_t^2 - \Delta_{g_0}$ be the wave operator. If $(x_0, t_0) \in M$ and if there is a geodesic $\eta(s)$ in M_0 so that $\eta(0) = x_0$ and some Jacobi field along η only vanishes when s = 0, then by looking at the corresponding variation through geodesics one can show that (x_0, t_0) is in the set B of Theorem 1.7. In particular, $(x_0, t_0) \in B$ if there is at least one geodesic through x_0 with no conjugate points, but one may have $(x_0, t_0) \in B$ even if all geodesics through x_0 have conjugate points as long as some variation field only vanishes when s = 0. A similar argument can be given for general real principal type operators by looking at variation fields of null bicharacteristic curves. However, this argument fails in the special situation when $x_0 \in M_0$ and every M_0 -geodesic through x_0 has a conjugate point y of maximal order (i.e. the space of normal Jacobi fields vanishing both at x_0 and y has maximal order dim $(M_0) - 1$). The sphere is the standard example of a manifold where any point is maximally conjugate to its antipodal point. If M_0 contains a neighborhood of the hemisphere, then the set B in Theorem 1.7 would be empty.

1.6. Special cases. We next give a few basic examples illustrating the above results, related to

- the boundary / scattering rigidity problem; and
- the inverse boundary problem for the Lorentzian wave equation.

Example 1.8. (Boundary / scattering rigidity) Let (M_0, g_0) be a compact Riemannian manifold with strictly convex boundary, and assume that the geodesic flow in (M_0, g_0) is nontrapping. Let $M := SM_0 = \{(x, v) \in TM_0; |v|_{g_0} = 1\}$ be the unit sphere bundle of M_0 , and let

$$P := X + \mathcal{A}$$

where X is the geodesic vector field in SM_0 and $\mathcal{A} \in C^{\infty}(SM_0)$. Since P is a first order operator, projections of null bicharacteristic curves to M are integral curves of P (i.e. geodesics in SM_0) and the nontrapping assumption ensures that P is of real principal type. Moreover, the scattering relation α_P induces a corresponding relation $\beta : \partial(SM_0) \to \partial(SM_0)$ so that $\beta \circ \pi_M = \pi_M \circ \alpha_P$ where π_M is the projection $T^*M \to M$. Thus β is just the scattering relation for the metric g_0 in M_0 . The Cauchy data set is

$$C_P = \{ u |_{\partial(SM_0)}; (X + \mathcal{A})u = 0 \text{ in } SM_0 \}.$$

Theorem 1.1 states that C_P determines α_P , and hence β , as well as the quantities

$$\int_{\eta} \mathcal{A}(\eta(t)) \, dt \text{ modulo } 2\pi\mathbb{Z}$$

whenever η is a maximal geodesic in SM_0 . Conversely, the above information uniquely determines C_P (see Theorem 4.4). These results are quite elementary since P is of first order, but it is instructive to observe that the inverse boundary problem for $X + \mathcal{A}$ is equivalent to the scattering rigidity problem of determining a metric g_0 up to gauge from its scattering relation β , and to determining \mathcal{A} from its geodesic X-ray transform. It is well known that a general function $\mathcal{A} \in C^{\infty}(SM_0)$ is not determined uniquely by its X-ray transform. In fact, the kernel of the X-ray transform on $C^{\infty}(SM_0)$ consists precisely of the functions $\mathcal{A} = X\mathcal{B}$ where $\mathcal{B} \in C^{\infty}(SM_0)$ with $\mathcal{B}|_{\partial SM_0} = 0$ (the proof is an elementary argument combined with [PSU12, Proposition 5.2]). However, in applications \mathcal{A} often arises from a function or a tensor field on the base manifold M_0 and there are many available results in this case.

Example 1.9. (Lorentzian wave equation) Let (M, g) be a compact Lorentzian manifold with boundary, and consider the canonical wave operator \Box_g on M. In local coordinates,

$$\Box_g u = -|g|^{-1/2} \partial_{x_j} (|g|^{1/2} g^{jk} \partial_{x_k} u),$$

where $|g| = \det(g_{jk})$ and (g^{jk}) is the inverse of (g_{jk}) . The principal symbol p_2 of \Box_q is

(1.10)
$$p_2(x,\xi) = g^{jk}\xi_j\xi_k = \langle \xi, \xi \rangle_g.$$

In this case, results similar to Theorems 1.1-1.6 are contained in [SY18], with stability estimates.

The operator \Box_g is not necessarily of real principal type if no further assumptions on the geometry is made. Writing S^n for the unit sphere of dimension n, a counter-example is given by $M = S^1 \times M_0$, where M_0 is a compact domain with boundary on S^n , containing a closed geodesic, and g(t,x) = $-dt^2 + g_0(x)$ for $(t,x) \in S^1 \times M_0$ with dt and g_0 the natural metrics on S^1 and S^n , respectively. Indeed, in this case there is a closed null geodesic on M, and \Box_g is not of real principal type.

If all null geodesics exit M, then \Box_g is of real principal type. This is the case, for example, if (X,g) is a globally hyperbolic Lorentzian manifold and $M \subset X$ is a compact domain with smooth boundary. We refer to the books [O'N83, Ri09] for the definition and properties of globally hyperbolic Lorentzian manifolds. The classical case where $M = M_0 \times (0,T)$ for some T > 0 and

(1.11)
$$g(x,t) = -dt^2 + g_0(x),$$

with (M_0, g_0) a Riemannian manifold with boundary, is covered in Theorem 1.3.

Let a be a one form on M, and consider the wave operator $P = (d + ia)^*(d + ia)$. Here the adjoint is induced by the Lorentzian inner product g and d is the exterior derivative. Then the principal symbol of P is given by p_2 in (1.10). Defining a half density in a local coordinate system by $\mu = |g|^{1/4} |dx_1 \wedge \ldots \wedge dx_n|^{1/2}$, consistent with the volume form, yields that

$$\sigma_{\rm sub}[P^{\mu}](x,\xi) = g^{jk}a_j\xi_k = \langle a,\xi\rangle_g.$$

The expression $\exp\left[i\int_0^T \sigma_{\rm sub}[P^{\mu}](\gamma(t))\,dt\right]$, cf. (1.3), gives the parallel transport map with respect to the connection d+ia along the bicharacteristic $\gamma:[0,T] \to T^*M$ of P as follows: if $z(t) = \pi(\gamma(t))$ where $\pi: T^*M \to M$ is the natural projection, and if S(t) is a parallel section of the trivial bundle $M \times \mathbb{C}$ along z(t) (i.e. $\dot{S}(t) + ia(\dot{z}(t))S(t) = 0$), then $S(T) = \exp\left[-i\int_0^T \sigma_{\rm sub}[P^{\mu}](\gamma(t))\,dt\right]S(0)$.

Let $V \in C^{\infty}(M)$ have principal symbol $v_0(x,\xi) = V(x)$, and suppose that P has principal symbol p_2 in (1.10). Then the map in (1.4) corresponds to

$$LV(\gamma) = \int_0^T v_0(\gamma(t))dt = \int_0^T V(\pi(\gamma(t)))dt,$$

where $\gamma : [0,T] \to T^*M$ is a maximal bicharacteristic of P. This is the light ray transform of V. It is known that L is invertible in the case of certain stationary, globally hyperbolic manifolds [FI019], and in the case of certain real analytic manifolds [St17]. We refer also to [FIKO19] for the case of product geometries of the form (1.11).

We remark that the gauge invariances in Theorems 1.1-1.6, when specialized to the Lorentzian wave equation, are slightly different from those in [SY18]. In the boundary determination result in Theorem 1.5 there is no gauge invariance $a \to a - d\psi$ where $\psi \in C^{\infty}(M)$ satisfies $\psi|_{\partial M} = 0$, whereas such a gauge invariance appears in [SY18, Lemma 2]. This is due to the fact that our boundary measurements (Cauchy data set) for P are defined in terms of a reference Riemannian metric on M, whereas [SY18] defines boundary measurements in terms of the Lorentzian metric g and 1-form a appearing in P. The gauge invariances are discussed in more detail in Section 2.

1.7. Methods. The proofs of our theorems are essentially based on propagation of singularities for real principal type operators. If P is such an operator, propagation of singularities is often understood as a regularity result stating that if a solution u of Pu = 0 is smooth at some point on a null bicharacteristic curve, then u has to be smooth at all points on this curve. This result has a complement stating that for any suitable non-trapped null bicharacteristic segment γ for P, there is a function u solving Pu = 0 away from the end points of γ so that the wave front set of u is precisely on γ , see [DH72] and [Hö85, Theorem 26.1.7]. This complementary result, which is basically a quasimode (or approximate solution) construction, produces interesting special solutions to the equation Pu = 0. It is this result and its variants that will be useful in the solution of inverse problems.

It is natural to define the Cauchy data set for sufficiently smooth solutions of Pu = 0. Thus the singular solutions mentioned above are not directly useful, but rather we will use a semiclassical version of such a quasimode construction. There are various methods for constructing quasimodes. If the null bicharacteristic segment has no cusps, a classical geometrical optics or Gaussian beam type construction is sufficient (see e.g. [BL67, Hö71, Ra82, Ra01]). The general case with cusps is more involved, and in [DH72] it is dealt with by conjugating P microlocally in a full neighborhood of the null bicharacteristic segment to the normal form D_{x_1} using Fourier integral operators. In principle the quasimode construction in this article could be done by implementing a semiclassical version of the Fourier integral operator conjugation argument (see [DKLS16] for a special case). However, in the proofs of the main theorems we need to perform various computations with these quasimodes on manifolds with boundary. To handle the boundary effects it is beneficial to have a more direct construction. We will use a modification of the Gaussian beam construction based on representing the quasimode as an integral of Gaussian wave packets (coherent states) over the bicharacteristic as in [PU93, KV98]. This approach resolves the problems near cusps and gives an elementary and direct construction. In [GUW16], which describes a general framework, such quasimodes are called semiclassical states associated with isotropic submanifolds of phase space.

Given a quasimode v concentrating near a null bicharacteristic, we employ the solvability theory of real principal type operators to produce exact solutions of Pu = 0 that are close to v. We use these special solutions to pass from the Cauchy data set C_P to the scattering relation or null bicharacteristic ray transform by using two different methods: a mix-and-match construction combined with semiclassical propagation of singularities, or an integral identity (Lemma 2.8) analogous to standard identities used in the Calderón and Gel'fand problems. For the boundary determination results we again use integral identities and solutions concentrating near null bicharacteristics in the hyperbolic region. However, in the elliptic region we need different special solutions that concentrate near a boundary point and are exponentially decreasing in the interior (Theorem 5.1). These solutions are analogous to those used in boundary determination for elliptic second order equations. Similar arguments appear in the construction of a boundary parametrix for the wave equation in the elliptic region (see e.g. [Ta81, Chapter IX]). The coefficients are eventually recovered via the complex stationary phase method, see [Hö85, Section 7.7] or [MS74] for the original approach.

The proof of Theorem 1.7 for nonlinear equations is based on higher order linearizations. The idea is that given solutions v_1, \ldots, v_N of the linearized equation (i.e. $Pv_j = 0$) and small parameters $\varepsilon_1, \ldots, \varepsilon_N$, there is a small solution u of the nonlinear equation Pu + a(x, u) = 0 so that u is close to $\varepsilon_1v_1 + \ldots + \varepsilon_Nv_N$. Taking suitable derivatives with respect to ε_j and using the knowledge of the Cauchy data of small solutions, one recovers information about Taylor coefficients of a(x, u) integrated against products of the solutions v_j . The key point is to use special solutions v_j that concentrate along different null bicharacteristic curves, which leads to the geometric condition in Theorem 1.7.

1.8. **Previous literature and outlook.** There is a large literature on inverse boundary problems of the above type, and we only give a few relevant references. The prototype for Theorems 1.1–1.3 are corresponding results for the wave equation, going back to [RS88, Ra90, St89] in the Euclidean case. The scattering relation was considered in [Gu76], and [Uh04] outlines how to determine the scattering relation for a Riemannian metric from the hyperbolic Dirichlet-to-Neumann map. The case of Lorentzian wave equations is discussed in [SY18], together with references to earlier work. Boundary determination results for wave equations may be found in [SU90, SU05, Mo14, SY18]. Boundary determination for second order elliptic equations goes back to [KV84, SU88], with [LU89, DKSU09] considering the geometric case.

Inverse problems for nonlinear hyperbolic equations have been studied in [KLU18, KLOU14, LUW18], and we have also used ideas from related results for elliptic equations [LLLS19, FO19]. We also mention the work [Is91] studying inverse boundary problems for certain general equations of the form P(D)u + Vu = 0, where P(D) is a constant coefficient operator. The present article addresses similar questions for operators with variable coefficients.

As mentioned above, we consider this article to be a starting point in approaching inverse problems from a more general point of view. This point of view, in addition to the particular theorems stated here, is one of the main contributions of this article. It suggests many future directions including the following:

- 1. What kind of information about P (more precisely, about the principal symbol $\sigma_{\rm pr}[P]$) can be determined from the null bicharacteristic scattering relation α_P ? If P is the geodesic vector field on the unit sphere bundle of a compact Riemannian manifold with boundary, this is the scattering/lens/boundary rigidity problem studied in [PU05, SUV16, SUV17].
- 2. For which operators P and for which classes of functions on T^*M is the null bicharacteristic ray transform invertible? If P is the geodesic vector field, this transform is the geodesic ray transform for which there is a substantial literature, see the survey paper [IM19]. If P is the wave operator on a Lorentzian manifold, this transform is the light ray transform studied in [FIKO19, FIO19, LOSU19, St17, VW19].
- 3. The real principal type condition includes a nontrapping assumption for the null bicharacteristic flow. Can one obtain results in the presence of (sufficiently mild) trapping? For the geodesic vector field and hyperbolic trapping, this has been studied in detail in [Gu17].
- 4. We have only considered uniqueness results stating that C_P uniquely determines some information about P. Is it possible to study stability, reconstruction, range characterization, or further partial data cases? Moreover, instead of looking at scalar operators, can one give similar results for real principal type systems?
- 5. In this article we have studied the consequences of propagation of singularities for inverse problems for real principal type operators. What other classes of operators or general mechanisms for inverse problems could be studied in this way?
- 6. In examples such as the Calderón or Gel'fand problem, knowing the Cauchy data set is equivalent to knowing a boundary map (Dirichlet-to-Neumann map) which is a pseudodifferential or a Fourier integral operator. In these cases one can determine information about the coefficients by symbol computations. Our proofs are similar, but we use integral identities as a substitute for symbol computations. If P is a general operator, is it possible to associate a symbol or Lagrangian manifold directly to the Cauchy data set C_P ?
- 7. The symbol computations mentioned in the previous item only use the singularities of the integral kernel of the boundary map. The arguments in this article are in a similar spirit. Is it possible to extract information from the C^{∞} part of the integral kernel? This is what happens in the Calderón problem, see e.g. [Sa17] for a related discussion. Also the Boundary

Control method, originating from [Be87], uses the smooth part in the case of wave equations with time-independent coefficients.

The rest of the article is organized as follows. In Section 2 we collect some preliminaries related to real principal type operators, semiclassical wave front sets, and Cauchy data sets. Section 3 gives the construction of semiclassical quasimodes associated with null bicharacteristic segments. Section 4 proves Theorems 1.1–1.3 related to determining the scattering relation and bicharacteristic ray transforms from the Cauchy data set. The boundary determination results, Theorems 1.4–1.6, are established in Section 5. Finally, Section 6 considers semilinear equations and proves Theorem 1.7. The proofs of some auxiliary results are given in Appendix A.

Acknowledgements. L.O. was supported by EPSRC grants EP/P01593X/1 and EP/R002207/1. M.S. was supported by the Academy of Finland (Finnish Centre of Excellence in Inverse Modelling and Imaging, grant numbers 312121 and 309963) and by the European Research Council under Horizon 2020 (ERC CoG 770924). P.S. was partly supported by NSF Grant DMS-1900475. G.U. was partly supported by NSF, a Si-Yuan Professorship at HKUST and a Walker Family Professorship at UW. This material is based upon work supported by the Clay Foundation and NSF under Grant No. 1440140, while M.S. and G.U. were in residence at MSRI in Berkeley, California, during the semester on Microlocal Analysis in 2019.

2. Preliminaries

Real principal type operators. We begin by recalling basic facts about real principal type operators from [Hö85, Section 26.1].

First assume that $P \in \Psi^m(X)$ is properly supported and has real homogeneous principal symbol $p = \sigma_{\rm pr}[P]$ of order m. The bicharacteristic curves of P are integral curves of the Hamilton vector field H_p on $T^*X \setminus 0$. In local coordinates, a bicharacteristic curve $\gamma(t) = (x(t), \xi(t))$ solves the Hamilton equations

$$\dot{x}(t) = d_{\xi} p(x(t), \xi(t)),$$

$$\dot{\xi}(t) = -d_x p(x(t), \xi(t))$$

The function p is constant along any bicharacteristic curve. In particular, the bicharacteristic curves in the characteristic set $p^{-1}(0)$ are called null bicharacteristic curves. A major feature of operators with real principal symbol is the fact that singularities propagate along null bicharacteristics: if Pu = f in X, then WF(u) \ WF(f) is contained in $p^{-1}(0)$ and is invariant under the flow of H_p .

The operator P is of real principal type if it additionally satisfies a nontrapping assumption: no bicharacteristic curve stays over a compact subset of X infinitely long. Combined with propagation of singularities, this implies that any solution of $Pu \in C^{\infty}$ in X with a compact singular set has to be smooth.

Remark 2.1. Several different conditions related to real principal type operators appear in the literature, such as $d_{\xi}p \neq 0$, or that dp and the canonical 1-form $\xi_j dx^j$ are linearly independent (equivalently, H_p is not *radial*, i.e. proportional to the radial direction $\xi_j \partial_{\xi_j}$). We wish to clarify the relations between these conditions. Clearly

$$d_{\xi}p \neq 0 \implies H_p$$
 is not radial.

Moreover, if $d_{\xi}p(x_0,\xi) \neq 0$ then the bicharacteristic through (x_0,ξ) moves away from x_0 . Thus

$$d_{\xi}p \neq 0$$
 at any $(x_0,\xi) \in p^{-1}(0) \implies P$ is real principal type near x_0 .

In the converse direction,

P is real principal type $\implies H_p$ is not radial anywhere on $p^{-1}(0)$,

since if H_p were radial at some $(x_0, \xi_0) \in p^{-1}(0)$, then there would be a bicharacteristic $\gamma(t) = (x_0, \xi(t))$ staying over x_0 for infinite time. The Tricomi operator $P = x_2 \partial_1^2 + \partial_2^2$ is of real principal type, but does not satisfy $d_{\xi}p \neq 0$ on $p^{-1}(0)$ (in fact null bicharacteristic curves may have cusps).

We will next give a solvability result for real principal type operators. This is a direct adaptation of [Hö85, Theorem 26.1.7], but we will state it in a form adapted to manifolds with boundary.

Let us introduce some notation for Sobolev spaces and distributions. Let X be an open manifold and let M be a compact subdomain with boundary. We fix some Riemannian metric on X with volume form dV, and consider Sobolev spaces $H^s(M)$ and $H^s_{loc}(X)$ with respect to the L^2 norm induced by dV. Here we slightly abuse notation and write $H^s(M)$ instead of $H^s(M^{int})$, where the latter space can be defined as the restriction of $H^s_{loc}(X)$ to M^{int} .

The distribution space $\mathscr{D}'(X)$ is the set of continuous linear functionals on $C_c^{\infty}(X)$, and we identify functions $u \in L^1_{loc}(X)$ with distributions using the pairing

$$(u, \varphi) = \int_X u \bar{\varphi} \, dV, \qquad \varphi \in C^\infty_c(X).$$

We continue to write (u, φ) for the distributional pairing. Here we deviate from standard practice by requiring that the pairing is conjugate linear in the second variable. This convention will be helpful below.

Write $E_M(X) = E_M$ for those elements in some space E(X) that are supported in M. With respect to the distributional pairing, one has for any $s \in \mathbb{R}$ the duality statements (see e.g. [AS19, Section 2])

$$(H^s(M))^* = H_M^{-s}, \qquad (H_M^s)^* = H^{-s}(M), \qquad (C^{\infty}(M))^* = \mathscr{D}'_M.$$

We also write (u, φ) for the pairings for spaces on M. Any differential operator P on M induces a map $P: \mathscr{D}'_M \to \mathscr{D}'_M$ via

$$(Pv,\varphi) = (v, P^*\varphi)$$

where P^* is the formal L^2 adjoint of P. Finally, given a set $S \subset C_M^\infty$, we consider its annihilator

$$S^{\perp} = \{ v \in \mathscr{D}'_M ; (v, \varphi) = 0 \text{ for all } \varphi \in S \}.$$

Proposition 2.2. Let P be a real principal type differential operator of order $m \ge 1$ on a compact manifold M with smooth boundary, and let $s \in \mathbb{R}$.

- (a) If $u \in \mathscr{D}'_M$ solves Pu = f in X where $f \in H^s_M$, then $u \in H^{s+m-1}_M$.
- (b) The set

$$N(P) = \{ v \in \mathscr{D}'_M ; Pv = 0 \}$$

is a finite dimensional space contained in C_M^{∞} .

(c) Define the space

$$Y_s = \{ v \in \mathscr{D}'_M ; P^* v \in H^s_M \}.$$

Then $Y_s \subset H_M^{s+m-1}$, and for some $C_s > 0$ one has the estimates

(2.1)
$$\|v\|_{H^{s+m-1}_M} \le C_s(\|P^*v\|_{H^s_M} + \|v\|_{H^{s+m-2}_M}), \qquad v \in Y_s,$$

and

(2.2)
$$\|v\|_{H^{s+m-1}_M} \leq C_s \|P^*v\|_{H^s_M}, \quad v \in Y_s \cap N(P^*)^{\perp}.$$

(d) Given any $f \in H^s(M) \cap N(P^*)^{\perp}$, there is a solution $u \in H^{s+m-1}(M)$ of the equation

$$Pu = f \text{ in } M.$$

Moreover, there is a bounded linear operator $E_s: H^s(M) \cap N(P^*)^{\perp} \to H^{s+m-1}(M)$ such that $PE_sf = f$.

Proof. Parts (a) and (b) follow from [Hö85, Theorems 26.1.4 and 26.1.7], after extending P as a real principal type operator to a slightly larger open manifold X. To prove part (c), note that by part (a) one has $Y_s \,\subset \, H_M^{s+m-1}$, and Y_s with norm $\|v\|_{Y_s} = \|v\|_{H_M^{s+m-2}} + \|P^*v\|_{H_M^s}$ becomes a Hilbert space. The inclusion map $\iota: Y \to H_M^{s+m-1}$ has closed graph, and the closed graph theorem yields (2.1). This in turn implies (2.2). Indeed, if (2.2) were false one would have a sequence (v_j) in $Y_s \cap N(P^*)^{\perp}$ with $\|v_j\|_{H_M^{s+m-1}} = 1$ and $\|P^*v_j\|_{H_M^s} \to 0$, and by compact Sobolev embedding some subsequence would converge strongly in H_M^{s+m-2} to some $v \in H_M^{s+m-2} \cap N(P^*)^{\perp}$ with $P^*v = 0$. Then (2.1) would give $\|v\|_{H_M^{s+m-2}} \ge 1/C$, and we would obtain a contradiction since v would be a nonzero element of both $N(P^*)$ and $N(P^*)^{\perp}$.

Finally, we prove part (d). Given $f \in H^s(M) \cap N(P^*)^{\perp}$, we define the (conjugate) linear functional

$$\ell_f: P^*C^\infty_M \to \mathbb{C}, \ \ell_f(P^*v) = (f,v) \text{ for } v \in C^\infty_M.$$

This is well defined by the assumption on f. One has

$$|(f,v)| \le \|f\|_{H^s(M)} \|v\|_{H^{-s}_M} \le C \|f\|_{H^s(M)} \|P^*v\|_{H^{-(s+m-1)}_M}, \qquad v \in C^\infty_M,$$

since this holds for $v \in C_M^{\infty} \cap N(P^*)^{\perp}$ by (2.2) (with *s* replaced by -s - m - 1) and both sides remain the same when a function in $N(P^*)$ is added to *v*. Thus, denoting by *W* the closure of $P^*C_M^{\infty}$ in $H_M^{-(s+m-1)}$, ℓ_f extends uniquely to a bounded functional on *W* such that

$$|\ell_f(w)| \le C ||f||_{H^s(M)} ||w||_{H^{-(s+m-1)}_M}, \qquad w \in W$$

Let Q be the orthonormal projection from ${\cal H}_M^{-(s+m-1)}$ to W, and define

$$\bar{\ell}_f: H_M^{-(s+m-1)} \to \mathbb{C}, \ \bar{\ell}_f(u) = \ell_f(Qu).$$

Then $\bar{\ell}_f$ is a bounded linear functional on $H_M^{-(s+m-1)}$, and it depends linearly on f by construction. By duality there exists $u_f \in H^{s+m-1}(M)$ with $\|u_f\|_{H^{s+m-1}(M)} \leq C \|f\|_{H^s(M)}$ such that

 $(u_f, P^*v) = (f, v), \qquad v \in C_M^{\infty}.$

In particular $Pu_f = f$ in M^{int} . Letting $E_s f = u_f$ concludes the proof.

Semiclassical wave front sets. Next, we will consider the action of real principal type operators on functions depending on a small parameter h > 0. We recall certain semiclassical notions following [Zw12]. A family $u = u_h \subset L^2_{loc}(X)$, $0 < h \leq 1$, is called L^2 -tempered if for any $\chi \in C^{\infty}_c(X)$ there is $N \geq 0$ so that

$$\|\chi u_h\|_{L^2} = O(h^{-N})$$

as $h \to 0$. We say that u_h is semiclassically smooth at $(x_0, \xi_0) \in T^*X$ if, in some local coordinates near x_0 , there is $\varphi \in C_c^{\infty}$ with $\chi = 1$ near x_0 and $\psi \in C_c^{\infty}$ with $\psi = 1$ near ξ_0 so that

$$\psi \mathscr{F}_h(\varphi u_h) = O_{L^2}(h^\infty)$$

where \mathscr{F}_h is the semiclassical Fourier transform,

$$\mathscr{F}_h f(\xi) = (2\pi h)^{-n} \int_{\mathbb{R}^n} e^{-i\frac{x\cdot\xi}{h}} f(x) \, dx.$$

The definition is independent of the choice of local coordinates. The semiclassical wave front set $WF_{scl}(u_h)$ is the complement of those points $(x_0, \xi_0) \in T^*X$ at which u_h is semiclassically smooth. Roughly, one can think of $WF_{scl}(u_h)$ as the set where the family u_h is localized in phase space as $h \to 0$.

We will need the following characterization of the semiclassical wave front set in terms of FBI (Fourier-Bros-Iagolnitzer) transforms. This is a local statement and reduces to the corresponding result in Euclidean space [Zw12, Theorem 13.14].

Proposition 2.3. Let $u = u_h$ be L^2 -tempered in X. One has $(x_0, \xi_0) \notin WF_{scl}(u)$ if and only if

$$\int_X u(y)\overline{e^{i\Psi(y;x,\xi)/h}b(y)} \, dV(y) = O(h^\infty)$$

uniformly over (x,ξ) near (x_0,ξ_0) , where $\Psi = \Psi(\cdot;x,\xi)$ and $b = b(\cdot)$ are smooth functions in X so that, in some local coordinates near x_0 ,

$$\Psi(y; x, \xi) = \xi \cdot (y - x) + \frac{i}{2}|y - x|^2, \quad \text{supp}(b) \text{ is close to } x_0, \quad b = 1 \text{ near } x_0.$$

If u_h and v_h are L^2 -tempered, the semiclassical wave front set has the following simple properties:

• If u_h and v_h are semiclassically smooth at (x_0, ξ_0) , then so is $u_h + v_h$. Thus

$$\operatorname{WF}_{\operatorname{scl}}(u_h) \triangle \operatorname{WF}_{\operatorname{scl}}(v_h) \subset \operatorname{WF}_{\operatorname{scl}}(u_h + v_h) \subset \operatorname{WF}_{\operatorname{scl}}(u_h) \cup \operatorname{WF}_{\operatorname{scl}}(v_h),$$

where $A \triangle B = (A \setminus B) \cup (B \setminus A)$.

• If $\|\chi v_h\|_{L^2} = O(h^{\infty})$ for $\chi \in C_c^{\infty}(X)$, then $\operatorname{WF}_{\operatorname{scl}}(v_h) = \emptyset$ and $\operatorname{WF}_{\operatorname{scl}}(u_h + v_h) = \operatorname{WF}_{\operatorname{scl}}(u_h)$.

The following result is a semiclassical version of the propagation of singularities theorem.

Proposition 2.4. Let P be a real principal type differential operator in an open manifold X, and let u_h be L^2 -tempered. If

$$Pu_h = f_h \ in \ X$$

where f_h is L^2 -tempered, then $WF_{scl}(u_h) \setminus WF_{scl}(f_h)$ is invariant under the Hamiltonian flow of the principal symbol of P.

Proof. Define $P_h := h^m P$, so that the semiclassical principal symbol of P_h is precisely the same as the classical principal symbol of P. Then $P_h u_h = h^m f_h$, and the result follows from [Zw12, Theorem 12.5].

Cauchy data sets. Next we will discuss Cauchy data sets, starting with three basic invariance properties.

Lemma 2.5. Let P be a differential operator of order m, and let $c \in C^{\infty}(M)$ be nonvanishing. Then

 $C_P = C_{cP}.$ If additionally $c|_{\partial M} = 1$ and $\nabla^j c|_{\partial M} = 0$ for $1 \le j \le m - 1$, then

$$C_P = C_{P+c^{-1}[P,c]}_{17}$$

Moreover, if $\Phi: M \to M$ is a diffeomorphism such that Φ and Id_M agree to order m-1 on ∂M , then

$$C_P = C_{\Phi^* P}.$$

Here Φ^*P is the differential operator on M defined via $\Phi^*P(\Phi^*v) = \Phi^*(Pv)$ for $v \in C^{\infty}(M)$.

Proof. The first part is clear, and the second part follows from the formula

$$P(cv) = cPv + [P, c]v$$

The third part holds since u and Φ^*u agree to order m-1 on ∂M by the assumption on Φ . \Box

It is instructive to verify how these invariances affect the scattering relation and bicharacteristic ray transforms. The next lemma shows that Theorem 1.1 is consistent with the above invariances. It also shows that if M is simply connected, then changing P to $P + c^{-1}[P, c]$ changes the subprincipal symbol by a term $\partial_{\xi_a} p_m \partial_a \varphi$ for some $\varphi \in C^{\infty}(M)$ with $\varphi|_{\partial M} = 0$. If P is a second order operator and A is a 1-form encoding the first order coefficients, this corresponds to the standard invariance $A \to A + d\varphi$ obtained by conjugating P with $e^{\pm i\varphi}$. See e.g. [DKSU09, SY18] for the Riemannian and Lorentzian cases. However, if M has nontrivial topology, there is a more general invariance (corresponding to $A \to A + d\varphi + h$ in the second order case) described by harmonic 1-forms h whose integrals over closed loops are in $2\pi i \mathbb{Z}$. This is related to the Aharonov-Bohm effect where in fact the Cauchy data sets may differ if the integrals of h over closed loops are not in $2\pi i \mathbb{Z}$, see [AB59] and the review [Es15]. This fact also explains the nonuniqueness modulo $2\pi \mathbb{Z}$ in Theorem 1.1.

Lemma 2.6. All three invariances in Lemma 2.5 preserve the scattering relation α_P (for the third one one needs to assume $m \geq 2$). Moreover, if M is connected and if $c \in C^{\infty}(M, \mathbb{C})$ satisfies $c|_{\partial M} = 1$, then

(2.3)
$$\sigma_{\rm sub}[(P+c^{-1}[P,c])^{\mu}] = \sigma_{\rm sub}(P^{\mu}) - i\sum_{a=1}^{n} \partial_{\xi_a} p_m(\partial_a \varphi + h_a)$$

for some $\varphi \in C^{\infty}(M,\mathbb{C})$ with $\varphi|_{\partial M} = 0$, and for some harmonic 1-form $h = h_a dx^a$ on M (with respect to the auxiliary Riemannian metric) whose tangential part vanishes on ∂M and which satisfies

$$\int_{\eta} h \in 2\pi i \mathbb{Z} \qquad \text{for any closed loop } \eta \text{ in } M.$$

The integral of $\sum_{a=1}^{n} \partial_{\xi_a} p_m(\partial_a \varphi + h_a)$ over any null bicharacteristic segment between boundary points is in $2\pi i \mathbb{Z}$.

Proof. Note that if P has real principal symbol and $c \in C^{\infty}(M, \mathbb{R})$ is nonvanishing, then changing P to cP or $P+c^{-1}[P,c]$ does not change the null bicharacteristic flow. Replacing P by Φ^*P changes the null bicharacteristics in the interior, but not at the boundary if Φ and Id_M agree to first order on ∂M . Thus the three invariances above preserve the scattering relation at least when $m \geq 2$.

If P has full symbol with polyhomogeneous expansion $\sum_{j=0}^{m} p_j(x,\xi)$ in some local coordinate system, then the order m-1 term in the corresponding expansion for $P + c^{-1}[P,c]$ is

$$p_{m-1}(x,\xi) - ic^{-1}\sum_{a=1}^n \partial_{\xi_a} p_m(x,\xi) \partial_a c(x).$$

Thus (2.3) follows if we can prove that

$$c^{-1}dc = d\varphi + h$$
18

where φ and h are as stated. This is just the Hodge decomposition for the 1-form $c^{-1}dc$, which can be obtained by using the auxiliary Riemannian metric g on M and solving

$$-\Delta_g \varphi = \delta_g(c^{-1}dc)$$
 in $M, \qquad \varphi|_{\partial M} = 0.$

It follows that $h := c^{-1}dc - d\varphi$ is harmonic (i.e. $dh = \delta_g h = 0$) with vanishing tangential part on ∂M . If η is a closed loop in M, we have

$$\int_{\eta} h = \int_{\eta} c^{-1} dc = \int_{c \circ \eta} \frac{dz}{z}$$

The last quantity is $2\pi i$ times the winding number of the curve $c \circ \eta$ in $\mathbb{C} \setminus \{0\}$, and hence belongs to $2\pi i\mathbb{Z}$.

Finally, using that M is connected, we may fix $x_0 \in \partial M$ and define the function

$$\psi: M \to \mathbb{C}/2\pi i\mathbb{Z}, \ \psi(x) = \int_{\eta_{x_0,x}} h \mod 2\pi i\mathbb{Z}$$

where $\eta_{x_0,x}$ is any smooth curve from x_0 to x in M. This is well defined by the condition on h, and $e^{\pm\psi}$ are well defined smooth functions $M \to \mathbb{C} \setminus \{0\}$. Moreover,

$$d(ce^{-\varphi}e^{-\psi}) = (c^{-1}dc - d\varphi - h)ce^{-\varphi}e^{-\psi} = 0.$$

Thus $ce^{-\varphi}e^{-\psi}$ is a constant. Evaluating at x_0 gives that $c = e^{\varphi}e^{\psi}$ and that $e^{\psi}|_{\partial M} = ce^{-\varphi}|_{\partial M} = 1$, i.e. $\psi(y) = 0$ modulo $2\pi i\mathbb{Z}$ whenever $y \in \partial M$. Now, if $\gamma : [0,T] \to M$, $\gamma(t) = (x(t),\xi(t))$ is a null bicharacteristic segment between boundary points, the fact that $\partial_{\xi_a} p_m(x(t),\xi(t)) = \dot{x}_a(t)$ yields

$$\int_0^T (\partial_{\xi_a} p_m \partial_a \varphi)(\gamma(t)) \, dt = \int_0^T \frac{d}{dt}(\varphi(x(t))) \, dt = 0$$

since $\varphi|_{\partial M} = 0$. For the part involving h we have

$$\int_0^T (\partial_{\xi_a} p_m h_a)(\gamma(t)) \, dt = \int_0^T h(\dot{x}(t)) \, dt.$$

Now we may write $\psi(x(t)) = \int_{\eta_{x_0,x(0)}} h + \int_0^t h(\dot{x}(s)) ds$ modulo $2\pi i \mathbb{Z}$, leading to

$$\int_0^T (\partial_{\xi_a} p_m h_a)(\gamma(t)) dt = \psi(x(T)) - \psi(x(0)) = 0 \text{ modulo } 2\pi i \mathbb{Z}$$

since $\psi(y) = 0$ modulo $2\pi i\mathbb{Z}$ for $y \in \partial M$.

Note also that the gauge invariances in Lemma 2.5 are formulated in a different way compared to the usual invariances in the Calderón or Gel'fand problems for second order operators. For instance, if \bar{g} is a Riemannian metric, A is a 1-form and q is a function on M, the corresponding second order operator (written in local coordinates) in the Calderón problem is

$$Pu = |\bar{g}|^{-1/2} (D_j + A_j) (|\bar{g}|^{1/2} \bar{g}^{jk} (D_k + A_k)u) + qu.$$

In this case one typically defines a normal derivative with respect to \bar{g} and A, leading to the Cauchy data set

$$\hat{C}_P = \{ (u|_{\partial M}, (du + iAu)(\nu_{\bar{g}})|_{\partial M} ; Pu = 0 \text{ in } M \},\$$

see [DKSU09]. The Cauchy data set in this article is instead defined in terms of a (known) reference Riemannian metric g on M as

$$C_P = \{ (u|_{\partial M}, \nabla_g u|_{\partial M}); Pu = 0 \text{ in } M \}.$$
¹⁹

Of course, if \bar{g} and A are known on ∂M , then C_P determines \tilde{C}_P and vice versa. However, the difference between C_P and \tilde{C}_P implies for instance that there is no analogue of the gauge invariance $A \to A + d\psi$ with $\psi|_{\partial M} = 0$ in our boundary determination result for the subprincipal symbol.

Finally, we discuss a simple but fundamental integral identity. We first give a characterization of the inclusion $C_{P_1} \subset C_{P_2}$ in terms of an estimate for a certain inner product.

Lemma 2.7. Let P_1 and P_2 be differential operators of order m. One has $C_{P_1} \subset C_{P_2}$ if and only if for any $u_1 \in H^m(M)$ solving $P_1u_1 = 0$ in M, there is $C_{u_1} > 0$ so that

(2.4)
$$|((P_1 - P_2)u_1, v)_{L^2(M)}| \le C_{u_1} \|P_2^*v\|_{H^{-m}(M)}, \qquad v \in H^m(M).$$

Proof. Assume that $C_{P_1} \subset C_{P_2}$ and that $u_1 \in H^m(M)$ solves $P_1u_1 = 0$ in M. Then we can find $\tilde{u}_2 \in H^m(M)$ with $P_2\tilde{u}_2 = 0$ in M so that $u_1 - \tilde{u}_2 \in H_0^m(M)$. For any $v \in H^m(M)$, one has in terms of $L^2(M)$ inner products

$$((P_1 - P_2)u_1, v) = -(P_2u_1, v) = -(P_2(u_1 - \tilde{u}_2), v) = -(u_1 - \tilde{u}_2, P_2^*v).$$

The inequality (2.4) follows from the duality of $H_0^m(M)$ and $H^{-m}(M)$.

Conversely, let $u_1 \in H^m(M)$ solve $P_1u_1 = 0$, and assume that (2.4) holds. Then $f := -P_2u_1$ satisfies

(2.5)
$$|(f,v)| \le C \|P_2^*v\|_{H^{-m}(M)}, \qquad v \in H^m(M).$$

Define a linear functional

$$\ell_f : P_2^* H^m(M) \to \mathbb{C}, \ \ell_f(P_2^* v) = (f, v).$$

This is well defined and bounded with respect to the $H^{-m}(M)$ norm by (2.5). By the Hahn-Banach theorem ℓ_f extends continuously to $H^{-m}(M)$, and by duality there is $w \in H_0^m(M)$ so that

$$(f, v) = \ell_f(P_2^* v) = (w, P_2^* v), \qquad v \in H^m(M).$$

It follows that $P_2w = f = -P_2u_1$. Thus $\tilde{u}_2 := u_1 + w$ solves $P_2\tilde{u}_2 = 0$ with $u_1 - \tilde{u}_2 \in H_0^m(M)$. This proves that $C_{P_1} \subset C_{P_2}$.

If we choose v in (2.4) to be a solution of $P_2^*v = 0$, we immediately obtain an integral identity from the condition $C_{P_1} = C_{P_2}$. For our applications we indeed need $P_1 - P_2$ to appear on the left hand side, and thus one of the functions needs to solve the adjoint equation.

Lemma 2.8. If $C_{P_1} = C_{P_2}$, then

$$((P_1 - P_2)u_1, u_2)_{L^2(M)} = 0$$

whenever $u_j \in H^m(M)$ solve $P_1u_1 = P_2^*u_2 = 0$ in M.

3. QUASIMODE CONSTRUCTION

We will now proceed to the construction of approximate solutions that concentrate near an injective segment of a null bicharacteristic curve. This can be done for any operator with real principal symbol.

Theorem 3.1. Let P be a differential operator of order m on a smooth manifold X, with real valued principal symbol p_m . Let $\gamma : [0,T] \to T^*X \setminus 0$ be a segment of a null bicharacteristic curve. If γ is injective on [0,T], there is a family $u = u_h \in C_c^{\infty}(X)$ for $0 < h \leq 1$ such that

$$WF_{scl}(u) = \gamma([0,T]), \qquad WF_{scl}(Pu) = \gamma(0) \cup \gamma(T).$$

Moreover, if \tilde{P} is another differential operator on X with the same principal symbol as P and if $v = v_h$ is the analogous family for \tilde{P}^* , then for any differential operator Q of order $\ell \geq 0$ on X whose coefficients vanish near the end points of γ one has

(3.1)
$$\lim_{h \to 0} h^{-\frac{n+1}{2}+\ell} \int_X (Qu)\bar{v} \, dV = c_0 \int_0^T q_\ell(\gamma(t)) \exp\left[-i \int_0^t (p_{m-1} - \tilde{p}_{m-1})(\gamma(s)) \, ds\right] \, dt$$

where $c_0 \neq 0$ is the constant in (3.37), q_ℓ is the principal symbol of Q, and $\sum_{j=0}^m p_j$ and $\sum_{j=0}^m \tilde{p}_j$ are the polyhomogeneous full symbols of P and \tilde{P} in a local coordinate system.

Note that the statement about $WF_{scl}(Pu)$ implies that $Pu = O(h^{\infty})$ away from the end points of γ , and hence u is an approximate solution for P. The formula (3.1) is related to the semiclassical limit measure for the family u. In fact, one could also prove that for any $a \in C_c^{\infty}(T^*X)$ vanishing near the end points of γ one has

$$\lim_{h \to 0} h^{-\frac{n+1}{2}} (\operatorname{Op}_h(a)u_h, u_h)_{L^2(X)} = \int_0^T r_{\gamma}(t) a(\gamma(t)) \, dt$$

where $Op_h(a)$ is the semiclassical Weyl quantization of a, and for any nonvanishing half density μ

$$r_{\gamma}(t) = c_0 \exp\left[-i \int_0^t (\sigma_{\rm sub}[P^{\mu}] - \sigma_{\rm sub}[(P^*)^{\mu}])(\gamma(s)) \, ds\right].$$

The formulation (3.1) will be convenient for our applications, and it does not involve semiclassical quantization. The expression on the right hand side of (3.1) must be coordinate invariant since the left hand side is, however, this can also be seen directly as follows:

Lemma 3.2. Let P and \tilde{P} be two differential operators of order m on X with the same principal symbol, and denote by $\sum_{j=0}^{m} p_j$ and $\sum_{j=0}^{m} \tilde{p}_j$ their polyhomogeneous full symbols in a local coordinate system. Then $p_{m-1} - \tilde{p}_{m-1}$ is an invariantly defined function on $T^*X \setminus 0$.

Proof. Let μ be a half density on X and write $\sum_{j=0}^{m} q_j$ and $\sum_{j=0}^{m} \tilde{q}_j$ for the polyhomogeneous full symbols of P^{μ} and \tilde{P}^{μ} in a local coordinate system. Then $q_m = p_m$ and

$$q_{m-1} = p_{m-1} - i\mu \partial_{\xi_j} p_m \,\partial_{x_j} \mu^{-1}$$

Analogous statements hold for \tilde{P}^{μ} , and recalling the definition of subprincipal symbol in (1.2), we see that the function

(3.2)
$$\sigma_{\rm sub}[P^{\mu}] - \sigma_{\rm sub}[\tilde{P}^{\mu}] = p_{m-1} - \tilde{p}_{m-1}$$

is defined invariantly on $T^*X \setminus 0$.

In the following, we will assume that P has order $m \ge 1$ and its principal symbol is denoted by $p_m \in C^{\infty}(T^*X)$. We will write in local coordinates

$$\gamma(t) = (x(t), \xi(t))$$

so that $(x(t),\xi(t))$ satisfies the Hamilton equations

$$\dot{x}(t) = d_{\xi} p_m(x(t), \xi(t)), \qquad \dot{\xi}(t) = -d_x p_m(x(t), \xi(t)).$$

More invariantly, we define $x(t) = \pi(\gamma(t))$ where $\pi: T^*X \to X$ is the natural projection.

We will use the following standard result related to conjugating a differential operator by exponentials. This is the "fundamental asymptotic expansion lemma" stated in [Ta81, Section VIII.7] or [Tr80, Section VI.3] for pseudodifferential operators and real valued phase functions, and in [Tr80, Section X.4] for complex valued phase functions. In our case of differential operators, the proof is just an elementary computation in local coordinates.

Lemma 3.3. Let Φ be a smooth real or complex valued function. Then

$$e^{-i\Phi/h}P(e^{i\Phi/h}u) = \sum_{j=0}^{m} h^{j-m}R_ju, \qquad h > 0,$$

where each R_j is a differential operator of order j. In particular, $R_1 u = \frac{1}{i}Lu + bu$ where L is a (possibly complex) vector field and $b = b_P$ is a function. In local coordinates one has

$$R_{0}u = p_{m}(x, \nabla\Phi(x))u,$$

$$Lu = \partial_{\xi_{j}}p_{m}(x, \nabla\Phi(x))\partial_{j}u,$$

$$b = \frac{1}{2i}\partial_{\xi_{j}\xi_{k}}p_{m}(x, \nabla\Phi(x))\partial_{jk}\Phi(x) + p_{m-1}(x, \nabla\Phi(x))$$

where p_{m-1} is the order m-1 term in the polyhomogeneous expansion of the full symbol of P in these coordinates.

There are several methods for constructing quasimodes concentrating near an injective null bicharacteristic segment $\gamma : [0,T] \to T^*X \setminus 0$ where $\gamma(t) = (x(t),\xi(t))$. We first give a brief discussion of such methods, to motivate the construction given in this paper.

Local case. To construct a quasimode u locally near $x(t_0)$ with $\dot{x}(t_0) \neq 0$ it is enough to use the geometrical optics ansatz $u = e^{i\varphi/h}a$ where φ is a smooth real valued phase function solving the eikonal equation $p_m(x, d\varphi(x)) = 0$. If $\dot{x}(t_0) \neq 0$ this equation can always be solved locally near $x(t_0)$, but global solutions do not exist in general if there are caustic points (i.e. points where the projection $\Lambda \to X$ fails to have bijective differential where $\Lambda \subset T^*X \setminus 0$ is a Lagrangian manifold associated with the eikonal equation, see [Hö85, Section 6.4]).

Global case without cusps. More generally, if γ is injective on [0, T] and has no cusps in the sense that $\dot{x}(t) \neq 0$ for $t \in [0, T]$, one can use a Gaussian beam type construction. If x(t) is also injective on [0, T], this amounts to looking for a quasimode $u = e^{i\Phi/h}a$ where Φ and a are complex valued (more generally if x(t) is not injective the quasimode will be a finite sum of such functions). The phase function Φ is required to solve the eikonal equation only on the bicharacteristic in the sense that

(3.3)
$$p_m(x, d\Phi(x)) = 0$$
 to infinite order on $x([0, T])$.

The construction of Φ and a can be carried out very explicitly in global Fermi type coordinates near x([0,T]). This boils down to solving one nonlinear ODE (matrix Riccati equation) for the Hessian of Φ on x([0,T]), and linear ODEs for the derivatives of Φ and a along x([0,T]).

These Gaussian beam type constructions are very classical and go back at least to [BL67, Hö71] with further treatments e.g. in [Ra82, Ra01, KKL01]. A version of this construction is also given in [Hö85, Section 24.2], where the existence of the required complex phase function is explained in terms of properties of complex Lagrangian planes in the complexification of T^*X . However, the construction always breaks down when the bicharacteristic has a cusp. In fact, if (3.3) holds

near $x(t_0)$ where $\dot{x}(t_0) = 0$, then looking at first order derivatives in (3.3) and using the Hamilton equations implies that

$$0 = -\dot{\xi}(t_0) + \Phi''(x(t_0))\dot{x}(t_0) = -\dot{\xi}(t_0).$$

Hence one would have $dp_m(\gamma(t_0)) = 0$ and $\gamma(t) \equiv \gamma(t_0)$, which contradicts the assumption that γ is injective on [0, T].

We mention also the related construction of the propagator for hyperbolic equations as a global oscillatory integral with a complex-valued phase function [LSV94, SV97, CLV19].

General case. Let us now assume that γ is injective on [0,T] but $\dot{x}(t)$ may vanish. In the classical case (i.e. without a parameter), a construction of a quasimode u associated with γ such that $Pu \in C^{\infty}$ was given in [DH72] and [Hö85, Section 26.1]. The argument proceeds by constructing a canonical transformation χ near $\gamma([0,T])$ that straightens γ in phase space into the curve $\eta(x_1) = ((x_1,0), e_n)$ in $T^*\mathbb{R}^n$, and by quantizing χ using suitable amplitudes to obtain Fourier integral operators A and B such that BPA roughly corresponds to D_{x_1} microlocally near $\gamma([0,T])$. One then constructs an explicit quasimode U for D_{x_1} in \mathbb{R}^n associated with η , and u = AU will be the required quasimode for P. This phase space construction is not affected by the presence of cusps.

A semiclassical version of the above construction could be used to prove Theorem 3.1. However, we will give a direct proof based on a modification of the Gaussian beam construction. This construction is motivated by the fact that a standard semiclassical quasimode U for D_{x_1} associated with η ,

$$U(x_1, x') = e^{ix_n/h - |x'|^2/h}$$

can be thought of as a superposition of Gaussian wave packets along η . If A is a semiclassical Fourier integral operator quantizing χ , then u = AU would be a superposition of Gaussian wave packets along the curve x(t). We will thus look for the quasimode u directly in the form

$$u = \int_0^T e^{i\Phi(x,t)/h} a(x,t) \, dt$$

where Φ and a are smooth complex valued functions in $M \times [0, T]$, and each $e^{i\Phi(\cdot, t)/h}a(\cdot, t)$ is a Gaussian wave packet at x(t) oscillating in direction $\xi(t)$. The same idea appears in [PU93, KV98]. The phase function Φ will be chosen to satisfy

(3.4)
$$p_m(\cdot, d_x \Phi(\cdot, t)) + \partial_t \Phi(\cdot, t) = 0 \text{ to infinite order at } x(t) \text{ for } t \in [0, T].$$

This generalizes (3.3) to the case where Φ may depend on t. The construction boils down to solving the same matrix Riccati equation as in the usual Gaussian beam construction, but it is not affected by the presence of cusps. By (3.4) we have that $p_m(x, d_x \Phi(x, t))$ is small near x([0, T]) modulo terms of the form $\partial_t \Phi(x, t)$, but such terms can be dealt with using integration by parts in the formula for u.

After this motivating discussion, we will give a proof of the theorem. This will be done in detail since we will need to use the precise form of the quasimodes, including formulas within the proof, in the computations required for studying the inverse problems.

Proof of Theorem 3.1. The proof will be carried out in several steps.

Step 1. Setup.

We look for an approximate solution of Pu = 0 in the form

(3.5)
$$u(x) = \int_0^T e^{i\Phi(x,t)/h} a(x,t) dt$$

where Φ and a are smooth complex valued functions in $X \times [0, T]$. By Lemma 3.3, we have

(3.6)
$$Pu(x) = \int_0^T e^{i\Phi(x,t)/h} \left[h^{-m} p_m(x, d_x \Phi(x, t)) a(x, t) + h^{1-m} (\frac{1}{i} La + ba)(x, t) + \sum_{j=2}^m h^{j-m} R_j a(x, t) \right] dt.$$

We wish to find $\Phi(x,t)$ so that

(3.7)
$$\Phi(x(t),t) = 0, \qquad d_x \Phi(x(t),t) = \xi(t),$$

and so that

(3.8)
$$p_m(\cdot, d_x \Phi(\cdot, t)) + \partial_t \Phi(\cdot, t) = 0 \text{ to infinite order at } x(t)$$

Moreover, we wish to find $a(x,t) \sim \sum_{j=0}^{\infty} h^j a_j(x,t)$ so that each a_j is smooth and independent of h, and one has the transport equations

(3.9)
$$\frac{1}{i}(\partial_t + L)a_0 + ba_0 = 0 \text{ to infinite order at } x(t),$$

(3.10)
$$\frac{1}{i}(\partial_t + L)a_1 + ba_1 = -R_2a_0 \text{ to infinite order at } x(t),$$

:

(3.11)
$$\frac{1}{i}(\partial_t + L)a_2 + ba_2 = -R_3a_0 - R_2a_1 \text{ to infinite order at } x(t),$$

Step 2. Local construction of Φ .

1

Fix $t_0 \in [0, T]$. We first construct Φ in a small neighborhood of $(x(t_0), t_0)$. Computing in local coordinates, we have (writing $\partial_{x_j} p_m = \partial_{x_j} p_m(x, \nabla_x \Phi(x, t))$ etc):

(3.12)
$$\partial_{x_j}(p_m(x,\nabla_x\Phi)) = \partial_{x_j}p_m + \partial_{\xi_a}p_m\partial_{x_ax_j}\Phi,$$

$$(3.13) \qquad \qquad \partial_{x_j x_k} (p_m(x, \nabla_x \Phi)) = \partial_{x_j x_k} p_m + \partial_{x_j \xi_a} p_m \partial_{x_a x_k} \Phi + \partial_{x_k \xi_a} p_m \partial_{x_a x_j} \Phi + \partial_{\xi_a \xi_b} p_m \partial_{x_a x_j} \Phi \partial_{x_b x_k} \Phi + \partial_{\xi_a} p_m \partial_{x_a x_j x_k} \Phi,$$

(3.14)
$$\partial_x^{\gamma}(p_m(x,\nabla_x\Phi)) = \partial_{\xi_a} p_m \partial_{x_a} \partial_x^{\gamma} \Phi + F^{\gamma}(\nabla^{|\gamma|-1}p_m,\nabla_x^{|\gamma|-1}\Phi)\nabla_x^{|\gamma|}\Phi + G^{\gamma}(\nabla^{|\gamma|}p_m,\nabla_x^{|\gamma|-1}\Phi), \quad |\gamma| \ge 3.$$

In the last statement (which is easily proved by induction) $\nabla^k f$ collects all derivatives of f up to order k, and F^{γ} and G^{γ} are polynomials in their arguments.

Motivated by (3.7), we look for Φ locally having the form

$$\Phi(x,t) = \xi(t) \cdot (x - x(t)) + \frac{1}{2}H(t)(x - x(t)) \cdot (x - x(t)) + \Phi_3(x,t)$$

where H(t) is some complex symmetric matrix depending smoothly on t, and $\Phi_3(\cdot, t)$ vanishes to third order at x(t). Then

(3.15)
$$\Phi(x(t),t) = 0, \qquad \nabla_x \Phi(x(t),t) = \xi(t), \qquad \partial_{x_j x_k} \Phi(x(t),t) = H_{jk}(t).$$

One has $\xi(t) \cdot \dot{x}(t) = 0$ by the Hamilton equations and homogeneity. Thus it also follows that (3.16) $\partial_t \Phi(x,t) = (\dot{\xi}(t) - H(t)\dot{x}(t)) \cdot (x - x(t)) + G(x,t)(x - x(t)) \cdot (x - x(t))$

for some matrix valued function G(x, t).

We first show that (3.8) holds to first order. By (3.15) and (3.16), one has

$$p_m(x, \nabla_x \Phi) + \partial_t \Phi(x, t)\big|_{x=x(t)} = 0$$

We next use (3.12), the Hamilton equations and (3.15) to obtain

$$\partial_{x_j}(p_m(x,\nabla_x\Phi))\Big|_{x=x(t)} = -\dot{\xi}_j(t) + H_{aj}(t)\dot{x}_a(t)$$

By (3.16) we also have

$$\partial_{x_j}(\partial_t \Phi(x,t))\big|_{x=x(t)} = \dot{\xi}_j(t) - H_{ja}(t)\dot{x}_a(t)$$

and therefore (3.8) holds to first order.

To show that (3.8) holds to second order, we rewrite (3.13) as

$$\partial_{x_j x_k} (p_m(x, \nabla_x \Phi)) \Big|_{x=x(t)} = (D + BH + HB^t + HCH)_{jk}(t) + \dot{x}_a(t) \partial_{x_a x_j x_k} \Phi(x(t), t)$$

where B(t), C(t) and D(t) are the matrices

$$(3.17) D_{jk}(t) := \partial_{x_j x_k} p_m(\gamma(t)), B_{ja}(t) = \partial_{x_j \xi_a} p_m(\gamma(t)), C_{ab}(t) = \partial_{\xi_a \xi_b} p_m(\gamma(t)).$$

Moreover, we note that

$$\partial_{x_j x_k} (\partial_t \Phi(x, t)) \big|_{x=x(t)} = \partial_t (\partial_{x_j x_k} \Phi(x(t), t)) - \dot{x}_a(t) \partial_{x_a x_j x_k} \Phi(x(t), t).$$

Adding these two identities and using (3.15), we have

(3.18)
$$\partial_{x_j x_k} (p_m(x, \nabla_x \Phi) + \partial_t \Phi(x, t)) \big|_{x=x(t)} = (\dot{H} + HCH + BH + HB^t + D)_{jk}(t).$$

Given any complex symmetric matrix H_0 with $Im(H_0)$ positive definite, the matrix Riccati equation

(3.19)
$$\dot{H} + HCH + BH + HB^{t} + D = 0, \qquad H(t_0) = H_0,$$

has a smooth complex symmetric matrix solution H(t) where Im(H(t)) is positive definite, see [KKL01, Lemma 2.56]. The solution exists globally in the set where B, C and D are smooth, i.e. in the whole coordinate patch. Choosing such a solution ensures that (3.8) holds to second order.

Finally, let $r \ge 3$ and assume that we have prescribed $\partial_x^\beta \Phi(x(t), t)$ for $|\beta| \le r - 1$ so that (3.8) holds to order r - 1. Let $|\gamma| = r$. Evaluating (3.14) at x(t) gives

$$\partial_x^{\gamma}(p_m(x,\nabla_x\Phi))\big|_{x(t)} = \dot{x}_a(t)\partial_{x_a}\partial_x^{\gamma}\Phi + F^{\gamma}(\nabla^{|\gamma|-1}p_m,\nabla_x^{|\gamma|-1}\Phi)\nabla_x^{|\gamma|}\Phi + G^{\gamma}(\nabla^{|\gamma|}p_m,\nabla_x^{|\gamma|-1}\Phi)\big|_{x(t)}.$$

Moreover, one has

$$\partial_x^{\gamma} \partial_t \Phi(x(t),t) = \partial_t (\partial_x^{\gamma} \Phi(x(t),t)) - \dot{x}_a(t) \partial_{x_a} \partial_x^{\gamma} \Phi(x(t),t)$$

Adding the above two equations gives

$$(3.20) \qquad \partial_x^{\gamma}(p_m(x,\nabla_x\Phi)+\partial_t\Phi)\big|_{x(t)} \\ = \partial_t(\partial_x^{\gamma}\Phi(x(t),t)) + F^{\gamma}(\nabla^{|\gamma|-1}p_m,\nabla_x^{|\gamma|-1}\Phi)\nabla_x^{|\gamma|}\Phi + G^{\gamma}(\nabla^{|\gamma|}p_m,\nabla_x^{|\gamma|-1}\Phi)\big|_{x(t)}.$$

Thus (3.8) will hold to order r provided that

$$\partial_t (\partial_x^{\gamma} \Phi(x(t), t)) + F^{\gamma} (\nabla^{|\gamma| - 1} p_m, \nabla_x^{|\gamma| - 1} \Phi) \nabla_x^{|\gamma|} \Phi + G^{\gamma} (\nabla^{|\gamma|} p_m, \nabla_x^{|\gamma| - 1} \Phi) = 0, \qquad |\gamma| = r.$$

These equations over all γ with $|\gamma| = r$ can be understood as a linear system of ODEs for the vector $(\partial_x^{\gamma} \Phi(x(t), t))_{|\gamma|=r}$, and given any initial data at t_0 this system has a smooth solution. Thus

we have constructed the formal Taylor series of Φ at (x(t), t), and applying Borel summation gives the required function Φ locally.

Step 3. Global construction of Φ .

We wish to glue together the local constructions in Step 2. To do this, we cover [0, T] by finitely many open intervals I_1, \ldots, I_R so that $0 \in I_1, T \in I_R, I_j \cap I_{j+1} \neq \emptyset$, and each $x(I_j)$ is contained in some coordinate patch (U_j, y_j) . We first construct Φ_1 near $x(I_1)$ as in Step 2 with Im(H(0))positive definite. Choosing some $t_1 \in I_1 \cap I_2$, we then construct Φ_2 near $x(I_2)$ as in Step 2 so that the initial data for the ODEs for Φ_2 at $(x(t_1), t_1)$ matches the Taylor series of Φ_1 at $(x(t_1), t_1)$ when Φ_1 is written in the y_2 coordinates.

We claim that one has

(3.21)
$$\Phi_1 = \Phi_2$$
 to infinite order at any $(x(t), t)$ where $t \in I_1 \cap I_2$.

This holds to first order since $\Phi_j(x(t), t) = 0$ and $d_x \Phi_j(x(t), t) = \xi(t)$. For the higher order case, it is enough to use the invariant statements

$$p_m(\cdot, d_x \Phi_j(\cdot, t)) + \partial_t \Phi_j(\cdot, t)|_{x(t)} = 0$$
 to high order for $t \in I_1 \cap I_2$ and $j = 1, 2$

and the formulas (3.18) and (3.20), which imply that in some local coordinates y, both $\partial_y^{\gamma} \Phi_1(x(t), t)$ and $\partial_y^{\gamma} \Phi_2(x(t), t)$ satisfy the same first order ODEs and have the same initial data when $t = t_1$.

Repeating the above process finitely many times, we see that the formal Taylor series of $\Phi(\cdot, t)|_{x(t)}$ varies smoothly with t for $t \in [0, T]$. We can obtain a global smooth function Φ with this Taylor series by a Borel summation scheme that is compatible across coordinate patches. To do this, we use the auxiliary Riemannian metric on X with covariant derivative ∇ . Then the tensor fields

$$T_j(t) = \nabla^j_x \Phi(\cdot, t)|_{x(t)},$$

defined in terms of the formal Taylor series of Φ , are invariantly defined and depend smoothly on $t \in [0,T]$. The map $E(t,v) = (\exp_{x(t)}(v),t)$ for $t \in [0,T]$ and $v \in T_{x(t)}M$ is a local diffeomorphism near each (t,0) and it is injective on $[0,T] \times \{0\}$. Hence it is a diffeomorphism onto a neighborhood of $\{(x(t),t); t \in [0,T]\}$ (see e.g. [KS13, Lemma 7.3]). We may define a smooth function Φ as the Borel sum

$$\Phi(x,t) = \sum_{j=0}^{\infty} \frac{T_j(t)(V(x,t),\dots,V(x,t))}{j!} \chi(|V(x,t)|/\varepsilon_j)$$

where V(x,t) is the projection of $E^{-1}(x,t)$ to the v component, $\chi \in C_c^{\infty}(\mathbb{R})$ satisfies $\chi = 1$ near 0, and ε_j are suitable small numbers (see [Hö85, Theorem 1.2.6]). To check that Φ indeed has the right Taylor series, observe that for small s

$$\Phi(\exp_{x(t)}(sv), t) = \sum_{j=0}^{\infty} \frac{T_j(t)(v, \dots, v)}{j!} s^j \chi(s|v|/\varepsilon_j).$$

If $\eta(s)$ is a geodesic, the formula $\partial_s^j(f(\eta(s)) = \nabla_{\dot{\eta}(s)}(\nabla_{\dot{\eta}(s)}(\cdots(\nabla_{\dot{\eta}(s)}f))) = \nabla^j f(\dot{\eta}(s), \ldots, \dot{\eta}(s))$ shows that Φ defined as above satisfies $\nabla_x^j \Phi(\cdot, t)|_{x(t)} = T_j(t)$.

Step 4. Construction of a.

Recall that we wish to find $a(x,t) \sim \sum_{j=0}^{\infty} h^j a_j(x,t)$ so that a_j solve the transport equations (3.9)–(3.11). Evaluating the first equation at x(t) gives that

$$\frac{1}{i}(\partial_t + L)a_0 + ba_0\big|_{x=x(t)} = \frac{1}{i}\partial_t(a_0(x(t), t)) + b(x(t))a_0(x(t), t).$$

This will vanish provided that

(3.22)
$$a_0(x(t),t) = \exp\left[-i\int_0^t b(x(s))\,ds\right]$$

where, for the sake of definiteness, we have chosen the initial value

$$a(x(0), 0) = 1.$$

Taking first order derivatives in (3.9) gives

$$\partial_{x_j} \left[\frac{1}{i} (\partial_t + L) a_0 + b a_0 \right] \Big|_{x = x(t)} = \frac{1}{i} \partial_t (\partial_{x_j} a_0(x(t), t)) + q_j^k(t) \partial_{x_k} a_0(x(t), t)) + r_j(t) a_0(x(t), t)$$

where q_j^k, r_j are independent of a_0 . Requiring this to vanish leads to a linear ODE system for $(\partial_{x_j} a_0(x(t), t))_{j=1}^n$, which can be solved as above. Continuing this process gives the formal Taylor series of $a_0(\cdot, t)$ at x(t) within a coordinate patch. As in Step 4, covering x([0, T]) by finitely many coordinate patches and solving the ODEs for a_0 in the next coordinate patch with initial data obtained from the previous coordinate patch, we see that the formal Taylor series of $a_0(\cdot, t)|_{x(t)}$ depends smoothly on $t \in [0, T]$. Applying Borel summation as in Step 4 yields a smooth function a_0 satisfying (3.9). The amplitudes a_1, a_2, \ldots are constructed analogously. The support of a can be chosen to be contained in any fixed small neighborhood of $\{(x(t), t); t \in [0, T]\}$.

Step 5. Further properties of Φ .

Here we collect some further properties of the phase function $\Phi(x,t)$ for later use. Again, in order to have coordinate invariant statements, we use the auxiliary Riemannian metric g on X. We also equip $X \times [0,T]$ with the product metric $g \oplus e$, where e is the Euclidean metric on \mathbb{R} .

First we fix the choice for the initial data for H(t) by assuming that the quasimode near x(0) is constructed using Riemannian normal coordinates at x(0), and when H(t) is written in these coordinates one chooses in (3.19) the initial value

$$H(0) = i \mathrm{Id}.$$

Next recall from (3.7) that

(3.23)
$$\Phi(x(t),t) = 0, \qquad d_x \Phi(x(t),t) = \xi(t), \qquad \partial_{x_j x_k} \Phi(x(t),t) = H_{jk}(t).$$

In particular, $\operatorname{Im}(\Phi(\cdot, t))$ vanishes to first order at x(t). Consequently the Riemannian Hessian $\nabla_x^2 \operatorname{Im}(\Phi)|_{(x(t),t)} = (\partial_{x_j x_k} \operatorname{Im}(\Phi) - \Gamma_{jk}^l \partial_{x_l} \operatorname{Im}(\Phi)) dx^j \otimes dx^k$ is equal to $\operatorname{Im}(H_{jk}(t)) dx^j \otimes dx^k$. This shows that $\operatorname{Im}(H(t))$ is an invariantly defined 2-tensor field along x(t), and

$$\nabla_x^2 \operatorname{Im}(\Phi)|_{(x(t),t)} = \operatorname{Im}(H(t)).$$

Now Im(H(0)) is positive definite, and this property is preserved for solutions of the matrix Riccati equation. This shows that there is c > 0 with

$$\nabla_x^2 \operatorname{Im}(\Phi)|_{(x(t),t)}(v,v) \ge c|v|^2, \quad \text{uniformly over } t \in [0,T].$$

Thus if supp(a) is chosen sufficiently small, we will have

(3.24)
$$\operatorname{Im}(\Phi(x,t)) \ge cd(x,x(t))^2, \qquad (x,t) \in \operatorname{supp}(a).$$

We will next consider derivatives of Φ with respect to t. By (3.16) one has

(3.25)
$$\partial_t \Phi(x(t), t) = 0$$

Differentiating (3.16) further, we have

(3.26)
$$d_x \partial_t \Phi(x(t), t) = (\dot{\xi}_j(t) - H_{jk}(t)\dot{x}^k(t)) dx^j$$

which is an invariantly defined 1-form along x(t) since the left hand side is invariant. Similarly

(3.27)
$$\partial_t^2 \Phi(x(t), t) = -(\dot{\xi}_j(t) - H_{jk}(t)\dot{x}^k(t))\dot{x}^j(t)$$

Step 6. Wave front set of Pu.

We now study the wave front set of Pu using the formula (3.6). By (3.8) we have

$$p_m(x, d_x \Phi(x, t)) = -\partial_t \Phi(x, t) + r(x, t)$$

where $r(\cdot, t)$ vanishes to infinite order at x(t). We observe that

$$\int_0^T e^{i\Phi(x,t)/h} h^{-m}(-\partial_t \Phi(x,t)) a(x,t) dt = -h^{-m} \int_0^T \frac{h}{i} \partial_t (e^{i\Phi(x,t)/h}) a(x,t) dt$$
$$= h^{1-m} \int_0^T e^{i\Phi(x,t)/h} \frac{1}{i} \partial_t a(x,t) dt + \frac{1}{i} h^{1-m} (e^{i\Phi(x,0)/h} a(x,0) - e^{i\Phi(x,T)/h} a(x,T))$$

Thus, using the formula for Pu in (3.6), we have

$$Pu(x) = \frac{1}{i}h^{1-m}(e^{i\Phi(x,0)/h}a(x,0) - e^{i\Phi(x,T)/h}a(x,T)) + \int_0^T e^{i\Phi(x,t)/h} \left[h^{-m}(ra)(x,t) + h^{1-m}(\frac{1}{i}(\partial_t a + La) + ba)(x,t) + \sum_{j=2}^m h^{j-m}R_ja(x,t)\right] dt.$$

By (3.9)–(3.11) we further have

$$(3.28) \quad Pu(x) = \frac{1}{i}h^{1-m}(e^{i\Phi(x,0)/h}a(x,0) - e^{i\Phi(x,T)/h}a(x,T)) + \int_0^T e^{i\Phi(x,t)/h} \left[\sum_{j=0}^m h^{j-m}r_j(x,t)\right] dt$$

where each r_j is a smooth function with $\operatorname{supp}(r_j) \subset \operatorname{supp}(a)$ and $r_j(\cdot, t)$ vanishes to infinite order at x(t). By (3.24), for any N there is C_N so that

$$\left|\int_{0}^{T} e^{i\Phi(x,t)/h} \left[\sum_{j=0}^{m} h^{j-m} r_{j}(x,t)\right] dt\right| \leq C_{N} \int_{0}^{T} e^{-cd(x,x(t))^{2}/h} \left[\sum_{j=0}^{m} h^{j-m} d(x,x(t))^{N}\right] \chi(x,t) dt$$

where χ is the characteristic function of $\operatorname{supp}(a)$. Thus the $L^2(X)$ norm of the last term in (3.28) is $O(h^{\infty})$. Since the first two terms on the right of (3.28) are Gaussian wave packets at $\gamma(0)$ and $\gamma(T)$ with nonvanishing amplitudes, it follows (see [Zw12, Section 8.4.2]) that

$$WF_{scl}(Pu) = \gamma(0) \cup \gamma(T)$$

Step 7. Wave front set of u.

Let $(x_0,\xi_0) \in T^*X$. We will test whether (x_0,ξ_0) is in WF_{scl}(u) by integrating u against the wave packet $e^{i\Psi/h}b$, where $\Psi = \Psi(\cdot; x_0,\xi_0)$ and b are as in Proposition 2.3. Note in particular that

$$\Psi(x_0) = 0, \qquad d\Psi(x_0) = \xi_0, \qquad \nabla^2 \operatorname{Im}(\Psi)(x_0) > 0, \qquad b(x_0) \neq 0.$$

Moreover, since b is supported close to x_0 , we have

$$d\Psi \neq 0$$
 in supp $(b) \setminus \{x_0\}$, $\operatorname{Im}(\Psi)(x) \ge cd(x, x_0)^2$ in supp (b) .

We thus need to study the integral

$$I(h) := \int_X u \overline{e^{i\Psi/h}b} \, dV = \int_0^T \int_X e^{i\Theta(x,t)/h} r(x,t) \, dV \, dt$$

where

$$\Theta(x,t) := \Phi(x,t) - \overline{\Psi(x)}, \qquad r(x,t) := a(x,t)\overline{b(x)}.$$

We will show that $I(h) = O(h^{\infty})$ if $(x_0, \xi_0) \notin \gamma([0, T])$ (see cases 7a and 7b below), and $I(h) \sim h^{\frac{n+1}{2}}$ if $(x_0, \xi_0) \in \gamma([0, T])$ (see case 7c below). Then Proposition 2.3 proves that $WF_{scl}(u) = \gamma([0, T])$.

Case 7a. First assume that $x_0 \notin x([0,T])$. By the properties of Φ and Ψ and by the triangle inequality,

(3.29)
$$\operatorname{Im}(\Theta(x,t)) \ge c(d(x,x(t))^2 + d(x,x_0)^2) \ge \frac{c}{2}d(x(t),x_0)^2 \ge c' > 0$$

uniformly over supp(r). Thus in this case $I(h) = O(e^{-C/h})$.

Case 7b. Let $x_0 \in x([0,T])$, but $\xi_0 \neq \xi(t)$ for any $t \in I_0$ where $I_0 = \{t \in [0,T]; x(t) = x_0\}$. Note that

$$d_{x,t}\Theta(x(t),t) = (\xi(t) - d_x\Psi(x(t)), 0).$$

It follows that $d_{x,t}\Theta(x(t),t)$ is nonvanishing for any $t \in I_0$. Thus if we choose $\chi \in C_c^{\infty}([0,T])$ supported close to I_0 with $\chi = 1$ near I_0 , then nonstationary phase [Hö85, Theorem 7.7.1], together with its boundary version [Hö85, Theorem 7.7.17(i)] if $0 \in I_0$ or $T \in I_0$, implies that

$$\int_0^T \int_X e^{i\Theta(x,t)/h} r(x,t)\chi(t) \, dV \, dt = O(h^\infty).$$

If we replace χ by $1 - \chi$ in the integrand, we are in the region where $d(x(t), x_0) \ge c > 0$ and by (3.29) the resulting integral is $O(e^{-C/h})$.

Case 7c. Let now $(x_0,\xi_0) = (x(t_0),\xi(t_0))$ for some $t_0 \in [0,T]$. The phase function satisfies

$$\operatorname{Im}(\Theta) \ge 0, \qquad \Theta(x(t_0), t_0) = 0, \qquad d_{x,t} \Theta(x(t_0), t_0) = 0$$

In order to use stationary phase, we need to show that the Hessian $\nabla_{x,t}^2 \Theta|_{(x(t_0),t_0)}$ is invertible. Computing in Riemannian normal coordinates at $(x(t_0), t_0)$ and using (3.26)–(3.27), we have

(3.30)
$$\nabla_{x,t}^2 \Theta|_{(x(t_0),t_0)} = \begin{pmatrix} H+G & \dot{\xi} - H\dot{x} \\ (\dot{\xi} - H\dot{x})^t & (H\dot{x} - \dot{\xi}) \cdot \dot{x} \end{pmatrix}$$

where everything is evaluated at $t = t_0$ and $G := -\nabla^2 \bar{\Psi}(x_0)$ satisfies $\operatorname{Im}(G) > 0$. If $(\nabla_{x,t}^2 \Theta)\zeta = 0$ where $\zeta = \begin{pmatrix} v \\ z \end{pmatrix} \in \mathbb{C}^{n+1}$ with $v \in \mathbb{C}^n$ and $z \in \mathbb{C}$, looking at imaginary parts gives $0 = \operatorname{Im}(\nabla_x^2 \Phi)\zeta \cdot \bar{\zeta} = |\operatorname{Im}(G)^{1/2}v|^2 + |\operatorname{Im}(H)^{1/2}(v - \dot{x}z)|^2.$

Thus
$$v = 0$$
 and $\dot{x}z = 0$. If $\dot{x}(t_0) \neq 0$ this implies that $\zeta = 0$, showing that $\nabla^2_{x,t}\Theta|_{(x(t_0),t_0)}$ is
invertible. If $\dot{x}(t_0) = 0$ we additionally use that $0 = \nabla^2_{x,t}\Theta\begin{pmatrix}0\\z\end{pmatrix} = \begin{pmatrix}\dot{\xi}z\\0\end{pmatrix}$, so $\dot{\xi}(t_0)z = 0$. But

the condition $\dot{x}(t_0) = 0$ implies that $\dot{\xi}(t_0) \neq 0$ (otherwise dp would vanish at $\gamma(t_0)$ so $\gamma(t) \equiv \gamma(t_0)$, contradicting the assumption that γ is injective). Thus z = 0, showing that $\nabla_{x,t}^2 \Theta|_{(x(t_0),t_0)}$ is invertible also when $\dot{x}(t_0) = 0$.

Now if $\chi \in C_c^{\infty}([0,T])$ is supported near t_0 and satisfies $\chi = 1$ near t_0 , then stationary phase [Hö85, Theorem 7.7.5], or its boundary version [Hö85, Theorem 7.7.17(iii)] if $t_0 = 0$ or $t_0 = T$, implies that

$$\int_0^T \int_X e^{i\Theta(x,t)/h} r(x,t)\chi(t) \, dV \, dt = c_0 h^{\frac{n+1}{2}} + O(h^{\frac{n+1}{2}+1})$$

where $c_0 \neq 0$ since $r(x_0, t_0) \neq 0$. When we replace χ by $1 - \chi$ in the integrand, then we are either in the region where x(t) is away from x_0 (case 7a) or x(t) is close to x_0 but $\xi(t)$ is away from ξ_0 (case 7b), and the resulting integral is $O(h^{\infty})$. Combining these facts gives that $I(h) \sim h^{\frac{n+1}{2}}$.

Step 8. Semiclassical measure.

Let u be as above, and let v be the corresponding quasimode for \tilde{P}^* . Then u and v have the form

$$u(x) = \int_0^T e^{i\Phi(x,t)/h} a(x,t) \, dt, \qquad v(x) = \int_0^T e^{i\Phi(x,s)/h} \beta(x,s) \, ds$$

Note that u and v have the same phase function Φ since both P and \tilde{P} have principal symbol p_m . Moreover, β is an amplitude analogous to a satisfying

$$\beta(x(t),t) = \exp\left[-i\int_0^t b_{\tilde{P}^*}(x(s))\,ds\right] + O(h).$$

Using Lemma 3.3, we have

$$Qu(x) = \int_0^T e^{i\Phi(x,t)/h} \left[h^{-\ell} q_\ell(x, d_x \Phi(x,t)) a(x,t) + \sum_{j=1}^\ell h^{j-\ell} R_j a(x,t) \right] dt$$

where R_j is now defined in terms of Q. We only need to consider the $h^{-\ell}$ term, since the argument below will show that the other terms will be lower order as $h \to 0$. Then

$$\int_{X} Q u \bar{v} \, dV = h^{-\ell} \int_{0}^{T} \left[\int_{0}^{T} \int_{X} e^{i\Phi(x,t)/h} q_{\ell}(x, d_x \Phi(x,t)) a(x,t) \overline{e^{i\Phi(x,s)/h} \beta(x,s)} \, dV \, dt \right] ds$$

$$+ \text{lower order terms.}$$

For any fixed $s \in [0,T]$ we will study the integral inside the brackets, i.e. the integral

$$I_s(h) := \int_0^T \int_X e^{i\Theta_s(x,t)/h} r_s(x,t) \, dV \, dt$$

where

$$\Theta_s(x,t) := \Phi(x,t) - \overline{\Phi(x,s)}, \qquad r_s(x,t) = a(x,t)\overline{\beta(x,s)}q_\ell(x,d_x\Phi(x,t)).$$

Using the conditions for Φ , one has

$$\mathrm{Im}(\Theta_s) \geq 0 \text{ on } \mathrm{supp}(r_s), \qquad \Theta_s(x(s),s) = 0, \qquad d_{x,t}\Theta_s(x(s),s) = 0.$$

The formula (3.30) with the choices $\Psi(x) = \Phi(x,s)$ and $t_0 = s$ shows that

(3.31)
$$M(s) := \nabla_{x,t}^2 \Theta_s|_{(x(s),s)} = \begin{pmatrix} 2i \mathrm{Im}(H) & \dot{\xi} - H\dot{x} \\ (\dot{\xi} - H\dot{x})^t & (H\dot{x} - \dot{\xi}) \cdot \dot{x} \end{pmatrix}.$$

Again, this is the representation of $\nabla_{x,t}^2 \Theta_s|_{(x(s),s)}$ in Riemannian normal coordinates at (x(s), s). The argument after (3.30) shows that M(s) is invertible and hence we can use stationary phase to evaluate $I_s(h)$. However, since there is an additional integration over s, we need to ensure that all constants are uniform over $s \in [0, T]$. To do this, first we show that there is c > 0 with

(3.32)
$$|M(s)\zeta| \ge c|\zeta|,$$
 uniformly over $s \in [0,T].$

In fact, it is enough to prove that

(3.33)
$$\det(M(s)) = \alpha_0 \det(\operatorname{Im}(H(s)))$$

for some constant $\alpha_0 \neq 0$. The computation below will show that

$$\alpha_0 = \frac{i}{2} (2i)^n (|\dot{x}(0)|^2 + |\dot{\xi}(0)|^2).$$

Since Im(H(s)) is positive definite for $s \in [0, T]$, (3.32) is a consequence of (3.33) and the Cramer rule for computing $M(s)^{-1}$.

For proving (3.33), we write

$$\rho(s) := H(s)\dot{x}(s) - \dot{\xi}(s), \qquad \mathcal{R}(s) := \operatorname{Re}(H(s)), \qquad \mathcal{I}(s) := \operatorname{Im}(H(s)).$$

Here $\mathcal{R}(s)$ and $\mathcal{I}(s)$ are real symmetric matrices. The Schur complement formula for the determinant of a block matrix yields

$$\det(M(s)) = \det(2i\mathcal{I}(s))f(s)$$

where

$$f(s) := \rho(s) \cdot (\dot{x}(s) - (2i\mathcal{I}(s))^{-1}\rho(s)) = \frac{i}{2}\mathcal{I}(s)^{-1}\rho(s) \cdot \overline{\rho(s)}$$

In the last equality we used the fact that $\dot{x} = \text{Im}(\mathcal{I}^{-1}\rho)$. We thus have

(3.34)
$$\dot{f}(s) = \frac{i}{2} \left[-\mathcal{I}^{-1} \dot{\mathcal{I}} \mathcal{I}^{-1} \rho \cdot \bar{\rho} + \mathcal{I}^{-1} \dot{\rho} \cdot \bar{\rho} + \mathcal{I}^{-1} \rho \cdot \bar{\rho} \right]$$

To compute $\dot{\rho}$, note that from the Hamilton equations and (3.17) we obtain the formulas

$$\ddot{x} = B^t \dot{x} + C \dot{\xi}, \qquad \ddot{\xi} = -D \dot{x} - B \dot{\xi}$$

Using these formulas together with the matrix Riccati equation (3.19), we obtain

$$\dot{\rho} = -(HC + B)\rho.$$

Finally, taking the imaginary part of the matrix Riccati equation (3.19) we have

$$(3.36) \qquad \qquad \dot{\mathcal{I}} + \mathcal{I}C\mathcal{R} + \mathcal{R}C\mathcal{I} + B\mathcal{I} + \mathcal{I}B^t = 0$$

Inserting the formulas (3.35) and (3.36) into (3.34) proves that $\dot{f}(s) \equiv 0$. This shows (3.33), where $\alpha_0 = (2i)^n f(0)$ and $f(0) = \frac{i}{2}\rho(0) \cdot \overline{\rho(0)} = \frac{i}{2}(|\dot{x}(0)|^2 + |\dot{\xi}(0)|^2)$.

Having proved (3.32), the Taylor formula shows that there is $\delta > 0$ with

$$|d_{x,t}\Theta_s(\exp_{x(s)}(v), s+z)| \ge \frac{c}{2} \left| \begin{pmatrix} v \\ z \end{pmatrix} \right|, \qquad \left| \begin{pmatrix} v \\ z \end{pmatrix} \right| \le \delta, \quad s \in [0,T].$$

In normal coordinates at (x(s), s) this means that

$$\frac{|(x - x(s), t - s)|}{|\Theta'_s(x, t)|} \le C, \qquad |(x - x(s), t - s)| \le \delta, \quad s \in [0, T].$$

This is precisely the condition in the stationary phase theorem [Hö85, Theorem 7.7.5] that will make the constants uniform over $s \in [0, T]$. Note that since Q vanishes near the end points of γ , we do not need to worry about boundary contributions. Thus, if $\chi \in C_c^{\infty}(\mathbb{R})$ satisfies $\chi = 1$ for

 $|t| \leq \delta/2$ and $\operatorname{supp}(\chi) \subset (-\delta, \delta)$, and if the support of $\beta(x, s)$ is chosen sufficiently small depending on δ , then the stationary phase theorem implies that

$$\left| \int_0^T \int_X e^{i\Theta_s(x,t)/h} r_s(x,t) \chi(t-s) \, dV \, dt - h^{\frac{n+1}{2}} (\det(M(s)/2\pi i))^{-1/2} r_s(x(s),s) \right| \le Ch^{\frac{n+3}{2}}$$

uniformly over $s \in [0, T]$. There is no $|g|^{1/2}$ term coming from dV since the computation is done in Riemannian normal coordinates. Moreover, if we replace $\chi(t-s)$ by $1-\chi(t-s)$ in the integrand, then one is either in the region where x(t) is close to x(s) but $|t-s| \ge \delta/2$, so that the integral will be $O(h^{\infty})$ uniformly over $s \in [0, T]$ by nonstationary phase as in Case 7b above, or one is in the region where x(t) is away from x(s) and one can use Gaussian decay as in Case 7a above. It follows that

$$|I_h(s) - h^{\frac{n+1}{2}} (\det(M(s)/2\pi i))^{-1/2} r_s(x(s), s)| \le Ch^{\frac{n+3}{2}}$$

uniformly over $s \in [0, T]$. By (3.33) this may be rewritten as

$$I_h(s) = c_0 h^{\frac{n+1}{2}} \det(\mathcal{I}(s))^{-1/2} \exp\left[-i \int_0^s \left[b_P(x(\sigma)) - \overline{b_{\tilde{P}^*}(x(\sigma))}\right] d\sigma\right] q_\ell(\gamma(s)) + O(h^{\frac{n+3}{2}})$$

uniformly over $s \in [0, T]$, where

(3.37)
$$c_0 = (2\pi i)^{\frac{n+1}{2}} \alpha_0^{-\frac{1}{2}} = \frac{2\pi^{\frac{n+1}{2}}}{(|\dot{x}(0)|^2 + |\dot{\xi}(0)|^2)^{1/2}}$$

This proves that, as $h \to 0$,

$$h^{\ell-\frac{n+1}{2}} \int_X Q u \bar{v} \, dV \to c_0 \int_0^T \det(\mathcal{I}(s))^{-1/2} \exp\left[-i \int_0^s \left[b_P(x(\sigma)) - \overline{b_{\tilde{P}^*}(x(\sigma))}\right] \, d\sigma\right] q_\ell(\gamma(s)) \, ds$$

Using the properties of Φ and the notation (3.17), the function $b = b_P$ in Lemma 3.3 satisfies

$$b_P(x(t)) = \frac{1}{2i} \operatorname{tr}(C(t)H(t)) + p_{m-1}(\gamma(t)).$$

As $\tilde{p}_m = p_m$ is real, the adjoint \tilde{P}^* , defined with respect the $L^2(M)$ inner product associated to g, has in some local coordinates the full symbol

$$p_m + \frac{1}{i} \sum_{j=1}^n (\partial_{x_j \xi_j} p_m + |g|^{-1/2} \partial_{\xi_j} p_m \partial_{x_j} |g|^{1/2}) + \overline{\tilde{p}_{m-1}} + r_j$$

where r contains the terms of order $\leq m-2$. Thus, in normal coordinates at x(t),

$$b_{\tilde{P}^*}(x(t)) = \frac{1}{2i} \operatorname{tr}(C(t)H(t)) + \frac{1}{i} \operatorname{tr}(B(t)) + \overline{\tilde{p}_{m-1}(\gamma(t))}.$$

It follows that

$$b_P(x(t)) - \overline{b_{\tilde{P}^*}(x(t))} = \frac{1}{i} \operatorname{tr}(C(t)\mathcal{R}(t) + B(t)) + p_{m-1}(\gamma(t)) - \tilde{p}_{m-1}(\gamma(t)).$$

Moreover, using (3.36) and symmetries of trace,

 $\partial_t (\log \det \mathcal{I}) = \operatorname{tr}(\mathcal{I}^{-1}\dot{\mathcal{I}}) = -\operatorname{tr}(C\mathcal{R} + \mathcal{I}^{-1}\mathcal{R}C\mathcal{I} + \mathcal{I}^{-1}B\mathcal{I} + B^t) = -2\operatorname{tr}(C\mathcal{R} + B)$ which gives, using that $\mathcal{I}(0) = \operatorname{Id}$,

$$\det \mathcal{I}(t) = \exp\left[-2\int_0^t \operatorname{tr}(C(s)\mathcal{R}(s) + B(s))\,ds\right].$$

$$32$$

These facts imply that

$$\det(\mathcal{I}(t))^{-1/2} \exp\left[-i \int_0^t \left[b_P(x(s)) - \overline{b_{\tilde{P}^*}(x(s))}\right] ds\right] = \exp\left[-i \int_0^t (p_{m-1}(\gamma(s)) - \tilde{p}_{m-1}(\gamma(s))) ds\right].$$

This finally proves that

$$h^{\ell-\frac{n+1}{2}} \int_X Qu\bar{v} \, dV \to c_0 \int_0^T q_\ell(\gamma(t)) \exp\left[-i \int_0^t (p_{m-1}(\gamma(s)) - \tilde{p}_{m-1}(\gamma(s))) \, ds\right] \, dt$$

$$\downarrow \neq 0 \text{ is given in (3.37).} \qquad \Box$$

where $c_0 \neq 0$ is given in (3.37).

Remark 3.4. For later purposes, we observe that if M is a compact subset of X disjoint from the end points of γ , then for $k \ge 0$ one has

(3.38)
$$||u||_{H^k(M)} = O(h^{\frac{n+1}{4}-k}).$$

This follows from (3.1) upon taking $\tilde{P} = P^*$, u = v and $Q = R^* \chi^2 R$ where R is elliptic of order k (one can take k even) and $\chi \in C_c^{\infty}(X)$ satisfies $\chi = 1$ near M and $\chi = 0$ near the end points of γ .

Remark 3.5. It is possible to view the construction in Theorem 3.1 formally in terms of a quasimode construction in $X \times \mathbb{R}$ for the operator

$$Q(x, t, D_x, D_t) := h^{1-m} D_t + P(x, D_x).$$

If $\gamma(s) = (x(s), \xi(s))$ is an injective null bicharacteristic segment for P, then $(x(s), s, \xi(s), 0)$ is a similar bicharacteristic segment for Q and its spatial projection $\eta(s) = (x(s), s)$ satisfies $\dot{\eta}(s) \neq 0$ everywhere. One can then think of $e^{i\Phi(x,t)/h}a(x,t)$ as a guasimode for Q near the curve η , and integrating in t produces a quasimode for P.

We will also formulate a result related to computing the values of u at x(t) away from cusp points. This involves stationary phase in the t variable. Near cusp points the phase function will have a degenerate critical point in t and the behaviour is more complicated. For example, if P is the Tricomi operator, one can check from the construction that the phase function will have cubic behaviour near cusps. For such phase functions, at least if they are real valued, the asymptotics involve Airy functions [Hö85, Theorem 7.7.18].

Lemma 3.6. If u is the function in Theorem 3.1 and if $t_0 \in (0,T)$ is such that $\dot{x}(t_0) \neq 0$ and $x(t) \neq x(t_0)$ for $t \neq t_0$, then

$$u(x(t_0)) = h^{1/2} \frac{1}{c_{\gamma}(t_0)} \exp\left[-i \int_0^{t_0} b_P(x(s)) \, ds\right] + O(h^{3/2})$$

where c_{γ} is a smooth function depending on γ that is nonvanishing at any t_0 satisfying the assumption above.

Proof. We will give the proof under the more general assumption that $\dot{x}(t) \neq 0$ for any t with $x(t) = x(t_0)$, and $x(t_0)$ is not one of x(0) or x(T). Write $x_0 = x(t_0)$, and recall that

$$u(x_0) = \int_0^T e^{i\Phi(x_0,t)/h} a(x_0,t) \, dt$$

Let $I_{t_0} = \{t \in [0,T]; x(t) = x_0\}$. Since $\dot{x}(t) \neq 0$ when $t \in I_{t_0}$, points of I_{t_0} are isolated and hence $I_{t_0} = \{s_1, \ldots, s_N\}$ where $s_1 < s_2 < \ldots < s_N$ and one of the s_j is t_0 .

Note first that if t is away from I_{t_0} , then x(t) is bounded away from x_0 and thus by (3.24) the contribution of this region to $u(x_0)$ is $O(e^{-C/h})$. At points of I_{t_0} , since $x(s_j) = x_0$, the formulas (3.23)–(3.27) imply that

$$\Phi(x_0, s_j) = 0, \qquad \partial_t \Phi(x_0, s_j) = 0, \qquad \partial_t^2 \Phi(x_0, s_j) = (H\dot{x} - \dot{\xi}) \cdot \dot{x}|_{s_j}$$

where $H, \dot{x}, \dot{\xi}$ are written in terms of Riemannian normal coordinates at x_0 . Since $\text{Im}(H(s_j)) > 0$ and since $\dot{x}(s_j) \neq 0$ by assumption, we have $\partial_t^2 \Phi(x_0, s_j) \neq 0$ and hence we can use stationary phase [Hö85, Theorem 7.7.5] near each s_j . This implies that

$$u(x_0) = h^{1/2} \sum_{j=1}^{N} \frac{1}{c_{\gamma}(s_j)} a(x_0, s_j) + O(h^{3/2})$$

where

$$c_{\gamma}(s_j) = \left(\frac{2\pi i}{H(s_j)\dot{x}(s_j) \cdot \dot{x}(s_j)}\right)^{1/2}$$

and, since $x_0 = x(s_j)$, by (3.22) one has

$$a(x_0, s_j) = \exp\left[-i \int_0^{s_j} b_P(x(s)) \, ds\right].$$

Now if t_0 satisfies the assumption of the theorem, then N = 1 and the result follows.

4. Determining the scattering relation and ray transforms

We will next prove Theorem 1.1. The main idea for recovering the scattering relation is the following: if γ_1 is a maximal null bicharacteristic for P_1 in M starting at (x_0, ξ_0) , one constructs a solution u_1 of $P_1u_1 = 0$ in a neighborhood X of M so that $WF_{scl}(u_1)$ is on the extension $\check{\gamma}_1$ of γ_1 to X. Using the assumption $C_{P_1} = C_{P_2}$, there is a solution \tilde{u}_2 of $P_2\tilde{u}_2 = 0$ in M having the same Cauchy data as u_1 on ∂M . In fact one can extend \tilde{u}_2 so that $P_2\tilde{u}_2 = 0$ near M and $u_1 = \tilde{u}_2$ outside M (this is the mix-and-match construction mentioned in the introduction). One then considers the maximal null bicharacteristic γ_2 of P_2 in M going through (x_0, ξ_0) .

The above construction ensures that $WF_{scl}(u_1|_{X\setminus M})$ contains a part near (x_0, ξ_0) and also a part near $\alpha_{P_1}(x_0, \xi_0)$. Since $u_1 = \tilde{u}_2$ outside M, also $WF_{scl}(\tilde{u}_2|_{X\setminus M})$ contains these two parts. On the other hand, since $P_2\tilde{u}_2 = 0$ near M and since γ_2 contains (x_0, ξ_0) , by semiclassical propagation of singularities $WF_{scl}(\tilde{u}_2|_{X\setminus M})$ must also contain parts near the start and end points of γ_2 . Combining these conditions eventually leads to the fact that $\alpha_{P_1}(x_0, \xi_0) = \alpha_{P_2}(x_0, \xi_0)$. In the proof we need to deal with the possibility of tangential contacts with ∂M , which adds some technicalities.

If the principal parts of P_1 and P_2 agree, the recovery of subprincipal information in Theorem 1.1 is also based on constructing solutions concentrating near a maximal null bicharacteristic. However, instead of using the mix-and-match construction, it is more convenient to employ the integral identity in Lemma 2.8 and the formula for the semiclassical limit measure in (3.1).

Proof of Theorem 1.1. The proof will be done in several steps.

Step 1. Preparation.

We begin by extending both P_1 and P_2 smoothly to a slightly larger manifold X, in such a way that $P_1 = P_2$ in $X \setminus M$ (this is possible since $P_1 = P_2$ to infinite order at ∂M). The principal symbols $p_j = p_{j,m}$ of P_j then satisfy

(4.1)
$$p_1 = p_2 \text{ in } T^*(X \setminus M^{\text{int}}).$$

In particular, the null directions on $\partial(T^*M)$ are the same for P_1 and P_2 . Note also that if X is chosen sufficiently small, both P_1 and P_2 will be real principal type in X (i.e. the nontrapping condition will hold in X), see [Hö85, proof of Theorem 26.1.7].

Let $(x_0,\xi_0) \in \partial'_{\operatorname{null},P_1}(T^*M)$, and let $\gamma_1: [0,T_1] \to T^*M \setminus 0$ be the maximal null bicharacteristic for P_1 with $\gamma_1(0) = (x_0,\xi_0)$ (the case where $\gamma_1(T_1) = (x_0,\xi_0)$ is analogous). We denote by $\check{\gamma}_1$ the maximal continuation of γ_1 as a null bicharacteristic of P_1 in X. Since γ_1 is maximal, there is a strictly increasing sequence $(t_j^-)_{j=1}^\infty$ with $t_j^- \to 0$ and a strictly decreasing sequence $(t_j^+)_{j=1}^\infty$ with $t_j^+ \to T_1$ such that

$$\gamma_1(t_i^{\pm}) \in T^*(X \setminus M).$$

Since $d(x_1(t_j^{\pm}), M) \to 0$ as $j \to \infty$, where $x_1(t) = \pi(\gamma_1(t))$, after passing to subsequences we may choose a strictly decreasing sequence (κ_j) with

(4.2)
$$d(x_1(t_{j+1}^{\pm}), M) < \kappa_j < d(x_1(t_j^{\pm}), M), \qquad \kappa_j \to 0.$$

Then $X_j := \{x \in X ; d(x, M) < \kappa_j\}$ is an open manifold so that $M \subset X_j \subset \subset X$ and

(4.3)
$$\gamma_1(t_j^{\pm}) \in T^*(X \setminus \overline{X}_j).$$

Let $\gamma_2 : [-S_2, T_2] \to T^*M \setminus 0$ be the maximal null bicharacteristic for P_2 in M with $\gamma_2(0) = (x_0, \xi_0)$ and $S_2, T_2 \ge 0$.

Step 2. Strategy.

We wish to show that $S_2 = 0$ or $T_2 = 0$, i.e. $(x_0, \xi_0) \in \partial'_{\text{null}, P_2}(T^*M)$, and that

$$\{\gamma_1(0), \gamma_1(T_1)\} = \{\gamma_2(-S_2), \gamma_2(T_2)\}.$$

This will imply that $\partial'_{\operatorname{null},P_1}(T^*M) \subset \partial'_{\operatorname{null},P_2}(T^*M)$ and $\alpha_{P_1} = \alpha_{P_2}$ on $\partial'_{\operatorname{null},P_1}(T^*M)$. Changing the roles of P_1 and P_2 then implies that $\partial'_{\operatorname{null},P_1}(T^*M) = \partial'_{\operatorname{null},P_2}(T^*M)$ and $\alpha_{P_1} = \alpha_{P_2}$.

Below we will work with j fixed, and we will suppress j from the notation of u_1 etc.

Step 3. There is $u_1 \in H^m(X_j)$ with $P_1u_1 = 0$ in X_j and $WF_{scl}(u_1) = \check{\gamma}_1([t_j^-, t_j^+]) \cap T^*X_j$.

Let v_1 be the quasimode associated to $\check{\gamma}_1|_{[t_i^-, t_i^+]}$ in X constructed in Theorem 3.1, satisfying

$$WF_{scl}(v_1) = \breve{\gamma}_1([t_j^-, t_j^+]), \qquad WF_{scl}(P_1v_1) = \breve{\gamma}_1(t_j^-) \cup \breve{\gamma}_1(t_j^+).$$

Then $||P_1v_1||_{H^s(X_j)} = O(h^\infty)$ for any fixed s since $WF_{scl}(P_1v_1)$ is away from \overline{X}_j using (4.3). Let $u_1 = v_1|_{X_j} + r_1$, where $r_1 \in H^{s+m-1}(X_j)$ is the solution given in Proposition 2.2 of the equation

$$P_1 r_1 = -P_1 v_1 \text{ in } X_j$$

with

$$\|r_1\|_{H^{s+m-1}(X_j)} \lesssim \|P_1v_1\|_{H^s(X_j)} = O(h^{\infty}).$$

Then WF_{scl}(u₁) = WF_{scl}(v₁|_{X_j}) = $\check{\gamma}_1([t_j^-, t_j^+]) \cap T^*X_j$, and $P_1u_1 = 0$ in X_j .

Step 4. There is $\tilde{u}_2 \in H^m(X_j)$ so that $P_2\tilde{u}_2 = 0$ in X_j , $\tilde{u}_2 = u_1$ in $X_j \setminus M$, and \tilde{u}_2 is L^2 -tempered.

To prove this we invoke the assumption $C_{P_1} = C_{P_2}$, which implies that there is $w_2 \in H^m(M)$ such that

$$P_2w_2 = 0$$
 in M , $\nabla^k w_2|_{\partial M} = \nabla^k u_1|_{\partial M}$ for $k \le m - 1$.

Define the function

$$\tilde{u}_2 := \begin{cases} w_2, & \text{in } M, \\ u_1, & \text{in } X_j \setminus M \end{cases}$$

Then $\tilde{u}_2 \in H^m(X_j)$, and $P_2\tilde{u}_2 = 0$ in X_j since this holds separately in M and $X_j \setminus M$. It remains to show that \tilde{u}_2 is L^2 -tempered. First note that $u_1 - \tilde{u}_2$ is in $H^m_M(X_j)$ and solves

$$P_2(u_1 - \tilde{u}_2) = -P_2 u_1.$$

We may add a function in $N(P_2)$ to \tilde{u}_2 so that one still has $P_2\tilde{u}_2 = 0$ in X_j and $\tilde{u}_2 = u_1$ in $X_j \setminus M$, and additionally $u_1 - \tilde{u}_2 \in N(P_2)^{\perp}$. Then (2.2) implies that

(4.4)
$$\|u_1 - \tilde{u}_2\|_{L^2_M} \lesssim \|P_2(u_1 - \tilde{u}_2)\|_{L^2_M} = \|P_2u_1\|_{L^2_M} \lesssim \|u_1\|_{H^m(M)}.$$

Thus $\|\tilde{u}_2\|_{L^2_M} \lesssim \|u_1\|_{H^m(M)} = O(h^{\frac{n+1}{4}-m})$ by (3.38). It follows that \tilde{u}_2 is L^2 -tempered.

Step 5. $\gamma_1(\sigma_-) = \gamma_2(-S_2)$ and $\gamma_1(\sigma_+) = \gamma_2(T_2)$ for some $\sigma_{\pm} \in \{0, T_1\}$.

By Step 3 we have $WF_{scl}(u_1) = \check{\gamma}_1([t_j^-, t_j^+]) \cap T^*X_j$. Since $\tilde{u}_2 = u_1$ in $X_j \setminus M$, we further have (4.5) WF_1(\tilde{u}_2|_{X_j \setminus X_j}) = (\check{\gamma}_1([t_j^-, 0)) \sqcup \check{\gamma}_1((T_1, t_j^+))) \cap T^*(X_j \setminus M)

(4.5)
$$WF_{scl}(u_2|_{X_j \setminus M}) = (\gamma_1([t_j, 0]) \cup \gamma_1((T_1, t_j^+])) \cap T^*(X_j \setminus M).$$

Using (4.2) we have $\check{\gamma}_1(t_2^+) \in WF_{scl}(\tilde{u}_2|_{X \setminus M})$ for $k > i$, and since the wave front set

Using (4.2) we have $\check{\gamma}_1(t_k^{\pm}) \in \operatorname{WF}_{\operatorname{scl}}(\tilde{u}_2|_{X_j \setminus M})$ for k > j, and since the wave front set is closed this implies that $\gamma_1(0), \gamma_1(T_1) \in \operatorname{WF}_{\operatorname{scl}}(\tilde{u}_2)$. However, since $\gamma_1(0) = \gamma_2(0)$, semiclassical propagation of singularities for the solution \tilde{u}_2 of $P_2\tilde{u}_2 = 0$ in X_j , together with the maximality of γ_2 , implies that $\check{\gamma}_2(\tau_k^{j,\pm}) \in \operatorname{WF}_{\operatorname{scl}}(\tilde{u}_2|_{X_j \setminus M})$ for some increasing sequence $\tau_k^{j,-} \to -S_2$ and decreasing sequence $\tau_k^{j,+} \to T_2$ where $\check{\gamma}_2$ is the continuation of γ_2 to X. After passing to subsequences, we may assume that $-S_2 - 1/k < \tau_k^{j,-} < -S_2$ and $T_2 < \tau_k^{j,+} < T_2 + 1/k$.

Up to now we have been working with a fixed j. Next we consider the diagonal sequences $\tau_j^{j,\pm}$ as $j \to \infty$, and note that by (4.5) one has

(4.6)
$$\breve{\gamma}_2(\tau_j^{j,\pm}) = \breve{\gamma}_1(\sigma_j^{\pm})$$

for some σ_j^{\pm} with $\sigma_j^{\pm} \in [t_j^-, 0] \cup [T_1, t_j^+].$

It follows that $\tau_j^{j,-} \to -S_2$ and $\tau_j^{j,+} \to T_2$, as well as (possibly after passing to subsequences) $\sigma_j^{\pm} \to \sigma^{\pm}$ for some $\sigma_{\pm} \in \{0, T_1\}$. Thus by (4.6) one has

$$\gamma_1(\sigma_-) = \gamma_2(-S_2), \qquad \gamma_1(\sigma_+) = \gamma_2(T_2).$$

Step 7. $\partial'_{\operatorname{null},P_1}(T^*M) = \partial'_{\operatorname{null},P_2}(T^*M)$ and $\alpha_{P_1} = \alpha_{P_2}$.

Now if $\sigma_{-} \neq \sigma_{+}$, then necessarily $T_1 > 0$ and $\{\sigma_{-}, \sigma_{+}\} = \{0, T_1\}$. Thus

$$\{\gamma_1(0), \gamma_1(T_1)\} = \{\gamma_2(-S_2), \gamma_2(T_2)\}.$$

In particular, $(x_0, \xi_0) = \gamma_1(0) = \gamma_2(0)$ is one of the endpoints of γ_2 , so either $S_2 = 0$ or $T_2 = 0$.

On the other hand, assume that $\sigma_{-} = \sigma_{+} = 0$ (the case $\sigma_{\pm} = T_1$ is analogous). Then $\gamma_2(-S_2) = \gamma_2(T_2)$ and consequently $S_2 = T_2 = 0$ (since $\gamma_2|_{[-S_2,T_2]}$ is injective by the real principal type

assumption). We wish to prove that also $T_1 = 0$. If this is not true, then $T_1 > 0$ and $\gamma_1(0) \neq \gamma_1(T_1)$ using that $\gamma_1|_{[0,T_1]}$ is injective. Returning to (4.6), after passing to subsequences if necessary, it follows that $\sigma_j^{\pm} \leq 0$ for all j. Now in local coordinates in T^*M , the curves $\check{\gamma}_j(t)$ for small t are very close to the line $(x_0, \xi_0) + t\dot{\gamma}_1(0)$ where $\dot{\gamma}_1(0) = \dot{\gamma}_2(0) \neq 0$ by (4.1) and the real principal type assumption. Since $\tau_j^{j,-} < 0$, $\tau_j^{j,+} > 0$ and $\sigma_j^{\pm} \leq 0$, this leads to a contradiction with (4.6). This proves that $T_1 = 0$ and

$$\{\gamma_1(0), \gamma_1(T_1)\} = \{\gamma_2(-S_2), \gamma_2(T_2)\}\$$

also when $\sigma_+ = \sigma_-$.

We have now proved the claims stated in Step 2, and thus Step 7 is complete.

Step 8. If P_1 and P_2 have the same principal symbol and if μ is a half density on M, one has

$$\exp\left[i\int_0^T \sigma_{\rm sub}[P_1^{\mu}](\gamma(t))\,dt\right] = \exp\left[i\int_0^T \sigma_{\rm sub}[P_2^{\mu}](\gamma(t))\,dt\right]$$

whenever $\gamma : [0,T] \to T^*M \setminus 0$ is a maximal null bicharacteristic in M.

By the Remark after Theorem 1.1, it is enough to prove that

$$\exp\left[i\int_0^T (p_{m-1,1} - p_{m-1,2})(\gamma(t))\,dt\right] = 0.$$

Write $Q = P_1 - P_2$. Since $P_1 = P_2$ in $X \setminus M$, the operator Q is a differential operator with smooth coefficients in X that vanishes in $X \setminus M$ and has principal symbol

$$q_{m-1} = p_{m-1,1} - p_{m-1,2}$$

We denote by $\check{\gamma}$ the continuation of γ as a null bicharacteristic curve in X.

We now use Theorem 3.1 with $P = P_1$ and $\tilde{P} = P_2$ and construct quasimodes v_1, v_2 associated with $\check{\gamma}|_{[-\varepsilon_1, T+\varepsilon_2]}$ with ε_j chosen so that the end points are outside M. Then $\|P_1v_1\|_{H^s(M)} =$ $\|P_2^*v_2\|_{H^s(M)} = O(h^{\infty})$. As in Step 3, we construct solutions $u_j = v_j + r_j$ of $P_1u_1 = 0$ and $P_2^*u_2 = 0$ in M with $\|r_j\|_{H^{s+m-1}(M)} = O(h^{\infty})$.

Since $C_{P_1} = C_{P_2}$ and $Q = P_1 - P_2$ vanishes outside M, the integral identity in Lemma 2.8 implies that

$$0 = (Qu_1, u_2)_{L^2(M)} = (Qv_1, v_2)_{L^2(X)} + O(h^{\infty}).$$

Multiplying this identity by $h^{-\frac{n+1}{2}+m-1}$ and using (3.1), we obtain

$$\int_{0}^{T} q_{m-1}(\gamma(t)) \exp\left[-i \int_{0}^{t} q_{m-1}(\gamma(s)) \, ds\right] \, dt = 0.$$

The integrand is equal to $i\partial_t(\exp\left[-i\int_0^t q_{m-1}(\gamma(s))\,ds\right])$, hence it follows that

$$\exp\left[-i\int_0^T q_{m-1}(\gamma(t))\,dt\right] = 1$$

This is the required statement.

Remark 4.1. If the maximal null bicharacteristic $\gamma : [0,T] \to T^*M \setminus 0$ is sufficiently nice, e.g. $\dot{x}(t) \neq 0$ and x(t) is injective on [0,T], one can give an alternative proof of Step 8 by using the mixand-match construction and a similar argument as in Steps 3–5. That is, one constructs solutions of $P_j u_j = 0$ associated with γ , and defines another solution of $P_2 \tilde{u}_2 = 0$ in X_1 so that $\tilde{u}_2 = u_1$

outside M. By semiclassical propagation of singularities, $u_2 - \tilde{u}_2$ is $O(h^{\infty})$ at the end point x(T). Comparing the values of u_1 and u_2 at x(T) using Lemma 3.6 proves Step 8.

Proof of Theorem 1.2. The proof is the same as in Step 8 in the proof of Theorem 1.1. By Lemma 2.8, the assumption $C_{P+Q_1} = C_{P+Q_2}$ implies the integral identity

$$((Q_1 - Q_2)u_1, u_2)_{L^2(M)} = 0$$

whenever $u_j \in H^m(M)$ solve $(P + Q_1)u_1 = (P^* + Q_2^*)u_2 = 0$ in M. We wish to find such solutions u_j that are close to the quasimodes constructed in Theorem 3.1.

Let $\gamma : [0,T] \to T^*M \setminus 0$ be a maximal null bicharacteristic in M. We extend P as a smooth differential operator to some slightly larger open manifold X and let $\check{\gamma}$ be the continuation of γ to X. We also extend Q_j smoothly to X so that the coefficients of $Q_1 - Q_2$ vanish outside X. Choose some $\varepsilon_j > 0$ so that $\check{\gamma}(-\varepsilon_1)$ and $\check{\gamma}(T+\varepsilon_2)$ are in $X \setminus M$. Let $v_1 = v_{1,h} \in C_c^{\infty}(X)$ be the quasimode given by Theorem 3.1 related to $\check{\gamma}|_{[-\varepsilon_1,T+\varepsilon_2]}$ satisfying $\|(P+Q_1)v_1\|_{H^s(M)} = O(h^{\infty})$, and let $v_2 = v_{2,h}$ be the corresponding quasimode satisfying $\|(P^* + Q_2^*)v_2\|_{H^s(M)} = O(h^{\infty})$.

We define $u_j = v_j + r_j$, where r_j are correction terms satisfying

$$(P+Q_1)r_1 = f_1 \text{ in } M, \qquad (P^*+Q_2^*)r_2 = f_2 \text{ in } M,$$

where $f_1 = -(P + Q_1)v_1|_M$ and $f_2 = -(P^* + Q_2^*)v_2|_M$. Note that $f_j \in C^{\infty}(M)$ with $f_1 \in N(P_1^* + Q_1^*)^{\perp}$ and $f_2 \in N(P + Q_2)^{\perp}$. By Proposition 2.2, there exist solutions $r_j \in H^{s+m-1}(M)$ of the above equations with the norm estimates

$$||r_j||_{H^{s+m-1}(M)} \lesssim ||f_j||_{H^s(M)} = O(h^\infty).$$

Then $u_j \in H^{s+m-1}(M)$ also solve $(P+Q_1)u_1 = (P^*+Q_2^*)u_2 = 0$ in M.

Inserting the solutions u_i in the integral identity, we obtain that

$$\int_M (Q_1 - Q_2) v_1 \bar{v}_2 \, dV = O(h^\infty)$$

Using Theorem 3.1 and the fact that the coefficients of $Q_1 - Q_2$ vanish outside M, this implies that

$$\int_0^T \sigma_{\rm pr}[(Q_1 - Q_2)](\gamma(t)) \, dt = 0.$$

Here we also used that the subprincipal symbols of $P+Q_1$ and $P+Q_2$ agree, so that the exponential factor in (3.1) goes away.

We next consider the case of strictly hyperbolic operators and prove Theorem 1.3. First we verify that strictly hyperbolic operators are indeed real principal type (this should be well known, but we give a proof for completeness in Appendix A).

Lemma 4.2. Let X be an open manifold, let $\phi \in C^{\infty}(X, \mathbb{R})$, and let P be a strictly hyperbolic differential operator of order m in X with respect to the level surfaces of ϕ . Then P is of real principal type in X.

Proof of Theorem 1.3. We begin by establishing suitable analogues of Proposition 2.2 (see Step 1 below) and Lemma 2.8 (Step 2 below), and then indicate the minor modifications required in the proofs of Theorems 1.1 and 1.2 by giving a proof of (1.6) (Step 3 below). The other claims can be proven using analogous modifications.

Choose an open manifold $Y_1 \subset X$ so that $M_1 \subset Y_1$ and $\overline{Y_1}$ is compact. Define

$$X_1 = \{ x \in Y_1 ; 0 < \phi(x) < T \}.$$

Step 1. Given any $f \in H_0^s(X_1)$, with $s \ge 1$, there is a solution $u \in H^{s+m-1}(X_1)$ of $P_1 u = f$ satisfying $\operatorname{tr}_{\Gamma_-}^{m-1} u = 0$ and

$$\|u\|_{H^{s+m-1}(X_1)} \le C \|f\|_{H^s(X_1)}.$$

We define the space

$$\mathcal{H} = \{ u \in H^{s+m-1}(X_1) ; P_1 u \in H^s_0(X_1), \text{ tr}_{\Gamma_-}^{m-1} u = 0 \},\$$

equipped with the norm defined by

$$||u||_{\mathcal{H}}^2 = ||u||_{H^{s+m-1}(X_1)}^2 + ||P_1u||_{H^s(X_1)}^2.$$

Now \mathcal{H} is a Hilbert space and writing $P = P_1|_{\mathcal{H}}$, $P : \mathcal{H} \to H_0^s(X_1)$ is clearly continuous. Let us show that P is surjective. Let $f \in H_0^s(X_1)$. We may extend f by zero to obtain a function in $H^s(X)$, still denoted by f. By [Hö85, Theorem 23.2.4] there is $u \in H^{s+m-1}_{loc}(X)$ such that $\operatorname{supp}(u) \subset \phi^{-1}([0,\infty))$ and $P_1u = f$ on X_1 . Hence P is surjective.

It follows now (see [EHN96, Proposition 2.3 and 2.4]) that the pseudoinverse P^{\dagger} of P is continuous $P^{\dagger}: H_0^s(X_1) \to \mathcal{H}$ and $PP^{\dagger}f = f$. Thus $u = P^{\dagger}f$ solves Pu = f with the continuous dependence (4.7).

Step 2. If
$$C_{P_1}^{\text{lat}} = C_{P_2}^{\text{lat}}$$
, then
(4.8) $((P_1 - P_2)u_1, u_2)_{L^2(M)} = 0$

whenever $u_j \in H^m(M)$ solve $P_1 u_1 = P_2^* u_2 = 0$ in M, $\operatorname{tr}_{\Gamma_-}^{m-1} u_1 = 0$ and $\operatorname{tr}_{\Gamma_+}^{m-1} u_2 = 0$.

As $C_{P_1}^{\text{lat}} \subset C_{P_2}^{\text{lat}}$ we can find $\tilde{u}_2 \in H^m(M)$ such that $P_2 \tilde{u}_2 = 0$ in M and $\operatorname{tr}_{\Gamma \cup \Gamma_-}^{m-1} \tilde{u}_2 = \operatorname{tr}_{\Gamma \cup \Gamma_-}^{m-1} u_1$. Then, using also $\operatorname{tr}_{\Gamma_+}^{m-1} u_2 = 0$,

$$((P_1 - P_2)u_1, u_2) = -(P_2u_1, u_2) = -(P_2(u_1 - \tilde{u}_2), u_2) = -(u_1 - \tilde{u}_2, P_2^*u_2) = 0.$$

Step 3. If P_1 and P_2 have the same principal symbol and if μ is a half density on M, one has

$$\exp\left[i\int_0^T \sigma_{\rm sub}[P_1^{\mu}](\gamma(t))\,dt\right] = \exp\left[i\int_0^T \sigma_{\rm sub}[P_2^{\mu}](\gamma(t))\,dt\right]$$

whenever $\gamma : [0,T] \to T^*M \setminus 0$ is a maximal null bicharacteristic in M that does not meet $\partial(T^*M)$ away from Γ .

It is again enough to prove

(4.9)
$$\exp\left[i\int_0^T q_{m-1}(\gamma(t))\,dt\right] = 1,$$

where q_{m-1} is the principal symbol of $Q = P_1 - P_2$. We denote by $\check{\gamma}$ the continuation of γ as a null bicharacteristic curve in X.

We use Theorem 3.1 with $P = P_1$ and $\tilde{P} = P_2$ to obtain quasimodes v_1, v_2 associated with $\check{\gamma}|_{[-\varepsilon_1, T+\varepsilon_2]}$ with ε_j chosen so that the end points are outside X_1 . Observe that, when X_1 is chosen small enough, v_j can be constructed so that, writing $\Sigma = \phi^{-1}(0) \cup \phi^{-1}(T)$, one has $\operatorname{supp}(v_j) \cap \Sigma = \emptyset$. We choose $\chi \in C^{\infty}(X_1)$ so that $\chi = 1$ in M and that $\chi = 0$ near $\partial X_1 \setminus \Sigma$. Finally we choose $r_1 \in H^m(X_1)$ so that $P_1r_1 = -\chi P_1v_1$, $\operatorname{tr}_{\Gamma_-}^{m-1} r_1 = 0$ and that (4.7) holds with s = 1. Then

$$||r_1||_{H^m(X_1)} \le C ||\chi P_1 v_1||_{H^1(X_1)} = O(h^\infty),$$

and setting $u_1 = v_1 + r_1$, it follows that $P_1 u_1 = 0$ in M.

Note that P_2 is strictly hyperbolic also with respect to the level surfaces of $-\phi$. Repeating the argument in Step 1, with P_1 replaced by P_2 and ϕ by $-\phi$, we see that there is a solution $r_2 \in H^{s+m-1}(X_1)$ of $P_2r_2 = -\chi P_2v_2$ satisfying $\operatorname{tr}_{\Gamma_+}^{m-1}r_2 = 0$ and $||r_2||_{H^1(X_1)} = O(h^{\infty})$. We set $u_2 = v_2 + r_2$.

Since $C_{P_1} = C_{P_2}$ and $Q = P_1 - P_2$ vanishes outside M, the integral identity (4.8) implies that

$$0 = (Qu_1, u_2)_{L^2(M)} = (Qv_1, v_2)_{L^2(X)} + O(h^{\infty}).$$

Multiplying this identity by $h^{-\frac{n+1}{2}+m-1}$ and using (3.1), we obtain (4.9) as in the proof of Theorem 1.1.

We will show next that by small modifications of the above proofs, one can obtain partial data results in the corresponding inverse problems. Let $\Gamma \subset \partial M$ be a nonempty open set. We use the notation

$$\|\operatorname{tr}_{\Gamma}^{m-1} u\|_{\mathcal{H}^{m-1}(\Gamma)}^{2} = \sum_{j=0}^{m-1} \|\nabla^{j} u\|_{\Gamma}\|_{L^{2}(\Gamma)}^{2},$$

and consider the partial Cauchy data set

$$C_{P,\Gamma} = \{ (u|_{\Gamma}, \nabla u|_{\Gamma}, \dots, \nabla^{m-1}u|_{\Gamma}) ; u \in H^{m}(M) \text{ solves } Pu = 0 \text{ in } M \text{ and} \\ \| \operatorname{tr}_{\partial M \setminus \Gamma}^{m-1} u \|_{\mathcal{H}^{m-1}(\partial M \setminus \Gamma)} \leq \| \operatorname{tr}_{\Gamma}^{m-1} u \|_{\mathcal{H}^{m-1}(\Gamma)} \}.$$

The last condition means that we are only using solutions of Pu = 0 whose Cauchy data on $\partial M \setminus \Gamma$ is not much larger than the Cauchy data on Γ (the constant 1 in the inequality is quite arbitrary and could be replaced by any other fixed constant). For simplicity we do not consider the scattering relation in the following result, whose proof is given in Appendix A.

Theorem 4.3. Let M be a compact manifold with smooth boundary, let P_1, P_2 be real principal type differential operators of order $m \ge 1$ in M, and let $\Gamma \subset \partial M$ be a nonempty open set. Assume that $P_1 = P_2$ to infinite order on ∂M . If the principal symbols of P_1 and P_2 coincide, then for any nonvanishing half density μ on M one has

(4.10)
$$\exp\left[i\int_0^T \sigma_{\rm sub}[P_1^{\mu}](\gamma(t))\,dt\right] = \exp\left[i\int_0^T \sigma_{\rm sub}[P_2^{\mu}](\gamma(t))\,dt\right]$$

whenever $\gamma : [0,T] \to T^*M$ is a maximal null bicharacteristic curve for P_1 whose spatial projection intersects Γ transversally and does not meet ∂M away from Γ .

Finally, if $m \ge 2$ and if $P_j = P + Q_j$ where Q_j has order $\le m - 2$ for j = 1, 2, then then

(4.11)
$$\int_0^T \sigma_{\rm pr}[Q_1](\gamma(t)) \, dt = \int_0^T \sigma_{\rm pr}[Q_2](\gamma(t)) \, dt$$

whenever $\gamma : [0,T] \to T^*M$ is a maximal null bicharacteristic curve for P whose spatial projection intersects Γ transversally and does not meet ∂M away from Γ .

Finally, we show that for first order operators a converse of Theorem 1.1 holds: the scattering relation and integrals of the subprincipal symbol modulo $2\pi\mathbb{Z}$ determine the Cauchy data set, at least under a strict convexity assumption. The result is stated for C^{∞} Cauchy data sets since we use a regularity result from [PU05, PSU12] stated in the C^{∞} case. Note that any first order differential operator with real principal symbol in M is of the form $P = \frac{1}{i}L + V$, where L is a real vector field in M and $V \in C^{\infty}(M)$. Let X be an open manifold containing M, and extend Lsmoothly to X. We say that M is strictly convex for L if

$$\left. \frac{d^2}{dt^2} (\rho(x(t)) \right|_{t=0} < 0$$

whenever $\rho \in C^{\infty}(X)$ is a boundary defining function for M (i.e. $M = \{\rho \geq 0\}$ and $d\rho \neq 0$ on ∂M), and x(t) is an integral curve of L so that $x(0) \in \partial M$ and $\dot{x}(0) \in T(\partial M)$. A local coordinate computation shows that this condition does not depend on the extension or on the choice of ρ . The proof of the following theorem is in Appendix A.

Theorem 4.4. Let M be compact with smooth boundary, and let $P_j = \frac{1}{i}L_j + V_j$ be real principal type differential operators of order 1 in M. Assume that is M is strictly convex for L_1 and L_2 . Suppose that

$$\alpha_{P_1} = \alpha_{P_2}, \qquad \exp\left[-i\int_0^{T_1} V_1(x_1(t))\,dt\right] = \exp\left[-i\int_0^{T_2} V_2(x_2(t))\,dt\right]$$

whenever $x_j : [0, T_j] \to M$ are maximal integral curves of L_j with $x_1(0) = x_2(0)$. Then the C^{∞} Cauchy data sets of P_1 and P_2 are equal:

 $\{u|_{\partial M} ; u \in C^{\infty}(M), P_1u = 0 \text{ in } M\} = \{u|_{\partial M} ; u \in C^{\infty}(M), P_2u = 0 \text{ in } M\}.$

5. Boundary determination

In this section we assume that M is a compact manifold with smooth boundary and that P is a real principal type differential operator of order $m \ge 2$. We consider the problem of determining the boundary values of coefficients of P and their derivatives (possibly up to gauge) from the knowledge of C_P . For $(x_0, \xi_0) \in T^*(\partial M)$, we give two arguments for determining boundary values at x_0 (these arguments were already described in the introduction, recall also that ν is the *inner* unit conormal vector to ∂M with respect to the auxiliary Riemannian metric on M):

- (Elliptic region) If $t \mapsto p_m(x_0, \xi_0 + t\nu)$ has a simple non-real root, we use exponentially decaying solutions that concentrate near x_0 to give an analogue of boundary determination results for second order elliptic equations.
- (Hyperbolic region) If $t \mapsto p_m(x_0, \xi_0 + t\nu)$ has at least two distinct real roots, we use solutions concentrating near two null bicharacteristics through x_0 and obtain an analogue of boundary determination results for the wave equation.

In fact the regions could be mixed, and we will use a combination of both methods. Here are some examples:

- If $P = \partial_t^2 \Delta_{g_0}$ is the wave operator in $M := M_0 \times (0, T)$, then the elliptic region (resp. hyperbolic region) at $(x_0, t_0) \in \partial M_0 \times (0, T)$ is the set of spacelike covectors (resp. timelike covectors) in $T^*_{(x_0, t_0)}(\partial M)$.
- If P is elliptic, then all roots of $t \mapsto p_m(x_0, \xi_0 + t\nu)$ are non-real.
- If m is odd, the map $t \mapsto p_m(x_0, \xi_0 + t\nu)$ always has a real root unless x_0 is a characteristic boundary point.

- If P is strictly hyperbolic with respect to the level surfaces of $\phi \in C^{\infty}(X, \mathbb{R})$ (see Section 1), and if x_0 in an interior point of $\partial M \cap \{\phi = c\}$ for some c, then by definition $t \mapsto p_m(x_0, \xi_0 + t\nu)$ has m distinct real roots.
- If $P = (\partial_t^2 \Delta_{g_0})(\partial_t^2 + \Delta_{g_0})$ in $M_0 \times (0, T)$, there can be both real and non-real roots.

The special solutions in the hyperbolic region will be the ones constructed in Theorem 3.1. In the elliptic region, new special solutions will be required. Their construction is based on the fact that whenever $x \in \partial M$ and $t \mapsto p_m(x, \xi + t\nu)$ has a simple non-real root, there is an approximate solution of Pu = 0 in M with complex phase that concentrates near x, decays exponentially in the interior, and whose boundary value oscillates in direction ξ . The argument is similar to the construction of a boundary parametrix for the wave equation in the elliptic region, see e.g. [Ta81, Chapter IX].

Theorem 5.1. Let P be a differential operator of order $m \ge 2$ in M (here the principal symbol p_m is allowed to be complex valued). Let $x \in \partial M$, and assume that for some $\xi \in T_x^* \partial M$ the map $z \mapsto p_m(x, \xi + z\nu)$ has a simple root z with positive imaginary part. Fix s > 0. For $0 < h \le 1$ there exists $u = u_h \in C^{\infty}(M)$ having the form

$$u = e^{i\Phi/h}a$$

where $d\Phi|_x = \xi + z\nu$, a is supported near x, $a|_{\partial M}$ can be prescribed arbitrarily near x independently of h, $\operatorname{Im}(\Phi) \geq 0$ on $\operatorname{supp}(a)$, and

$$||u||_{L^2(M)} \sim h^{1/2}, \qquad ||u||_{H^k(M)} \lesssim h^{-k+1/2}, \qquad ||Pu||_{H^s(M)} = O(h^\infty).$$

Proof. Choose local coordinates near x so that x corresponds to 0, M is given by $\{x_n \ge 0\}$ near 0, the unit conormal ν corresponds to (0, 1), and the cotangent vector ξ corresponds to $(\xi', 0)$ with $\xi' \in \mathbb{R}^{n-1}$. Write $u = e^{i\Phi/h}a$. By Lemma 3.3, we have

$$Pu = e^{i\Phi/h} \left[h^{-m} p_m(x, d\Phi)a + h^{1-m} (\frac{1}{i}La + ba) + \sum_{j=2}^m h^{j-m} R_j a \right].$$

We first construct a smooth complex function Φ so that near 0,

(5.1)
$$\begin{cases} \Phi(x',0) = x' \cdot \xi', \\ \operatorname{Im}(\partial_n \Phi(x',0)) > 0, \\ p_m(x,d\Phi) = 0 \text{ to infinite order on } \{x_n = 0\}. \end{cases}$$

In fact, if we define $\Phi(x', 0) = x' \cdot \xi'$, then

$$p_m(x, d\Phi)|_{x_n=0} = p_m(x', 0, \xi', \partial_n \Phi(x', 0)).$$

This vanishes if $\partial_n \Phi(x', 0)$ is a root of $z \mapsto p_m(x', 0, \xi', z)$. By assumption there exists a simple root with positive imaginary part when x' = 0, hence also for x' near 0, and we denote this root by z(x'). Then z(x') depends smoothly on x' since it is a simple root, and we may define

(5.2)
$$\partial_n \Phi(x',0) = z(x').$$

Next we compute

$$\partial_{x_n}(p_m(x, d\Phi))|_{x_n=0} = \partial_{x_n} p_m(x', 0, \xi', z(x')) + \partial_{\xi_j} p_m(x', 0, \xi', z(x')) \partial_{j_n} \Phi(x', 0)$$

We wish to choose $\partial_n^2 \Phi(x', 0)$ so that this vanishes. However, writing

$$p_m(x', 0, \xi', \xi_n) = (\xi_n - z(x'))q(x', \xi', \xi_n)$$
42

where $q(x',\xi',z(x')) \neq 0$, we see that $\partial_{\xi_n} p_m(x',0,\xi',z(x')) \neq 0$, and hence we may define

$$\partial_n^2 \Phi(x',0) = -\frac{1}{\partial_{\xi_n} p_m} (\partial_{x_n} p_m + \sum_{\alpha=1}^{n-1} \partial_{\xi_\alpha} p_m \partial_{\alpha n} \Phi) \Big|_{(x',0,\xi',z(x'))}$$

With this choice we have $\partial_{x_n}(p_m(x, d\Phi))|_{x_n=0} = 0$. Continuing this process allows us to prescribe $\partial_n^3 \Phi(x', 0), \ \partial_n^4 \Phi(x', 0), \ \dots$ so that $\partial_{x_n}^2(p_m(x, d\Phi)), \ \partial_{x_n}^3(p_m(x, d\Phi)), \ \dots$ vanish on $\{x_n = 0\}$. Using Borel summation we obtain a smooth function Φ with the required Taylor series at $x_n = 0$, so that Φ satisfies (5.1).

We now construct the amplitude a by Borel summation in the form

$$a \sim \sum_{j=0}^{\infty} h^j a_j$$

where the a_k are smooth functions independent of h, and they satisfy the following transport equations to infinite order on $\{x_n = 0\}$:

$$\frac{1}{i}La_0 + ba_0 = 0,$$

$$\frac{1}{i}La_1 + ba_1 = -R_2a_0,$$

$$\frac{1}{i}La_2 + ba_2 = -R_3a_0 - R_2a_1$$

:

The first equation on $\{x_n = 0\}$ reads as

$$\frac{1}{i}\partial_{\xi_j}p_m(x',0,\xi',z(x'))\partial_j a_0(x',0) + b(x',0)a_0(x',0) = 0.$$

We define $a_0(x',0) = \eta(x')$ where η is any given function in $C_c^{\infty}(\mathbb{R}^{n-1})$ supported close enough to 0. We have seen that $\partial_{\xi_n} p_m(x',0,\xi',z(x')) \neq 0$, hence we may solve $\partial_n a(x',0)$ from the above equation as

$$\partial_n a_0(x',0) = -\frac{i}{\partial_{\xi_n} p_m} \left(\frac{1}{i} \sum_{\alpha=1}^{n-1} \partial_{\xi_\alpha} p_m \partial_\alpha a_0 + b a_0\right) \Big|_{(x',0,\xi',z(x'))}$$

Continuing in this way, we may define $\partial_n^j a_0(x', 0)$ for $j \ge 2$ and apply Borel summation to obtain a smooth function a_0 with the required Taylor series at $\{x_n = 0\}$, so that $\frac{1}{i}La_0 + ba_0 = 0$ to infinite order on $\{x_n = 0\}$. The functions a_j for $j \ge 1$ are constructed in a similar way (one may set $a_j(x', 0) = 0$ for $j \ge 1$), so that all the transport equations are satisfied to infinite order on $\{x_n = 0\}$. This completes the construction of the amplitude.

We have now constructed a smooth function

$$u = e^{i\Phi/h}a$$

so that

$$Pu \sim e^{i\Phi/h} \sum_{j=0}^{\infty} h^{j-m} f_j$$

where each f_j vanishes to infinite order on $\{x_n = 0\}$ and $|f_j| \leq 1$ uniformly over $h \leq 1$. Near 0 one has

$$cx_n \le \operatorname{Im}(\Phi(x', x_n)) \le Cx_n$$
43

for some c, C > 0. Thus if supp(a) is chosen to be sufficiently close to 0, one has

$$\|u\|_{L^{2}(M)}^{2} \sim \int_{\{|x'| \leq 1\}} \int_{0}^{1} e^{-x_{n}/h} dx_{n} dx' \sim h,$$

$$\|u\|_{H^{k}(M)}^{2} \lesssim h^{-2k} \int_{\{|x'| \leq 1\}} \int_{0}^{1} e^{-x_{n}/h} dx_{n} dx' \sim h^{1-2k}$$

and

$$||Pu||_{H^N(M)} \lesssim h^{-m-N} ||e^{-cx_n/h}O(x_n^{\infty})||_{L^2(\mathrm{supp}(a))} = O(h^{\infty}).$$

The result follows.

The next example gives a useful interpretation for a boundary determination result for the wave equation. This will motivate the corresponding proof for general real principal type operators.

Example 5.2. Let (M_0, g_0) be compact with smooth boundary, let $V \in C^{\infty}(M_0 \times (0, T))$, and consider the Dirichlet problem for the wave equation

$$(\partial_t^2 - \Delta_{g_0} + V)u = 0$$
 in $M_0 \times (0, T)$, $u|_{\partial M_0 \times (0, T)} = f$, $u|_{t=0} = \partial_t u|_{t=0} = 0$.

The Dirichlet-to-Neumann map is given by

$$\Lambda_V: f \mapsto \partial_\nu u|_{\partial M_0 \times (0,T)}.$$

The boundary determination result in [SY18] shows that for a small neighborhood $\Gamma \subset \partial M_0 \times (0,T)$ of a boundary point (x_0,t_0) , the localized map $\Lambda_V^{\Gamma} : C_c^{\infty}(\Gamma) \to C^{\infty}(\Gamma), f \mapsto \Lambda_V f|_{\Gamma}$ is a pseudodifferential operator and the Taylor series of V at (x_0,t_0) can be computed from the symbol of Λ_V^{Γ} .

Consider coordinates (x_1, x', t) and assume that Γ corresponds to $\{x_1 = 0\}$. The symbol of Λ_V^{Γ} can be computed by testing against highly oscillatory functions $f(x', t) = \chi(x', t)e^{i(x'\cdot\xi'+t)/h}$ where χ is a smooth cutoff and ξ' is in the hyperbolic region (i.e. there are two null directions at (x_0, t_0) whose projection to $T^*(\partial M_0 \times (0,T))$ is $(0,\xi',1)$). The argument in [SY18] gives roughly (when the cutoff χ is made to depend on h in a suitable way) that

$$\lim_{h \to 0} h^{\alpha}((\Lambda_V - \Lambda_0)f, f)_{L^2(\Gamma)} = cV(x_0, t_0)$$

where $c \neq 0$ and α is a suitable number depending on n.

On the other hand, one has the integral identity

$$((\Lambda_V - \Lambda_0)f, f)_{L^2(\Gamma)} = \int_{M_0} \int_0^T V u \bar{u}_0 \, dt \, dV$$

where u is the solution given above, and u_0 solves $(\partial_t^2 - \Delta_{g_0})u_0 = 0$ in $M_0 \times (0, T)$ with Dirichlet data f and vanishing Cauchy data on $\{t = T\}$. Since u has vanishing Cauchy data on $\{t = 0\}$ and highly oscillatory boundary data, it is related to a null bicharacteristic (possibly with reflections) that starts at (x_0, t_0) and moves forward in time. Similarly, since u_0 has vanishing Cauchy data on $\{t = T\}$, it is related to a null bicharacteristic starting at (x_0, t_0) and moving backward in time. Thus the product $u\bar{u}_0$ concentrates near the intersection of the projections of these two bicharacteristics, i.e. near the point (x_0, t_0) . Analyzing the integral over $M_0 \times (0, T)$ when $h \to 0$ then leads to a proof for recovering $V(x_0, t_0)$. Proof of Theorem 1.4. Let P_1, P_2 be real principal type differential operators of order $m \geq 2$. Assume that

$$C_{P_1} = C_{P_2}.$$

The integral identity in Lemma 2.8 implies that

(5.3)
$$((P_1 - P_2)u_1, u_2)_{L^2(M)} = 0$$

whenever $u_j \in H^m(M)$ are solutions of $P_1u_1 = 0$ and $P_2^*u_2 = 0$ in M. Depending on the roots of the characteristic polynomial, we will use different special solutions u_j to prove the theorem. The strategy however will be the same: the main point is to prove that $p_{m,2}(x, \xi + \cdot \nu)$ vanishes at the roots of $p_{m,1}(x, \xi + \cdot \nu)$ whenever $(x, \xi) \in T^*(\partial M)$ is close to (x_0, ξ_0) , i.e. that

(5.4)
$$\begin{cases} p_{m,2}(x,\xi+\tau\nu) = 0 \text{ for any } \tau \in \mathbb{C} \text{ for which } p_{m,1}(x,\xi+\tau\nu) = 0, \\ \text{whenever } (x,\xi) \in T^*(\partial M) \text{ is close to } (x_0,\xi_0). \end{cases}$$

Here and below, we will use the fact that the assumptions of the theorem remain true for any $(x,\xi) \in T^*(\partial M)$ that is close enough to (x_0,ξ_0) .

Let us show how (5.4) implies the claim (1.7) for j = 0 (the case $j \ge 1$ will be done later). By (5.4) the polynomial $\tau \mapsto p_{m,2}(x, \xi + \tau \nu)$ vanishes at each of the *m* distinct roots of $\tau \mapsto p_{m,1}(x, \xi + \tau \nu)$. The fact that *x* is a noncharacteristic boundary point for P_j implies that both maps are polynomials of degree exactly *m* in τ . Thus one must have, for some $c(x, \xi) \ne 0$, that

$$p_{m,2}(x,\xi+\tau\nu) = c(x,\xi)p_{m,1}(x,\xi+\tau\nu), \qquad \tau \in \mathbb{R}.$$

Multiplying by τ^{-m} and letting $\tau \to \infty$ gives that

$$c(x,\xi) = \frac{p_{m,2}(x,\nu_x)}{p_{m,1}(x,\nu_x)}.$$

Thus in fact $c(x,\xi) = c(x)$. For $(x,\xi) \in T^*(\partial M)$ close to (x_0,ξ_0) we obtain

$$p_{m,2}(x,\xi+\tau\nu_x) = c(x)p_{m,1}(x,\xi+\tau\nu_x), \qquad \tau \in \mathbb{R}.$$

For any $x \in \partial M$ near x_0 , the set $\{\xi + \tau \nu_x; \tau \in \mathbb{R} \text{ and } (x,\xi) \text{ is near } (x_0,\xi_0)\}$ is open in T_x^*M . By real-analyticity in the fiber variable one then has

$$p_{m,2}(x,\xi) = c(x)p_{m,1}(x,\xi), \qquad x \in \partial M \text{ near } x_0, \ \xi \in T_x^*M,$$

where c is a smooth nonvanishing function near x_0 on ∂M given by

$$c(x) = \frac{p_{m,2}(x,\nu_x)}{p_{m,1}(x,\nu_x)}.$$

This proves the desired claim (1.7) for j = 0.

The proof of (5.4) will be divided in several cases.

Case (1). We first assume condition (1) in the theorem, i.e. that $p_{m,2}(x,\xi + \sigma\nu) = 0$ for some non-real $\sigma \in \mathbb{C}$ (if this holds for $p_{m,1}$ instead we obtain the result by interchanging P_1 and P_2 ; note that the conclusion of the theorem remains invariant under this change). Since P_2 has real principal symbol, the non-real roots come in complex conjugate pairs and hence we may assume that $\operatorname{Im}(\sigma) > 0$.

Using Theorem 5.1 for P_2^* , there are smooth functions

$$v_2 = v_{2,h} = \frac{h^{-1/2} e^{i\Phi_2/h} a_2}{45}$$

where $d\Phi_2|_x = \xi + \sigma \nu$, $a_2|_{\partial M}$ is independent of h with $a_2(x) \neq 0$, $||v_2||_{L^2} \sim 1$, and for any fixed but suitably large s

$$||P_2 v_2||_{H^s(M)} = O(h^\infty).$$

By the solvability result for real principal type operators in Proposition 2.2(d), there exist smooth solutions $u_2 = u_{2,h} = v_2 + r_2$ of $P_2 u_2 = 0$ in M such that

$$||r_2||_{H^{s+m-1}(M)} = O(h^\infty).$$

Let $\{\tau_1, \ldots, \tau_m\}$ be the complex roots of $\tau \mapsto p_{m,1}(x, \xi + \tau\nu)$. By assumption these are all distinct. Let τ be one of these roots. We wish to prove that $p_{m,2}(x, \xi + \tau\nu) = 0$, which establishes (5.4). We will consider two subcases.

Subcase (1a). Suppose that τ is non-real. We may assume that $\text{Im}(\tau) > 0$ (since if $\text{Im}(\tau) < 0$, then $\bar{\tau}$ is also a root and the argument below shows that $p_{m,2}(x,\xi+\bar{\tau}\nu)=0$, and hence $p_{m,2}(x,\xi+\tau\nu)=0$ since the roots come in complex conjugate pairs).

Using Theorem 5.1 for P_1 , we obtain solutions $u_1 = u_{1,h}$ of $P_1u_1 = 0$ in M of the form $u_1 = v_1 + r_1$, and

$$v_1 = h^{-1/2} e^{i\Phi_1/h} a_1$$

where $d\Phi_1|_x = \xi + \tau \nu$, $a_1|_{\partial M}$ is independent of h with $a_1(x) \neq 0$, $||v_1||_{L^2} \sim 1$, and

$$||r_1||_{H^{s+m-1}(M)} = O(h^{\infty})$$

Inserting the solutions u_1 and u_2 into (5.3), we obtain that

$$\int_{M} (P_1 - P_2)(v_1) \bar{v}_2 \, dV = O(h^{\infty}).$$

The explicit form of v_1 and v_2 together with Lemma 3.3 gives that

(5.5)
$$0 = \lim_{h \to 0} h^m \int_M (P_1 - P_2)(v_1) \bar{v}_2 \, dV$$
$$= \lim_{h \to 0} h^{-1} \int_M (p_{m,1}(x, d\Phi_1) - p_{m,2}(x, d\Phi_1)) e^{i(\Phi_1 - \bar{\Phi}_2)/h} a_1 \bar{a}_2 \, dV.$$

Write $q(x) := p_{m,1}(x, d\Phi_1(x)) - p_{m,2}(x, d\Phi_1(x))$ and $\Theta := \Phi_1 - \overline{\Phi}_2$. By the construction of the functions Φ_j , see (5.1)–(5.2), in boundary normal coordinates (for g) near 0 one has

$$\Theta(x', x_n) = (\tau - \bar{\sigma})x_n + O(x_n^2)$$

Note that τ and σ depend on x', and $\text{Im}(\tau - \bar{\sigma}) > 0$. If the support of a_j is chosen small enough, one has as $h \to 0$

$$h^{-1} \int_{M} e^{i\Theta/h} qa_1 \bar{a}_2 \, dV$$

= $h^{-1} \int_{\{x_n > 0\}} e^{i(\tau - \bar{\sigma})x_n/h + O(x_n^2)/h} qa_1 \bar{a}_2 |g|^{1/2} \, dx$
= $\int_{\{x_n > 0\}} e^{i(\tau - \bar{\sigma})x_n + hO(x_n^2)} (qa_1 \bar{a}_2 |g|^{1/2})(x', hx_n) \, dx$
 $\rightarrow \int_{\{x_n > 0\}} e^{i(\tau - \bar{\sigma})x_n} (qa_1 \bar{a}_2 |g|^{1/2})(x', 0) \, dx$
= $\int_{\mathbb{R}^{n-1}} \frac{i}{\tau - \bar{\sigma}} (qa_1 \bar{a}_2 |g|^{1/2})(x', 0) \, dx'.$

In the last step we used that $\int_0^\infty e^{iwx_n} dx_n = \frac{i}{w}$ when Im(w) > 0. Since $a_j(x', 0)$ can be chosen arbitrarily near 0, we obtain that q(x', 0) = 0 near 0. This means that

$$p_{m,1}(x',0,\xi',\tau(x')) = p_{m,2}(x',0,\xi',\tau(x')).$$

But $\tau(x')$ was a root of $t \mapsto p_{m,1}(x', 0, \xi', t)$, so it is also a root for $p_{m,2}$, i.e. $p_{m,2}(x, \xi + \tau \nu) = 0$. This proves (5.4) whenever τ is non-real.

Subcase (1b). Now suppose that τ is real. Let $\gamma : [0,T] \to T^*M$ be the maximal null P_1 bicharacteristic curve with $\gamma(0) = (x_0, \xi_0 + \tau \nu_{x_0})$. Extend M to a slightly larger open manifold X, extend P_1 smoothly as a real principal type operator to X, and let $\check{\gamma}(t) = (x(t), \xi(t))$ be the extension of γ . Choose $t_- < 0$ and $t_+ > T$ so that $\check{\gamma}(t_{\pm})$ are outside M, and let v_1 be the quasimode in Theorem 3.1 associated with $\check{\gamma}|_{[t_-,t_+]}$ with $\|P_1v_1\|_{H^s(M)} = O(h^{\infty})$ for some fixed large s > 0. Use Proposition 2.2(d) to find r_1 with $\|r_1\|_{H^{s+m-1}(M)} = O(h^{\infty})$ so that $u_1 = v_1 + r_1$ solves $P_1u_1 = 0$ in M. Recall that v_1 has the form

$$v_1(x) = \int_{t_-}^{t_+} e^{i\Phi_1(x,t)/h} a_1(x,t) \, dt$$

Recall also from (3.23)–(3.25) that

(5.6) $\Phi_1(x(t),t) = 0, \qquad d_x \Phi_1(x(t),t) = \xi(t), \qquad \partial_t \Phi_1(x(t),t) = 0,$

and

(5.7) $\operatorname{Im}(\Phi_1(x,t)) \ge cd(x,x(t))^2, \qquad (x,t) \in \operatorname{supp}(a_1).$

As discussed in (3.23)–(3.27), the quantity $\mathcal{I} = \text{Im}((\partial_{x_j x_k} \Phi_1))|_{(x_0,0)}$ is invariantly defined and we may write

(5.8)
$$\nabla_{x,t}^2 \operatorname{Im}(\Phi_1)|_{(x_0,0)} = \begin{pmatrix} \mathcal{I} & -\mathcal{I}\dot{x} \\ -(\mathcal{I}\dot{x})^t & \mathcal{I}\dot{x}\cdot\dot{x} \end{pmatrix}$$

Inserting the solutions u_1 and u_2 in (5.3), we obtain that

$$\int_{M} ((P_1 - P_2)v_1)\bar{v}_2 \, dV = O(h^{\infty}).$$

Inserting the expressions for v_1 and v_2 and using Lemma 3.3, it follows that

$$\int_{M} \int_{t-}^{t+} (p_{m,1}(x, d_x \Phi_1(x, t)) - p_{m,2}(x, d_x \Phi_1(x, t))) e^{i(\Phi_1(x, t) - \bar{\Phi}_2(x))/h} a_1(x, t) \bar{a}_2(x) dt dV + \text{lower order terms} = O(h^{\infty}).$$

Let x' be normal coordinates at x_0 on ∂M (for the metric induced by g), and let (x', x_n) be corresponding boundary normal coordinates so that x_0 corresponds to 0. Since a_2 is supported near x_0 , the *M*-integral can be written in these coordinates and we have

(5.9)
$$\int_{\{x_n>0\}} \int_{t_-}^{t_+} e^{i\Theta(x,t)/h} q(x,t) \, dt \, dx + \text{lower order terms} = O(h^\infty)$$

where $q(x,t) = (p_{m,1}(x, d_x \Phi_1(x,t)) - p_{m,2}(x, d_x \Phi_1(x,t)))a_1(x,t)\bar{a}_2(x)|g(x)|^{1/2}$ and the phase Θ is given by $\Theta(x,t) = \Phi_1(x,t) - \bar{\Phi}_2(x)$.

We wish to use stationary phase to show that the main contribution to the integral (5.9) comes from the region near t = 0. Note first that

$$d_x \Theta(0,0) = \xi(0) - \xi_0 - \bar{\sigma}\nu = (\tau - \bar{\sigma})\nu, \qquad \partial_t \Theta(0,0) = 0.$$
47

We next study the Hessian of $\text{Im}(\Theta)$ in the (x', x_n, t) coordinates. By (5.8) we have

$$\nabla_{x,t}^2 \mathrm{Im}(\Theta)|_{(0,0)} = \begin{pmatrix} \mathcal{I} & -\mathcal{I}\dot{x} \\ -(\mathcal{I}\dot{x})^t & \mathcal{I}\dot{x}\cdot\dot{x} \end{pmatrix} + \nabla_{x,t}^2 \mathrm{Im}(\Phi_2)|_{(0,0)}$$

By (5.1) we have $\nabla_{x',t}^2 \text{Im}(\Phi_2)|_{(0,0)} = 0$. If $\zeta' = (v',z)^t$ with $v' \in \mathbb{C}^{n-1}$, and if $\zeta = (v',0,z)^t$, it follows that

(5.10)
$$\nabla_{x',t}^{2} \operatorname{Im}(\Theta)|_{(0,0)} \zeta' \cdot \bar{\zeta}' = \begin{pmatrix} \mathcal{I} & -\mathcal{I}\dot{x} \\ -(\mathcal{I}\dot{x})^{t} & \mathcal{I}\dot{x} \cdot \dot{x} \end{pmatrix} \zeta \cdot \bar{\zeta} = |\mathcal{I}^{1/2}(\begin{pmatrix} v' \\ 0 \end{pmatrix} - \dot{x}z)|^{2}.$$

If the right hand side vanishes, then $\dot{x}_n z = 0$ and $v' = \dot{x}' z$. But the fact that τ is a simple root of $t \mapsto p_{m,1}(x_0, \xi_0 + t\nu)$ implies that

(5.11)
$$\dot{x}_n(0) = \partial_{\xi_n} p_{m,1}(x_0, \xi_0 + \tau \nu) \neq 0.$$

Thus we get z = 0 and v' = 0. This proves that $\nabla^2_{x',t} \Theta|_{(0,0)}$ is invertible.

If $\chi(t)$ is a smooth cutoff supported near 0 with $\chi = 1$ near 0, the above discussion together with stationary phase on manifolds with boundary [Hö85, Theorem 7.7.17(ii)] implies that

$$\int_{\{x_n>0\}} \int_{t_-}^{t_+} e^{i\Theta(x,t)/h} q(x,t)\chi(t) \, dt \, dx = c_0 h^{\frac{n+2}{2}} q(0,0) + O(h^{\frac{n+4}{2}}), \qquad c_0 \neq 0$$

If we replace $\chi(t)$ by $1 - \chi(t)$ in the integrand, then the fact that the *x*-integral is over a small neighborhood of x_0 and the assumption that γ never returns to x_0 after t = 0 implies that the corresponding integral is over the region where x(t) is bounded away from x_0 , and hence the corresponding integral is $O(e^{-C/h})$. Multiplying (5.9) by $h^{-\frac{n+2}{2}}$ and letting $h \to 0$ implies that q(0,0) = 0, i.e.

$$p_{m,1}(x, d_x \Phi_1(x, t)) - p_{m,2}(x, d_x \Phi_1(x, t))|_{(0,0)} = p_{m,1}(x_0, \xi_0 + \tau\nu) - p_{m,2}(x_0, \xi_0 + \tau\nu) = 0.$$

Since $p_{m,1}(x_0,\xi_0+\tau\nu)=0$, we obtain $p_{m,2}(x_0,\xi_0+\tau\nu)=0$. This also holds for $(x,\xi)\in T^*(\partial M)$ close to (x_0,ξ_0) , which proves (5.4) for real τ .

Case (2). We now assume condition (2) in the theorem, i.e. that all roots for $\tau \mapsto p_{m,j}(x_0, \xi_0 + \tau \nu)$ are real and simple and have the stated properties. Let σ be such that $p_{m,2}(x_0, \xi_0 + \sigma \nu) = 0$, and let τ be any root of $t \mapsto p_{m,1}(x_0, \xi_0 + t\nu) = 0$. We want to prove that $p_{m,2}(x_0, \xi_0 + \tau \nu) = 0$, which would imply (5.4).

We again use the integral identity (5.3) with suitable special solutions u_j . Let $\gamma_j : [0, T_j] \to T^*M$ be the maximal null P_j -bicharacteristic curves with $\gamma_1(0) = (x_0, \xi_0 + \tau\nu)$ and $\gamma_2(0) = (x_0, \xi_0 + \sigma\nu)$, and let $\check{\gamma}_j(t) = (x_j(t), \xi_j(t))$ be extensions to a slightly larger manifold X so that the end points of $\check{\gamma}_j|_{[t_j^-, t_j^+]}$ are outside M. As in Subcase (1b) above, let $u_j = v_j + r_j$ solve $P_1u_1 = 0$ and $P_2^*u_2 = 0$ in M, where v_j is the quasimode provided by Theorem 3.1 associated with $\check{\gamma}_j|_{[t_j^-, t_j^+]}$ and $||r_j||_{H^{s+m-1}(M)} = O(h^\infty)$ for some fixed large s. Then v_j has the form

$$v_j(x) = \int_{t_j^-}^{t_j^+} e^{i\Phi_j(x,t)/h} a_j(x,t) dt$$

where the phase functions Φ_j satisfy the counterparts of (5.6)–(5.8) with x(t), $\xi(t)$, and \mathcal{I} replaced by $x_j(t)$, $\xi_j(t)$, and \mathcal{I}_j . Moreover, a_j is supported in a small neighborhood of the curve $(x_j(t), t)$. Inserting the solutions u_i in (5.3), we obtain

(5.12)
$$\int_{M} \int_{t_{1}^{-}}^{t_{1}^{+}} \int_{t_{2}^{-}}^{t_{2}^{+}} e^{i\Theta(x,t,s)/h} q(x,t,s) \, ds \, dt \, dV(x) + \text{lower order terms} = O(h^{\infty})$$

where $\Theta(x,t,s) = \Phi_1(x,t) - \overline{\Phi_2(x,s)}$ and

$$q(x,t,s) = (p_{m,1}(x,d_x\Phi_1(x,t)) - p_{m,2}(x,d_x\Phi_1(x,t)))a_1(x,t)a_2(x,s).$$

The integral in (5.12) is over a fixed small neighborhood of the set

$$E := \{ (\bar{x}, t, s) \, ; \, \bar{x} \in M, \, x_1(t) = x_2(s) = \bar{x} \}.$$

Again, we want to use stationary phase to show that the main contribution comes from t = s = 0. First note that if $x_1(t) = x_2(s) = \bar{x}$, then

$$\Theta(\bar{x},t,s) = 0, \qquad d_x \Theta(\bar{x},t,s) = \xi_1(t) - \xi_2(s), \qquad \partial_{t,s} \Theta(\bar{x},t,s) = 0.$$

By the assumption that the bicharacteristics intersect nicely, one has $E = \{(x_0, 0, 0)\} \cup K$ where K is a compact subset of $M^{\text{int}} \times [\varepsilon, T_1] \times [\varepsilon, T_2]$ for some $\varepsilon > 0$, and in K one always has $\xi_1(t) \neq \xi_2(s)$. This shows that when (t, s) is away from (0, 0), and if the supports of a_j were chosen small enough, the integral in (5.12) is $O(h^{\infty})$ by nonstationary phase.

It remains to evaluate the integral

$$\int_M \int_{\mathbb{R}^2} e^{i\Theta(x,t,s)/h} q(x,t,s) \chi(t,s) \, ds \, dt \, dV(x)$$

where $\chi(t,s)$ is a cutoff function with small support and with $\chi = 1$ near (0,0). Note that the *M*-integral is over a small neighborhood of x_0 . Now

$$\Theta(x_0, 0, 0) = 0, \qquad d_x \Theta(x_0, 0, 0) = \xi_1(0) - \xi_2(0) = (\tau - \sigma)\nu, \qquad \partial_{t,s} \Theta(x_0, 0, 0) = 0.$$

Let x' be normal coordinates on ∂M near x_0 , and let (x', x_n) be corresponding boundary normal coordinates so that x_0 corresponds to 0. Let $v' \in \mathbb{C}^{n-1}$ and write $\zeta' = (v', z, w)^t$ and $\zeta = (v', 0, z, w)^t$. Arguing as in (5.10), we compute

$$\nabla_{x',t,s}^{2} \operatorname{Im}(\Theta)|_{(0,0,0)} \zeta' \cdot \bar{\zeta}' = \nabla_{x,t,s}^{2} \operatorname{Im}(\Theta)|_{(0,0,0)} \zeta \cdot \bar{\zeta}$$
$$= |\mathcal{I}_{1}^{1/2} \begin{pmatrix} v' \\ 0 \end{pmatrix} - \dot{x}_{1} z)|^{2} + |\mathcal{I}_{2}^{1/2} \begin{pmatrix} v' \\ 0 \end{pmatrix} - \dot{x}_{2} w)|^{2} + |\mathcal{I}_{2}^{1/2} \langle v' \rangle + |\mathcal{I}_{2}^{1/2} \langle v \rangle + |\mathcal{I}_{2}^$$

Assume that the last quantity vanishes. Using that τ and σ are simple roots, we have $\dot{x}_{1,n}(0) \neq 0$ and $\dot{x}_{2,n}(0) \neq 0$ as in (5.11). Since \mathcal{I}_1 and \mathcal{I}_2 are positive definite, we obtain z = w = 0 and v' = 0. This proves that $\nabla^2_{x',t,s}\Theta|_{(0,0,0)}$ is invertible. Thus stationary phase [Hö85, Theorem 7.7.17(ii)] yields that

$$\int_{M} \int_{\mathbb{R}^{2}} e^{i\Theta(x,t,s)/h} q(x,t,s)\chi(t,s) \, ds \, dt \, dV(x) = c_0 h^{\frac{n+3}{2}} q(x_0,0,0) + O(h^{\frac{n+5}{2}}), \qquad c_0 \neq 0.$$

Multiplying (5.12) by $h^{-\frac{n+3}{2}}$ and letting $h \to 0$ we obtain that $q(x_0, 0, 0) = 0$, i.e.

$$p_{m,1}(x_0,\xi_0+\tau\nu) = p_{m,2}(x_0,\xi_0+\tau\nu).$$

Thus $p_{m,2}(x_0,\xi_0+\tau\nu)=0$. This remains true for $(x,\xi)\in T^*(\partial M)$ close to (x_0,ξ_0) , which proves (5.4).

Concluding the proof. We have now proved (5.4) in all cases. As discussed after (5.4), this gives the claim (1.7) for j = 0. To prove this for all j, we assume that the claim has already been proved

for $j \leq k-1$. Extend the function c as a smooth nonvanishing function to M, and replace P_1 by $c^{-1}P_1$ (so $p_{m,1}$ is replaced by $c^{-1}p_{m,1}$). By Lemma 5.4 we still have $C_{P_1} = C_{P_2}$, and

(5.13)
$$\partial_{x_n}^j(p_{m,1}-p_{m,2})(x',0,\eta) = 0 \text{ when } j \le k-1, x' \text{ is close to } 0, \eta \in \mathbb{R}^n.$$

Thus by the Taylor formula, for x near 0 one has

(5.14)
$$p_{m,1}(x,\eta) - p_{m,2}(x,\eta) = x_n^k f(x,\eta), \qquad f(x',0,\eta) = \frac{1}{k!} \partial_{x_n}^k (p_{m,1} - p_{m,2})(x',0,\eta).$$

Let us assume that we are in Subcase (1a), with $p_{m,1}(x,\xi+\tau\nu) = 0$ and $\text{Im}(\tau) > 0$ where $\tau = \tau(x')$. Multiplying (5.3) by h^{m-k} , we have

$$0 = \lim_{h \to 0} h^{m-k} \int_M (P_1 - P_2)(v_1) \bar{v}_2 \, dV$$

=
$$\lim_{h \to 0} h^{-k-1} \int_M (p_{m,1}(x, d\Phi_1) - p_{m,2}(x, d\Phi_1)) e^{i(\Phi_1 - \bar{\Phi}_2)/h} a_1 \bar{a}_2 \, dV.$$

Note that $p_{m,1}(x, d\Phi_1(x)) - p_{m,2}(x, d\Phi_1(x)) = x_n^k r(x)$ where $r(x) := f(x, d\Phi_1(x))$. Arguing as in Subcase (1a), we get

$$h^{-k-1} \int_{M} e^{i\Theta/h} x_{n}^{k} r a_{1} \bar{a}_{2} dV$$

$$= h^{-k-1} \int_{\{x_{n}>0\}} e^{i(\tau-\bar{\sigma})x_{n}/h+O(x_{n}^{2})/h} x_{n}^{k} r a_{1} \bar{a}_{2} |g|^{1/2} dx$$

$$= \int_{\{x_{n}>0\}} x_{n}^{k} e^{i(\tau-\bar{\sigma})x_{n}+hO(x_{n}^{2})} (r a_{1} \bar{a}_{2} |g|^{1/2}) (x', hx_{n}) dx$$

$$\rightarrow \int_{\{x_{n}>0\}} x_{n}^{k} e^{i(\tau-\bar{\sigma})x_{n}} (r a_{1} \bar{a}_{2} |g|^{1/2}) (x', 0) dx'$$

$$= \int_{\mathbb{R}^{n-1}} \frac{\Gamma(k+1)}{(-i(\tau-\bar{\sigma}))^{k+1}} (r a_{1} \bar{a}_{2} |g|^{1/2}) (x', 0) dx'$$

using that $\int_0^\infty t^k e^{iwt} dt = \frac{\Gamma(k+1)}{(-iw)^{k+1}}$ when $\operatorname{Im}(w) > 0$. Since $a_j(x',0)$ can be chosen arbitrarily near 0, we obtain that r(x',0) = 0 near 0, i.e. that for $(x,\xi) \in T^*(\partial M)$ close to (x_0,ξ_0)

(5.15)
$$\partial_{x_n}^k(p_{m,1} - p_{m,2})(x,\xi + \tau\nu) = 0$$

This is true for each of the *m* distinct zeros τ of $t \mapsto p_{m,1}(x', 0, \xi', t)$, hence by real-analyticity in the fiber variable it follows that

$$\partial_{x_n}^k(p_{m,1} - p_{m,2})(x', 0, \eta) = \tilde{c}(x')p_{m,1}(x', 0, \eta)$$

where $\tilde{c}(x')$ is smooth near 0' and has the invariant expression

$$\tilde{c}(y) = \frac{\partial_{x_n}^k(p_{m,1} - p_{m,2})(y,\nu_y)}{p_{m,1}(y,\nu_y)}$$

We now replace P_1 by $(1 - \chi(x)\tilde{c}(x')x_n^k/(k!))P_1$ where χ is a cutoff to a small neighborhood of 0 with $\chi = 1$ near 0. Then C_{P_1} and condition (5.13) remain unchanged, and we have

$$\partial_{x_n}^k (p_{m,1} - p_{m,2})(x', 0, \eta) = 0.$$

The argument above shows that if (1.7) holds for $j \leq k - 1$, then possibly after replacing c by some c_k , (1.7) holds for $j \leq k$. The Taylor coefficients of c_k at $x_n = 0$ up to order k are uniquely determined by (1.7). Thus we may use Borel summation to construct a smooth nonvanishing function $c \in C^{\infty}(M)$ with this Taylor series at $x_n = 0$ near 0, and with this choice of c the condition (1.7) holds for all j. This concludes the proof if we are in Subcase (1a).

The arguments for Subcase (1b) and Case 2 are quite analogous, and we will only give a sketch for Subcase (1b). Again we arrange so that (5.14) holds. Then (5.9) is replaced by

$$\int_{\{x_n>0\}} \int_{t_-}^{t_+} e^{i\Theta(x,t)/h} x_n^k r(x,t) \, dt \, dx + \text{lower order terms} = O(h^\infty)$$

where $r(x,t) = f(x, d_x \Phi_1(x,t)) a_1(x,t) \bar{a}_2(x) |g(x)|^{1/2}$. As discussed in Subcase (1b), the main contribution comes from the integral

$$\int_{\{x_n>0\}} \int_{\mathbb{R}} e^{i\Theta(x,t)/h} x_n^k r(x,t)\chi(t) \, dt \, dx$$

where $\chi(t)$ is a cutoff to the region near t = 0. We write

$$e^{i\Theta/h} = \left(\frac{h}{i\partial_{x_n}\Theta}\partial_{x_n}\right)^k (e^{i\Theta/h}).$$

Integrating by parts and noting that the boundary terms always vanish due to the x_n^k factor, we see that the largest contribution with respect to h comes when all the ∂_{x_n} derivatives hit the x_n^k factor. This term has the form

$$(ih)^k(k!)\int_{\{x_n>0\}}\int_{\mathbb{R}}e^{i\Theta(x,t)/h}(\partial_{x_n}\Theta)^{-k}r(x,t)\chi(t)\,dt\,dx.$$

The stationary phase argument in Subcase (1b) gives that r(0,0) = 0, i.e. that $f(x_0, d\Phi_1(x_0,0)) = 0$, which means that

$$\partial_{x_n}^k (p_{m,1} - p_{m,2})(x_0, \xi_0 + \tau \nu) = 0.$$

This is the counterpart of (5.15). The argument after (5.15) can now be repeated to conclude the proof. \Box

Remark 5.3. The proof shows that even when the principal parts of P_j have complex coefficients (and if the equations $P_j u = f$ are solvable in a suitable sense), it is possible to obtain boundary determination results when $t \mapsto p_m(x, \xi + t\nu)$ has sufficiently many simple non-real roots whose imaginary parts have suitable signs.

Finally, we give the proof of Theorem 1.5 on boundary determination for lower order terms. The proof is very similar to that of Theorem 1.4, so we will only sketch the required modifications.

Proof of Theorem 1.5. Let $P_j := P + Q_j$ and $Q := Q_1 - Q_2$, and let $q(x, \xi)$ be the principal symbol of Q. We need to show that

$$\partial_{x_n}^j q(x_0,\eta) = 0, \qquad j \ge 0, \ \eta \in \mathbb{R}^n.$$

The integral identity (5.3) becomes

$$(Qu_1, u_2)_{L^2(M)} = 0$$

for solutions of $P_1u_1 = 0$ and $P_2^*u_2 = 0$ in M.

Now $p_{m,1} = p_{m,2} = p_m$. By assumption $t \mapsto p_m(x_0, \xi_0 + t\nu)$ has s distinct simple roots $\{\tau_1, \ldots, \tau_s\}$, and these roots have nonnegative imaginary parts. We want to prove that

(5.16)
$$q(x_0, \xi_0 + \tau_j \nu) = 0, \qquad 1 \le j \le s.$$

Assume first that some root $\sigma \in \{\tau_1, \ldots, \tau_s\}$ is non-real, with $\text{Im}(\sigma) > 0$. This is the counterpart of Case (1). Let τ be one of the roots τ_1, \ldots, τ_s . If τ is non-real, then $\text{Im}(\tau) > 0$, and we argue as in Subcase (1a) above. The counterpart of (5.5) is

$$0 = \lim_{h \to 0} h^{-1} \int_M q(x, d\Phi_1(x)) e^{i(\Phi_1 - \bar{\Phi}_2)/h} a_1 \bar{a}_2 \, dV.$$

Evaluating the limit as in Subcase (1a), we obtain that

$$q(x_0,\xi_0+\tau\nu)=0$$

Thus $t \mapsto q(x_0, \xi_0 + t\nu)$ vanishes at each non-real root τ in the set $\{\tau_1, \ldots, \tau_s\}$. However, if τ is a real root, the argument in Subcase (1b) implies that $q(x_0, \xi_0 + \tau\nu) = 0$ also in this case. This proves (5.16) if one of the roots is non-real.

Next assume that all roots in $\{\tau_1, \ldots, \tau_s\}$ are real, and $s \ge \max(r+1, 2)$. Let σ be one of these roots, and let $\tau \ne \sigma$ be another one of these roots. The argument in Case (2) implies that

$$q(x_0, \xi_0 + \tau \nu) = 0.$$

Now $q(x_0, \xi_0 + \tau \nu) = 0$ at each root $\tau \neq \sigma$, and choosing σ to be a different root (this is where we need $s \geq 2$) implies (5.16).

Finally, if $q(x,\xi)$ is real valued and if some $\tau = \tau_j$ is such that $\text{Im}(\tau) < 0$, then the argument above shows that $q(x_0,\xi_0 + \bar{\tau}\nu) = 0$. Taking complex conjugates gives $q(x_0,\xi_0 + \tau\nu) = 0$. Thus if $Q_1 - Q_2$ has real principal symbol, then (5.16) holds without the assumption that the roots τ_j have nonnegative imaginary parts.

Now, the polynomial $t \mapsto q(x_0, \xi_0 + t\nu)$ has degree r, and (5.16) implies that it has at least $s \geq r+1$ distinct roots. Thus

$$u(x_0,\xi_0+t\nu)=0, \qquad t\in\mathbb{R}.$$

The same result holds when (x_0, ξ_0) is varied slightly, and real-analyticity in η implies that

$$q(x_0,\eta) = 0, \qquad \eta \in T^*_{x_0}M.$$

Arguing as in the end of proof of Theorem 1.4 gives that

$$\partial_{x_n}^j q(x_0, \eta) = 0, \qquad j \ge 0, \ \eta \in T_{x_0}^* M$$

This finishes the proof.

6. Semilinear equations

In this section we prove Theorem 1.7 related to semilinear equations of the form

$$Pu + a(x, u) = 0$$
 in M

where a(x, z) is a nonlinearity satisfying the following conditions. Fix an integer $s > \max(m, n/2)$, and assume initially that

(6.1) a(x,z) is analytic in z near 0 as a $H^{s}(M)$ -valued function,

(6.2)
$$a(x,0) = \partial_z a(x,0) = 0.$$

The proof is based on constructing solutions $u_{\varepsilon_1,\ldots,\varepsilon_r}$ to the semilinear equation that are close to $\varepsilon_1 v_1 + \ldots + \varepsilon_r v_r$, where v_j are suitable solutions of the linearized equation $Pv_j = 0$ concentrating near null bicharacteristics, and on higher order linearization with respect to the parameters ε_j . The following result will allow us to construct such solutions under the weak uniqueness assumption $N(P^*) = \{0\}.$

Lemma 6.1. Let M be a compact manifold with smooth boundary, and let P be a real principal type differential operator on M with $N(P^*) = \{0\}$. Assume that a(x, z) satisfies (6.1)–(6.2). There exist $\delta, C > 0$ such that for any v in the set

$$X_{\delta} = \{ v \in H^{s}(M) \, ; \, \|v\|_{H^{s}(M)} < \delta \},\$$

there is a solution $u = S(v) \in H^s(M)$ of

$$Pu + a(x, u) = Pv$$
 in M

which satisfies

(6.3)
$$\|u - v\|_{H^s(M)} \le C \|v\|_{H^s(M)}^2.$$

The map $S: X_{\delta} \to H^{s}(M), v \mapsto u$ is C^{∞} . In particular, if we define

$$U_{\delta} = \{ v \in X_{\delta} ; Pv = 0 \},\$$

then for any $v \in U_{\delta}$ the function u = S(v) solves Pu + a(x, u) = 0 in M.

Proof. Since $N(P^*) = \{0\}$, Proposition 2.2 shows that there is a bounded linear map $E : H^s(M) \to H^{s+m-1}(M)$ with PE = Id. Given $v \in X_{\delta}$ we wish to find a solution u = v+r of Pu+a(x,u) = Pv. It is enough to find r solving the fixed point equation

(6.4)
$$r = T_v(r)$$

where $T_v(r) = -E(a(x, v+r)).$

We now study (6.4) for $v \in X_{\delta}$, and show that T_v is a contraction on X_{δ} for δ small enough. All constants below will be uniform over $v \in X_{\delta}$. Note first that by (6.1), for some $\delta_0, R > 0$ one has

(6.5)
$$\sum_{j=0}^{\infty} \frac{\|\partial_z^j a(x,0)\|_{H^s(M)}}{j!} \delta_0^j \le R$$

Then (6.2) and the fact that $H^s(M)$ for s > n/2 is an algebra imply that

(6.6)
$$\|a(x,w)\|_{H^{s}(M)} \leq \sum_{j=2}^{\infty} \|\frac{\partial_{u}^{j}a(x,0)}{j!}w^{j}\|_{H^{s}(M)} \leq C_{s} \sum_{j=2}^{\infty} \frac{\|\partial_{u}^{j}a(x,0)\|_{H^{s}(M)}}{j!}C_{s}^{j}\|w\|_{H^{s}(M)}^{j}.$$

If δ is chosen small enough, (6.5) yields

(6.7)
$$\|a(x,w)\|_{H^s(M)} \le C \|w\|_{H^s(M)}^2, \qquad w \in X_{\delta},$$

and similarly

(6.8)
$$\|\partial_u a(x,w)\|_{H^s(M)} \le C \|w\|_{H^s(M)}, \qquad w \in X_{\delta}$$

Now

(6.9) $||T_v(w)||_{H^s} \le ||E(a(x,v+w))||_{H^{s+m-1}} \le C||a(x,v+w)||_{H^s} \le C||v+w||^2_{H^s}, \quad w \in X_{\delta},$ showing that T maps X_{δ} to itself when δ is small. Similarly,

$$\begin{aligned} \|T_v(u) - T_v(w)\|_{H^s} &\leq \|E(a(x, v+u) - a(x, v+w))\|_{H^{s+m-1}} \leq C \|a(x, v+u) - a(x, v+w)\|_{H^s} \\ &\leq C \left(\int_0^1 \|\partial_u a(x, v+(1-t)u + tw)\|_{H^s} \, dt \right) \|u - w\|_{H^s}. \end{aligned}$$

By (6.8), if δ is small enough one has

(6.10)
$$\|T_v(u) - T_v(w)\|_{H^s} \le \frac{1}{2} \|u - w\|_{H^s}, \qquad u, w \in X_{\delta}.$$

We have proved that there is $\delta > 0$ so that for any $v \in X_{\delta}$, the map $T_v : X_{\delta} \to X_{\delta}$ satisfies (6.9)–(6.10). By the Banach fixed point theorem, there is a unique $r = r_v \in X_{\delta}$ solving $r = T_v(r)$. Writing u = S(v) = v + r we obtain a solution of Pu + a(x, u) = Pv. The estimate (6.9) yields

$$||r||_{H^s} = ||T_v(r)||_{H^s} \le C ||v+r||_{H^s}^2 \le C(||v||_{H^s}^2 + ||r||_{H^s}^2),$$

and since $r \in X_{\delta}$ it follows that for δ small enough

$$||u - v||_{H^s} = ||r||_{H^s} \le C ||v||_{H^s}^2$$

To show that S is C^{∞} , consider the map

$$F: X_{\delta} \times X_{\delta} \to X_{\delta}, \quad F(v,r) = r - T_v(r) = r + E(a(x,v+r)).$$

Now F is C^{∞} by (6.1). It satisfies F(0,0) = 0 and

$$D_r F_{(0,0)}(h) = h + E(\partial_u a(x,0)h) = h.$$

Thus $D_r F_{(0,0)}$ is a homeomorphism, and the implicit function theorem [RR04, Theorem 10.6] shows that there is a smooth map G near 0 in X_{δ} so that F(v, r) = 0 near 0 iff r = G(v). But F(v, r) = 0iff $r = r_v$, which shows that $G(v) = r_v$ and that $v \mapsto r_v$ is C^{∞} . It follows that also S is C^{∞} . \Box

The next technical lemma shows that if u is a small solution to Pu + a(x, u) = 0 depending smoothly on a solution v of Pv = 0, and if the Cauchy data sets for nonlinearities a and \tilde{a} coincide, then the corresponding small solution of $P\tilde{u} + \tilde{a}(x, \tilde{u}) = 0$ with $u - \tilde{u} \in H_0^m(M)$ also depends smoothly on v. The argument requires the uniqueness assumption $N(P) = \{0\}$.

Lemma 6.2. Let M be a compact manifold with smooth boundary, and let P be a real principal type differential operator on M with $N(P) = N(P^*) = \{0\}$. Suppose that a, \tilde{a} satisfy (6.1)–(6.2) and they agree to high order on ∂M in the sense that

(6.11)
$$a(\cdot, z) - \tilde{a}(\cdot, z) \in H_0^s(M) \text{ for } z \text{ near } 0.$$

Assume that for any sufficiently small $\delta > 0$ there is $\delta_1 < \delta$ so that $C_{a,\delta_1} \subset C_{\tilde{a},\delta}$.

If δ is small enough, given any $v \in U_{\delta}$ and solution u = S(v) of Pu + a(x, u) = 0 there is a small solution $\tilde{u} \in H^{s}(M)$ of $P\tilde{u} + \tilde{a}(x, \tilde{u}) = 0$ such that $u - \tilde{u} \in H^{m}_{0}(M)$, the map $v \mapsto \tilde{u}$ is C^{∞} from U_{δ} to $H^{s}(M)$, and

(6.12)
$$\|\tilde{u} - v\|_{H^s(M)} \le C \|v\|_{H^s(M)}^2$$

Proof. Fix any small $\delta > 0$. By assumption there is $\delta_1 < \delta$ with $C_{a,\delta_1} \subset C_{\tilde{a},\delta}$. Moreover, by (6.3) there is $\delta_2 < \delta_1$ so that $v \in U_{\delta_2}$ implies that $u = S(v) \in X_{\delta_1}$. Thus for any $v \in U_{\delta_2}$ there is $\tilde{u} \in X_{\delta}$ solving $P\tilde{u} + \tilde{a}(x, \tilde{u}) = 0$ such that $u - \tilde{u} \in H_0^m(M)$. We need to show that the map $\tilde{S}: U_{\delta_2} \to H^s(M), v \mapsto \tilde{u}$ is smooth. In order to do this, write $w = w_v := u - \tilde{u} \in H^s(M) \cap H_0^m(M)$, and note that w solves

$$Pw = f, \qquad f := \tilde{a}(x, u - w) - a(x, u).$$

We wish to express w as the solution of a fixed point equation as in Lemma 6.1, and show that the map $v \mapsto w_v$ is smooth.

We first prove that in fact $w \in H_0^{s+m-1}(M)$ and $Pw = f \in H_0^s(M)$. Let M be contained in some larger manifold X, let \breve{u} be some $H_{\text{comp}}^s(X)$ extension of u, and let \breve{w} be the zero extension of w outside M. Then $\breve{w} \in H_M^m(X)$ solves

$$P\breve{w} = \tilde{a}(x,\breve{u} - \breve{w}) - a(x,\breve{u})$$
 a.e. in X

since the right hand side vanishes outside M by the assumption (6.11). By the argument leading to (6.6), the right hand side is in fact in $H_M^m(X)$ since $H_{\text{comp}}^m(X) \cap L^\infty(X)$ is an algebra, see [KP88]

for the latter fact. Regularity for real principal type operators (see Proposition 2.2(a)) then implies that $\breve{w} \in H_M^{m+(m-1)}(X)$. Bootstrapping this regularity argument gives $\breve{w} \in H_M^{s+m-1}(X)$, which yields $w \in H_0^{s+m-1}(M)$ and $f \in H_0^s(M)$ as required.

Next we define the spaces

$$Y := \{ \varphi \in H_0^{s+m-1}(M) \, ; \, P\varphi \in H_0^s(M) \}$$

and

 $Z := \{f; f = P\varphi \text{ for some } \varphi \in Y\}.$

Now Y with norm $\|\varphi\|_Y = \|\varphi\|_{H^{s+m-1}} + \|P\varphi\|_{H^s}$ is a Hilbert space. Moreover, Z is a closed subspace of $H^s(M)$ (and of $H^s_0(M)$): if $P\varphi_j \to f$ in $H^s(M)$ where $\varphi_j \in Y$, then the condition $N(P) = \{0\}$ and the estimate (2.2) imply that

$$\|\varphi_j - \varphi_k\|_{H^{s+m-1}} \lesssim \|P\varphi_j - P\varphi_k\|_{H^s}.$$

Thus (φ_j) is a Cauchy sequence in $H_0^{s+m-1}(M)$ and converges to some φ in this space. Since $P\varphi_j \to f$ in $H_0^s(M)$, we have $f = P\varphi \in H_0^s(M)$ showing that Z is closed in $H^s(M)$.

The map $P: Y \to Z, \varphi \to P\varphi$ is linear, bounded and bijective by the assumption $N(P) = \{0\}$. The open mapping theorem ensures that there is a bounded inverse $G: Z \to Y$. Let Q be the orthonormal projection to Z in $H^s(M)$. We may now rewrite the equation

$$Pw = \tilde{a}(x, u - w) - a(x, u), \qquad w \in Y,$$

equivalently as

(6.13)
$$w = GQ(\tilde{a}(x, u - w) - a(x, u)).$$

Consider the map

$$F: X_{\delta} \times X_{\delta} \to H^{s}(M), \quad F(v, w) = w - GQ(\tilde{a}(x, S(v) - w) - a(x, S(v))).$$

Then F is C^{∞} and $D_w F_{(0,0)}h = h$, hence by the implicit function theorem there is a smooth map H near 0 in X_{δ} so that F(v, w) = 0 near 0 iff w = H(v). But $F(v, w_v) = 0$, showing that the map $v \mapsto w_v$ is smooth. Moreover, writing u = v + r where r satisfies

$$||r||_{H^s} \lesssim ||v||_{H^s}^2$$

the equation (6.13) together with (6.2) implies that

$$\|w\|_{H^s} \lesssim \|v+r-w\|_{H^s}^2 + \|v+r\|_{H^s}^2 \lesssim \|w\|_{H^s}^2 + \|v\|_{H^s}^2.$$

Since $||w||_{H^s}$ is small, this gives $||w||_{H^s} \lesssim ||v||_{H^s}^2$, and (6.12) follows.

Proof of Theorem 1.7. We begin by fixing solutions $v_1, v_2, v_3 \in H^s(M)$ of $Pv_j = 0$ (we will later choose v_j to concentrate near certain null bicharacteristics). Let $\varepsilon = (\varepsilon_1, \varepsilon_2, \varepsilon_3)$, and define $u_{\varepsilon} = S(\varepsilon_1 v_1 + \varepsilon_2 v_2 + \varepsilon_3 v_3)$. Then u_{ε} depends smoothly on each ε_j , and by (6.3) one has

$$u_0 = 0, \qquad \partial_{\varepsilon_j} u_{\varepsilon}|_{\varepsilon = 0} = v_j.$$

Differentiating the equation $Pu_{\varepsilon} + a(x, u_{\varepsilon}) = 0$ and using the conditions $\partial_u^k a(x, 0) = 0$ for k = 0, 1, 2, we see that $v_{123} := \partial_{\varepsilon_1 \varepsilon_2 \varepsilon_3} u_{\varepsilon}|_{\varepsilon=0}$ satisfies

(6.14)
$$Pv_{123} + \partial_u^3 a(x,0)v_1v_2v_3 = 0.$$

We now let $\tilde{u}_{\varepsilon} = \tilde{S}(\varepsilon_1 v_1 + \varepsilon_2 v_2 + \varepsilon_3 v_3)$ be the function in Lemma 6.2. Then \tilde{u}_{ε} depends smoothly on the ε_j , and using (6.12) we have

$$\tilde{u}_0 = 0, \qquad \partial_{\varepsilon_j} \tilde{u}_{\varepsilon}|_{\varepsilon = 0} = v_j$$

Similarly, differentiating the equation $P\tilde{u}_{\varepsilon} + \tilde{a}(x,\tilde{u}_{\varepsilon}) = 0$ shows that $\tilde{v}_{123} := \partial_{\varepsilon_1 \varepsilon_2 \varepsilon_3} \tilde{u}_{\varepsilon}|_{\varepsilon=0}$ satisfies

(6.15)
$$P\tilde{v}_{123} + \partial_u^3 \tilde{a}(x,0)v_1v_2v_3 = 0$$

Since $u_{\varepsilon} - \tilde{u}_{\varepsilon} \in H_0^m(M)$, also $v_{123} - \tilde{v}_{123} \in H_0^m(M)$. Subtracting the equations (6.14) and (6.15) and integrating against v_4 where $\bar{P}^*v_4 = 0$ (so that $P^*\bar{v}_4 = 0$) yields that

(6.16)
$$\int_{M} (\partial_{u}^{3} a(x,0) - \partial_{u}^{3} \tilde{a}(x,0)) v_{1} v_{2} v_{3} v_{4} \, dV = 0.$$

We have proved that (6.16) holds for any solutions v_j with $Pv_j = 0$ for $1 \le j \le 3$ and $\bar{P}^*v_4 = 0$. We now assume that $x_0 \in B$, i.e. there are two maximal null bicharacteristics $\gamma_j : [-S_j, T_j] \to T^*M \setminus 0$, $\gamma_j(t) = (x_j(t), \xi_j(t))$, j = 1, 2, so that the curves x_1 and x_2 only intersect at x_0 when t = 0 and $\dot{x}_1(0) \not\parallel \dot{x}_2(0)$. Let $\gamma_3(t) = (x_3(t), \xi_3(t))$ be the null bicharacteristic with $x_3(0) = x_0$ and $\xi_3(0) = -\xi_1(0)$ (if m is odd $\gamma_3(t) = (x_1(t), -\xi_1(t))$, while if m is even $\gamma_3(t) = (x_1(-t), -\xi_1(-t))$). Similarly, let $\gamma_4(t) = (x_4(t), \xi_4(t))$ be the null bicharacteristic with $x_4(0) = x_0$ and $\xi_4(0) = -\xi_2(0)$ (this is a null bicharacteristic for \bar{P}^* since P has real principal symbol). We consider solutions

$$v_j = q_j + r_j$$

where q_j is a quasimode given by Theorem 3.1 associated with γ_j extended slightly outside M, so that $\|Pq_j\|_{H^s(M)} = O(h^{\infty})$ and q_j is supported in a small neighborhood of the curve $x_j(t)$, and r_j are solutions of $Pr_j = -Pq_j$ in M obtained from Proposition 2.2 and satisfy $\|r_j\|_{H^{s+m-1}(M)} = O(h^{\infty})$. Then (6.16) gives

(6.17)
$$\int_{M} (\partial_{u}^{3} a(x,0) - \partial_{u}^{3} \tilde{a}(x,0)) q_{1} q_{2} q_{3} q_{4} \, dV = O(h^{\infty})$$

Since $x_1(t)$ and $x_2(t)$ only intersect at x_0 , the product $q_1q_2q_3q_4$ is supported in a small neighborhood of x_0 . By (3.5) each q_j has the form

$$q_j(x) = \int_{-S_j}^{T_j} e^{i\Phi_j(x,t_j)/h} a_j(x,t_j) \, dt_j$$

where the maximal null bicharacteristic γ_j is defined on $[-S_j, T_j]$ with $x_j(0) = x_0$ for $1 \le j \le 4$. Using (3.23)–(3.27), the phase functions satisfy

$$\operatorname{Im}(\Phi_j) \ge 0, \qquad \Phi_j(x_0, 0) = 0, \qquad d_{x, t_j} \Phi_j(x_0, 0) = (\xi_j(0), 0),$$

and

$$\nabla_{x,t_j}^2 \Phi_j(x_0,0) = \begin{pmatrix} H_j & -H_j \dot{x}_j + \dot{\xi}_j \\ (-H_j \dot{x}_j + \dot{\xi}_j)^t & (H_j \dot{x}_j - \dot{\xi}_j) \cdot \dot{x}_j \end{pmatrix}$$

where the last expression is computed in Riemannian normal coordinates at x_0 and evaluated at $t_j = 0$. Recall that $\text{Im}(H_j)$ are positive definite matrices.

Inserting the formulas for q_j into (6.17) and writing $\mathbf{t} = (t_1, \ldots, t_4)$, we obtain that

(6.18)
$$\int_M \int_{\mathbb{R}^4} (\partial_u^3 a(x,0) - \partial_u^3 \tilde{a}(x,0)) e^{i\Theta(x,\mathbf{t})/h} b(x,\mathbf{t}) \, d\mathbf{t} \, dV = O(h^\infty)$$

where $b(x, \mathbf{t})$ is supported in a small neighborhood of $(x_0, \mathbf{0})$ in $M^{\text{int}} \times \mathbb{R}^4$ by the assumption that $x_1(t)$ and $x_2(t)$ only intersect once at t = 0. Here we have written

$$\Theta(x, \mathbf{t}) = \Phi_1(x, t_1) + \ldots + \Phi_4(x, t_4), \qquad b(x, \mathbf{t}) = a_1(x, t_1) \cdots a_4(x, t_4).$$

Since we have arranged that

$$\xi_1(0) + \xi_2(0) + \xi_3(0) + \xi_4(0) = 0$$
56

the properties of Φ_i above ensure that

$$\operatorname{Im}(\Theta) \ge 0, \qquad \Theta(x_0, \mathbf{0}) = 0, \qquad d_{x, \mathbf{t}} \Theta(x_0, \mathbf{0}) = 0.$$

To show that the Hessian of Θ is invertible at $(x_0, \mathbf{0})$, we write $\mathbf{z} = (z_1, \dots, z_4)$ and observe that

(6.19)
$$\nabla_{x,\mathbf{t}}^{2}\Theta(x_{0},\mathbf{0})\left(\begin{array}{c}v\\\mathbf{z}\end{array}\right) = \left(\begin{array}{c}\sum_{j=1}^{4}[H_{j}(v-\dot{x}_{j}z_{j})+\dot{\xi}_{j}z_{j}]\\-(H_{1}\dot{x}_{1}-\dot{\xi}_{1})\cdot(v-\dot{x}_{1}z_{1})\\\vdots\\-(H_{4}\dot{x}_{4}-\dot{\xi}_{4})\cdot(v-\dot{x}_{4}z_{4})\end{array}\right)$$

If (6.19) vanishes, this would imply that

$$0 = \nabla_{x,\mathbf{t}}^{2} \operatorname{Im}(\Theta)(x_{0},\mathbf{0}) \begin{pmatrix} v \\ \mathbf{z} \end{pmatrix} \cdot \overline{\begin{pmatrix} v \\ \mathbf{z} \end{pmatrix}} = \sum_{j=1}^{4} |\operatorname{Im}(H_{j})^{1/2}(v - \dot{x}_{j}z_{j})|^{2}$$

and hence $v - \dot{x}_j z_j = 0$ for $1 \leq j \leq 4$. Thus in particular $\dot{x}_1 z_1 = \dot{x}_2 z_2$, and the condition $\dot{x}_1(0) \not\parallel \dot{x}_2(0)$ implies that $z_1 = z_2 = 0$. But the choice of γ_3 and γ_4 implies that when $t_j = 0$,

$$\dot{x}_3 = (-1)^{m-1} \dot{x}_1, \qquad \dot{\xi}_3 = (-1)^m \dot{\xi}_1, \qquad \dot{x}_4 = (-1)^{m-1} \dot{x}_2, \qquad \dot{\xi}_4 = (-1)^m \dot{\xi}_2.$$

Since $v - \dot{x}_j z_j = 0$ and since $\dot{x}_1(0)$ and $\dot{x}_2(0)$ are nonvanishing, this yields $z_3 = (-1)^{m-1} z_1$ and $z_4 = (-1)^{m-1} z_2$. Thus $z_j = 0$ for $1 \le j \le 4$ and also v = 0, showing that $\nabla^2_{x,\mathbf{t}} \Theta(x_0,\mathbf{0})$ is invertible.

Now using stationary phase in (6.18), we obtain

$$c_0(\partial_u^3 a(x_0,0) - \partial_u^3 \tilde{a}(x_0,0))h^{\frac{n+4}{2}} = O(h^{\frac{n+4}{2}+1})$$

where $c_0 \neq 0$ since the amplitudes $a_j(x, t_j)$ are nonvanishing at $(x_0, 0)$. This proves that $\partial_u^3 a(x_0, 0) = \partial_u^3 \tilde{a}(x_0, 0)$ whenever $x_0 \in B$.

We have proved that

$$\partial_u^k a(x_0, 0) = \partial_u^k \tilde{a}(x_0, 0), \qquad k \le 3$$

To prove that this holds also for k = 4, we consider $u_{\varepsilon} = S(\varepsilon_1 v_1 + \ldots + \varepsilon_4 v_4)$ where $Pv_j = 0$ and repeat the above argument (now looking at the $\partial_{\varepsilon_1 \varepsilon_2 \varepsilon_3 \varepsilon_4}$ derivative at $\varepsilon = 0$) to obtain that

$$\int_{M} (\partial_{u}^{4} a(x,0) - \partial_{u}^{4} \tilde{a}(x,0)) v_{1} v_{2} v_{3} v_{4} v_{5} \, dV = 0$$

whenever $\bar{P}^*v_5 = 0$. We now let v_j be a solution related to γ_j above for $1 \leq j \leq 4$, with the difference that now v_4 solves $Pv_4 = 0$ instead of $\bar{P}^*v_4 = 0$. We also choose v_5 to be any solution of $\bar{P}^*v_5 = 0$ with $v_5(x_0) \neq 0$ so that v_5 is independent of h. For instance, it is enough to choose v_5 to be related to the null bicharacteristic γ_1 through $x_0 = x_1(0)$, so that $\dot{x}_1(0) \neq 0$ and $x_1(t) \neq x_1(0)$ for $t \neq 0$ by the assumption that $x_0 \in B$. Then Lemma 3.6 gives that $v_5(x_0) \neq 0$, when $h = h_0$ and h_0 is fixed but sufficiently small. The argument above now gives that

$$(\partial_u^4 a(x_0, 0) - \partial_u^4 \tilde{a}(x_0, 0))v_5(x_0) = 0.$$

Since $v_5(x_0) \neq 0$, we have proved that

$$\partial_u^k a(x_0, 0) = \partial_u^k \tilde{a}(x_0, 0), \qquad k \le 4.$$

Continuing in this way shows that all derivatives of $a(x_0, \cdot) - \tilde{a}(x_0, \cdot)$ vanish at 0, which implies that $a(x_0, \cdot) = \tilde{a}(x_0, \cdot)$ by analyticity.

APPENDIX A. ADDITIONAL PROOFS

In this appendix we give for completeness the proofs of some auxiliary results used in the article.

Proof of Lemma 3.3. Fix some local coordinates and write

$$Pu = \sum_{r=0}^{m} \sum_{j_1,\dots,j_r=1}^{n} p_{j_1\dots j_r}(x) D_{j_1} \cdots D_{j_r}.$$

In terms of these local coordinates

$$p_m(x,\xi) = \sum_{j_1,\dots,j_m=1}^n p_{j_1\dots j_m}(x)\xi_{j_1}\dots\xi_{j_m},$$

$$\partial_{\xi_a} p_m(x,\xi) v_a = \sum_{j_1,\dots,j_m=1}^n p_{j_1\dots j_m}(x)(v_{j_1}\xi_{j_2}\dots\xi_{j_m}+\dots+\xi_{j_1}\dots\xi_{j_{m-1}}v_{j_m}),$$

$$\partial_{\xi_a\xi_b} p_m(x,\xi) v_{ab} = \sum_{j_1,\dots,j_m=1}^n p_{j_1\dots j_m}(x) \Big[(v_{j_1j_2}+v_{j_2j_1})\xi_{j_3}\dots\xi_{j_m} + (v_{j_1j_3}+v_{j_3j_1})\xi_{j_2}\xi_{j_4}\dots\xi_{j_m} + \dots + (v_{j_{m-1}j_m}+v_{j_mj_{m-1}})\xi_{j_1}\dots\xi_{j_{m-2}} \Big].$$

One also has $p_{m-1}(x,\xi) = \sum_{j_1,...,j_{m-1}=1}^n p_{j_1\cdots j_{m-1}}(x)\xi_{j_1}\cdots\xi_{j_{m-1}}$. We now compute

$$e^{-i\Phi/h}P(e^{i\Phi/h}u) = \sum_{r=0}^{m} \sum_{j_1,\dots,j_r=1}^{n} p_{j_1\dots j_r}(x)(D_{j_1} + h^{-1}\partial_{j_1}\Phi)\cdots(D_{j_r} + h^{-1}\partial_{j_r}\Phi)u$$
$$= \sum_{j=0}^{m} h^{j-m}R_ju$$

where each R_j is a differential operator of order j and

$$\begin{aligned} R_{0}u &= p_{m}(x,\nabla\Phi)u, \\ R_{1}u &= p_{m-1}(x,\nabla\Phi) + \sum_{j_{1},\dots,j_{m}} p_{j_{1}\cdots j_{m}}(x) \Big[D_{j_{1}}(\partial_{j_{2}}\Phi\cdots\partial_{j_{m}}\Phi u) \\ &+ \partial_{j_{1}}\Phi D_{j_{2}}(\partial_{j_{3}}\Phi\cdots\partial_{j_{m}}\Phi u) + \dots + \partial_{j_{1}}\Phi\cdots\partial_{j_{m-1}}\Phi D_{j_{m}}u \Big] \\ &= p_{m-1}(x,\nabla\Phi) + \frac{1}{i}\partial_{\xi_{a}}p_{m}(x,\nabla\Phi)\partial_{a}u + \frac{1}{2i}\partial_{\xi_{a}\xi_{b}}p_{m}(x,\nabla\Phi)\partial_{ab}\Phi. \end{aligned}$$

Proof of Lemma 4.2. We argue by contradiction and assume that there is a maximal null bicharacteristic $\gamma : I \to T^*X \setminus 0$ and a compact set $K \subset X$ with $\gamma(I) \subset T^*K$. As before, we write $x(t) = \pi(\gamma(t))$ where $\pi : T^*X \to X$ is the standard projection.

We fix an auxiliary Riemannian metric g on X and consider the projection $\eta(t)$ of $\gamma(t)$ to the unit cosphere bundle S^*X . We think of $\xi(t)$ (i.e. of $\gamma(t)$) as a covector field on x(t), and then $\eta(t)$ corresponds to $\omega(t) = \xi(t)/|\xi(t)|$. Note that $\eta(I)$ is contained in the compact set $S^*K \cap p^{-1}(0)$. The metric g induces the Sasaki metric on T^*X and corresponding horizontal and vertical subbundles, see [Pa99]. The Hamilton vector field H_p on T^*X decomposes in horizontal and vertical parts as

$$H_p = (d_{\xi}p, -\nabla_x p)$$
58

where $d_{\xi}p = \pi_*(H_p) \in TX$ and $\nabla_x p \in T^*X$. By the Hamilton equations and homogeneity

$$\dot{x}(t) = d_{\xi} p(\gamma(t)) = |\xi(t)|^{m-1} d_{\xi} p(\eta(t)),$$

$$D_t \xi(t) = -\nabla_x p(\gamma(t)) = -|\xi(t)|^m \nabla_x p(\eta(t)).$$

Here D_t is the covariant derivative along x(t).

If $[t_1, t_2] \subset I$, we have

$$\phi(x(t_2)) - \phi(x(t_1)) = \int_{t_1}^{t_2} \frac{d}{dt} \phi(x(t)) \, dt = \int_{t_1}^{t_2} d\phi(d_{\xi} p(\gamma(t)) \, dt = \int_{t_1}^{t_2} |\xi(t)|^{m-1} d\phi(d_{\xi} p(\eta(t))) \, dt.$$

Note that since $\tau \mapsto p(y, \zeta_0 + \tau d\phi(y))$ has only simple zeros, differentiating in τ yields that

$$d\phi(d_{\xi}p(y,\zeta)) \neq 0, \qquad (y,\zeta) \in p^{-1}(0).$$

Now $\eta(I)$ is contained in a compact set, so $|d\phi(d_{\xi}p(\eta(t)))| \sim 1$ uniformly over $t \in I$. Taking limits where t_j converge to the endpoints of I, we see that

(A.1)
$$\int_{I} |\xi(t)|^{m-1} dt < \infty.$$

On the other hand, we compute

$$\frac{d}{dt}\frac{1}{|\xi(t)|^{m-1}} = -(m-1)\frac{\langle D_t\xi(t),\xi(t)\rangle}{|\xi(t)|^{m+1}} = (m-1)\langle \nabla_x p(\eta(t)),\omega(t)\rangle$$

Again, since $\eta(I)$ is contained in a compact set, we have $1/|\xi(t)|^{m-1} \leq C(1+|t|)$. In other words, for some c > 0 one has

(A.2)
$$|\xi(t)|^{m-1} \ge \frac{c}{1+|t|}, \quad t \in I.$$

Now the combination of (A.1) and (A.2) implies that I must be a finite interval.

We have also

$$\frac{d}{dt}|\xi(t)| = \frac{\langle D_t\xi(t),\xi(t)\rangle}{|\xi(t)|} = -|\xi(t)|^m \langle \nabla_x p(\eta(t)),\omega(t)\rangle \le C|\xi(t)|^m, \quad t \in I.$$

Hence using (A.1) we see that $\ln|\xi(t)|$ is bounded from above on I. Due to this and (A.2) one has $C^{-1} \leq |\xi(t)| \leq C$ for some C > 0 and all $t \in I$. It follows that $\gamma(I)$ is contained in a compact set of $T^*X \setminus 0$. This again implies, together with $\dot{\gamma}(t) = H_p(\gamma(t))$, that $\gamma : I \to T^*X$ is uniformly continuous. Hence γ extends to \overline{I} as a continuous function. We may assume without loss of generality that $\inf I = 0$. Then $\lim_{t\to 0} \dot{\gamma}(t) = H_p(\gamma(0))$ and also $\dot{\gamma}$ is continuous up to 0. Therefore γ is the bicharacteristic from $\gamma(0)$ and thus it can be extended on $(-\epsilon, 0)$ for small $\epsilon > 0$, a contradiction with maximality of γ .

Proof of Theorem 4.3. Let us show (4.10). The proof is similar with Step 8 in the proof of Theorem 1.1, and it is enough to show

(A.3)
$$\exp\left[i\int_0^T q_{m-1}(\gamma(t))\,dt\right] = 1,$$

where q_{m-1} is the principal symbol of $Q = P_1 - P_2$. We denote by $\check{\gamma}(t) = (x(t), \xi(t))$ the continuation of γ as a null bicharacteristic curve in X.

We use Theorem 3.1 with $P = P_1$ and $\tilde{P} = P_2$ and construct quasimodes v_1, v_2 associated with $\check{\gamma}|_{[-\varepsilon_1, T+\varepsilon_2]}$ with ε_j chosen so that the end points are outside M and $x|_{[-\varepsilon_1, T+\varepsilon_2]}$ does not intersect $\partial M \setminus \Gamma$. As x(t) was assumed to intersect Γ transversally, we can also arrange that $x|_{[-\varepsilon_1,T+\varepsilon_2]}$ only intersects ∂M at x(0) and x(T). The quasimodes v_j can be constructed so that $\operatorname{supp}(v_j) \cap (\partial M \setminus \Gamma) = \emptyset$. Then $\|P_1v_1\|_{H^s(M)} = \|P_2^*v_2\|_{H^s(M)} = O(h^\infty)$. Using Proposition 2.2, we construct again solutions $u_j = v_j + r_j$ of $P_1u_1 = 0$ and $P_2^*u_2 = 0$ in M with $\|r_j\|_{H^m(M)} = O(h^\infty)$.

Let us show that $\operatorname{tr}_{\Gamma}^{m-1} u_1 \in C_{P_1,\Gamma}$. As

$$\|\operatorname{tr}_{\partial M\setminus\Gamma}^{m-1} u_1\|_{\mathcal{H}^{m-1}(\partial M\setminus\Gamma)} = \|\operatorname{tr}_{\partial M\setminus\Gamma}^{m-1} r_1\|_{\mathcal{H}^{m-1}(\partial M\setminus\Gamma)} = O(h^{\infty}),$$

it is enough to show that $||v_1|_{\Gamma}||_{L^2(\Gamma)}$ is not $O(h^{\infty})$. Since $x|_{[-\varepsilon_1, T+\varepsilon_2]}$ only meets ∂M at x(0) and x(T), it is enough to show that $||v_1|_{\tilde{\Gamma}}||_{L^2(\tilde{\Gamma})} \sim h^{\frac{n+1}{2}}$ where $\tilde{\Gamma} \subset \Gamma$ is a small neighborhood of x(0).

We argue similarly to Step 8 of the proof of Theorem 3.1. One has

$$\int_{\tilde{\Gamma}} |v_1|^2 \, dS = \int_{-\varepsilon_1}^{T+\varepsilon_2} \int_{-\varepsilon_1}^{T+\varepsilon_2} \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s) \, dS \, dt \, ds$$

where a does not vanish at (x(s), s, s) and

$$\Theta(x,t,s) = \Phi(x,t) - \overline{\Phi(x,s)},$$

with Φ as in the proof of Theorem 3.1. Note also that if $\chi(t,s)$ is a cutoff function supported near (0,0) one has

$$\int_{\tilde{\Gamma}} |v_1|^2 \, dS = \int_{-\varepsilon_1}^{T+\varepsilon_2} \int_{-\varepsilon_1}^{T+\varepsilon_2} \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{-C/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,s)/h} a(x,t,s)\chi(t,s) \, dS \, dt \, ds + O(e^{i\Theta(x,t,s)/h} ds \, ds + O(e^{i\Theta(x,t,s)/h}) \int_{\tilde{\Gamma}} e^{i\Theta(x,t,$$

since $\operatorname{Im}(\Theta) \ge c > 0$ when $x \in \tilde{\Gamma}$ and either s or t is bounded away from 0. We have

Im(
$$\Theta$$
) ≥ 0 on supp(a), $\Theta(x(0), 0, 0) = 0$, $d_{x,t,s}\Theta(x(0), 0, 0) = 0$,

and $\|v_1|_{\tilde{\Gamma}}\|_{L^2(\tilde{\Gamma})} \sim h^{\frac{n+1}{2}}$ follows from the method of stationary phase after we have shown that $\nabla^2_{x,t,s}\Theta$ is non-singular at (x(0), 0, 0). Analogously to (3.31), the Hessian of Θ is given by

$$\nabla^2_{x,t,s}\Theta|_{(x(0),0,0)} = \begin{pmatrix} 2i\mathrm{Im}(H) & \dot{\xi} - H\dot{x} & -\dot{\xi} + \overline{H}\dot{x} \\ (\dot{\xi} - H\dot{x})^t & (H\dot{x} - \dot{\xi}) \cdot \dot{x} & 0 \\ (-\dot{\xi} + \overline{H}\dot{x})^t & 0 & -(\overline{H}\dot{x} - \dot{\xi}) \cdot \dot{x} \end{pmatrix}.$$

Let $\zeta = (v, z, w)^t$, with $v = v_1 + iv_2$, $v_j \in T_{x(0)}\Gamma$, $z, w \in \mathbb{C}$, be in the kernel of the above matrix, then

$$0 = \operatorname{Im}(\nabla_{x,t,s}^2 \Theta|_{(x(0),0,0)}) \zeta \cdot \overline{\zeta} = |\operatorname{Im}(H)^{1/2} (v - z\dot{x})|^2 + |\operatorname{Im}(H)^{1/2} (v - w\dot{x})|^2.$$

As Im(H) is positive definite and $\dot{x}(0) \notin T_{x(0)}\Gamma$, we see that v = 0 and z = w = 0. Hence $\nabla_{x,t,s}^2 \Theta$ is non-singular at (x(0), 0, 0) and $\|v_1|_{\tilde{\Gamma}}\|_{L^2(\tilde{\Gamma})} \sim h^{\frac{n+1}{2}}$.

We have now proved that $\operatorname{tr}_{\Gamma}^{m-1} u_1 \in C_{P_1,\Gamma}$. We will need the following variation of Lemma 2.8, (A.4) $|((P_1 - P_2)u_1, u_2)_{L^2(M)}| = O(h^{\infty}).$

To show (A.4), we observe that due to $C_{P_1,\Gamma} \subset C_{P_2,\Gamma}$ there is $\tilde{u}_2 \in H^m(M)$ such that

(A.5) $P_2 \tilde{u}_2 = 0 \text{ and } \nabla^k u_1|_{\Gamma} = \nabla^k \tilde{u}_2|_{\Gamma} \text{ for } k = 0, 1, \dots, m-1.$

Therefore

$$|((P_1 - P_2)u_1, u_2)_{L^2(M)}| = |(P_2(u_1 - \tilde{u}_2), u_2)_{L^2(M)}| \leq C \|\operatorname{tr}_{\partial M \setminus \Gamma}^{m-1}(u_1 - \tilde{u}_2)\|_{\mathcal{H}^{m-1}(\partial M \setminus \Gamma)} \|\operatorname{tr}_{\partial M \setminus \Gamma}^{m-1} u_2\|_{\mathcal{H}^{m-1}(\partial M \setminus \Gamma)}.$$

We have $\|\operatorname{tr}_{\partial M\setminus\Gamma}^{m-1} u_2\|_{\mathcal{H}^{m-1}(\partial M\setminus\Gamma)} = O(h^{\infty})$. The definition of $C_{P_j,\Gamma}$, together with (A.5), implies that

$$\|\operatorname{tr}_{\partial M\setminus\Gamma}^{m-1}(u_1-\tilde{u}_2)\|_{\mathcal{H}^{m-1}(\partial M\setminus\Gamma)} \leq 2\|\operatorname{tr}_{\Gamma}^{m-1}u_1\|_{\mathcal{H}^{m-1}(\Gamma)}$$

Finally, for k large the trace theorem gives that $\|\operatorname{tr}_{\Gamma}^{m-1} u_1\|_{\mathcal{H}^{m-1}(\Gamma)} \lesssim \|u_1\|_{H^k(M)} = O(h^{\frac{n+1}{4}-k})$ by (3.38).

As $Q = P_1 - P_2$ vanishes outside M, the identity (A.4) implies that

$$O(h^{\infty}) = (Qu_1, u_2)_{L^2(M)} = (Qv_1, v_2)_{L^2(X)} + O(h^{\infty}).$$

Multiplying this identity by $h^{-\frac{n+1}{2}+m-1}$ and using (3.1), we obtain (A.3) as in the proof of Theorem 1.1.

We omit giving a proof of (4.11). The modifications needed in the proof of Theorem 1.2 are analogous to those above. $\hfill \Box$

Proof of Theorem 4.4. We first give some initial remarks related to a first order real principal type differential operator P on M. Note that P is of the form $P = \frac{1}{i}L + V$ where L is a real vector field on M and $V \in C^{\infty}(M)$. The null bicharacteristic curves for P are of the form $\gamma(t) = (x(t), \xi(t))$ where x(t) is an integral curve for L in M. The real principal type condition implies that for any $x_0 \in M$, the maximal integral curve $x : [-\tau_-^L(x_0), \tau_+^L(x_0)] \to M$ of L with $x(0) = x_0$ has finite length, i.e. $\tau_{\pm}^L(x_0) < \infty$. Moreover, the equation Pu = 0 in M becomes an ODE along each fixed integral curve, and thus any solution u satisfies

$$u(x(t)) = u(x(0)) \exp\left[-i \int_0^t V(x(s)) \, ds\right], \qquad 0 \le t \le T,$$

whenever $x: [0,T] \to M$ is a maximal integral curve of L.

Now suppose that $P_j = \frac{1}{i}L_j + V_j$ are as in the theorem. The assumption that $\alpha_{P_1} = \alpha_{P_2}$, together with the fact that the spatial projections of null bicharacteristics are integral curves, implies that

$$x_1(T_1) = x_2(T_2)$$

whenever $x_j : [0, T_j] \to M$ are maximal integral curves of L_j with $x_1(0) = x_2(0)$.

Let $f_1 \in C_{P_1}$, so that $f_1 = u_1|_{\partial M}$ where $u_1 \in C^{\infty}(M)$ and $P_1u_1 = 0$. For any maximal L_2 -integral curve $x_2 : [0, T_2] \to M$, define

$$u_2(x_2(t)) := f_1(x_2(0)) \exp\left[-i \int_0^t V_2(x_2(s)) \, ds\right]$$

This defines a function $u_2: M \to \mathbb{C}$ that is smooth along integral curves of L_2 and satisfies $P_2 u_2 = 0$. We claim that

$$u_2|_{\partial M} = u_1|_{\partial M}.$$

In fact, this holds at any $x \in \partial M$ with $\tau_{-}^{L_2}(x) = 0$ by definition. On the other hand, if $x \in \partial M$ and $\tau_{+}^{L_2}(x) = 0$, then $x = x_2(T_2)$ for some maximal L_2 -integral curve $x_2 : [0, T_2] \to M$, and

$$u_2(x) = u_2(x_2(T_2)) = f_1(x_2(0)) \exp\left[-i \int_0^{T_2} V_2(x_2(s)) \, ds\right].$$

By assumption, if $x_1 : [0, T_1] \to M$ is the maximal L_1 -integral curve with $x_1(0) = x_2(0)$, then also $x_1(T_1) = x_2(T_2)$ and $\exp\left[-i\int_0^{T_1} V_1(x_1(s)) ds\right] = \exp\left[-i\int_0^{T_2} V_2(x_2(s)) ds\right]$. It follows that

$$u_2(x) = f_1(x_1(0)) \exp\left[-i \int_0^{T_1} V_1(x_1(s)) \, ds\right] = u_1(x_1(T_1)) = u_1(x).$$

Thus we have proved that $u_1(x) = u_2(x)$ at each $x \in \partial M$ so that $\tau_{-}^{L_2}(x) = 0$ or $\tau_{+}^{L_2}(x) = 0$, but this covers all points of ∂M by strict convexity.

For any $f_1 \in C_{P_1}$, we have produced a function u_2 in M that is smooth along integral curves of L_2 and satisfies $P_2u_2 = 0$ with $u_2|_{\partial M} = f_1$. If we can show that $u_2 \in C^{\infty}(M)$, then we have

$$C_{P_1} \subset C_{P_2}.$$

The converse inclusion follows by changing the roles of P_1 and P_2 . Thus it remains to prove a regularity statement: if M is compact with smooth boundary, if L is a nontrapping real vector field in M so that M is strictly convex for L, and if $u: M \to \mathbb{C}$ solves $\frac{1}{i}Lu + Vu = 0$ in M with $u|_{\partial M} \in C^{\infty}(\partial M)$, then $u \in C^{\infty}(M)$. This is proved in [PU05, Lemma 1.1] when M is the unit sphere bundle, L is the geodesic vector field and V = 0, and in [PSU12, Lemma 5.1] for $V \neq 0$. However, inspecting the arguments in [PU05, PSU12] shows that the proofs also give the more general result stated above. The details will appear in [PSU19].

References

- [AB59] Y. Aharonov, D. Bohm, Significance of electromagnetic potentials in quantum theory, Phys. Rev. 115 (1959), 485–491.
- [AS19] Y.M. Assylbekov, P. Stefanov, *Sharp stability estimate for the geodesic ray transform*, Inverse Problems (to appear).
- [BL67] V.M. Babich, V.F. Lazutkin, The eigenfunctions which are concentrated near a closed geodesic (in Russian), 1967, Problems of Mathematical Physics, No. 2, Spectral Theory, Diffraction Problems, pp. 15–25, Izdat. Leningrad. Univ., Leningrad.
- [Be87] M.I. Belishev. An approach to multidimensional inverse problems for the wave equation, Doklady Akademii Nauk SSSR, **297** (1987), 524–527.
- [CLV19] M. Capoferri, M. Levitin, D. Vassiliev, Geometric wave propagator on Riemannian manifolds, arXiv:1902.06982.
- [DKSU09] D. Dos Santos Ferreira, C.E. Kenig, M. Salo, G. Uhlmann, Limiting Carleman weights and anisotropic inverse problems, Invent. Math. 178 (2009), 119–171.
- [DKLS16] D. Dos Santos Ferreira, Y. Kurylev, M. Lassas, M. Salo, The Calderón problem in transversally anisotropic geometries, J. Eur. Math. Soc. (JEMS) 18 (2016), 2579–2626.
- [DH72] J.J. Duistermaat, L. Hörmander, Fourier integral operators. II. Acta Math. 128 (1972), 183–269.
- [EHN96] H.W. Engl, M. Hanke, A. Neubauer, Regularization of inverse problems. Dordrecht: Kluwer Academic Publishers Group, 1996.
- [Es15] G. Eskin, Aharonov–Bohm effect revisited, Rev. Math. Phys. 27 (2015), no. 2, 1530001.
- [FIKO19] A. Feizmohammadi, J. Ilmavirta, Y. Kian, L. Oksanen, Recovery of time dependent coefficients from boundary data for hyperbolic equations, arXiv:1901.04211.
- [FIO19] A. Feizmohammadi, J. Ilmavirta, L. Oksanen, The light ray transform in stationary and static Lorentzian geometries, arXiv:1911.04834.
- [FO19] A. Feizmohammadi, L. Oksanen, An inverse problem for a semi-linear elliptic equation in Riemannian geometries, arXiv:1904.00608.
- [Ge54] I.M. Gel'fand, Some aspects of functional analysis and algebra. Proc. Intern. Cong. Math. 1 (1954), 253– 277.
- [Gu17] C. Guillarmou, Lens rigidity for manifolds with hyperbolic trapped sets, J. Amer. Math. Soc. **30** (2017), 561–599.
- [Gu76] V. Guillemin, Sojourn times and asymptotic properties of the scattering matrix, Proceedings of the Oji Seminar on Algebraic Analysis and the RIMS Symposium on Algebraic Analysis (Kyoto Univ., Kyoto, 1976). Publ. Res. Inst. Math. Sci., 12, supplement, 69–88 (1976/77).

- [GUW16] V. Guillemin, A. Uribe, Z. Wang, Semiclassical states associated with isotropic submanifolds of phase space, Lett. Math. Phys. 106 (2016), 1695–1728.
- [Hö71] L. Hörmander, On the existence and the regularity of linear pseudo-differential equations, L'Enseignement Math. 17 (1971), 99–163.
- [Hö85] L. Hörmander, The analysis of linear partial differential operators I-IV. Springer, 1983–1985.
- [IM19] J. Ilmavirta, F. Monard, Integral geometry on manifolds with boundary and applications, chapter in "The Radon Transform: The First 100 Years and Beyond" (Ronny Ramlau, Otmar Scherzer, eds.), de Gruyter, 2019.
- [Is91] V. Isakov, Completeness of products of solutions and some inverse problems for PDE, J. Diff. Eq. 92 (1991), 305–316.
- [KV98] M. Karasev, Y. Vorobjev, Integral representations over isotropic submanifolds and equations of zero curvature, Adv. Math. 135 (1998), 220–286.
- [KKL01] A. Katchalov, Y. Kurylev, M. Lassas, *Inverse Boundary Spectral Problems*, Monographs and Surveys in Pure and Applied Mathematics 123, Chapman Hall/CRC, 2001.
- [KP88] T. Kato, G. Ponce, Commutator estimates and the Euler and Navier-Stokes equations, Comm. Pure Appl. Math. 41 (1988), 891–907.
- [KS13] C.E. Kenig, M. Salo, The Calderón problem with partial data on manifolds and applications, Analysis & PDE 6 (2013), 2003–2048.
- [KV84] R. Kohn, M. Vogelius, Determining conductivity by boundary measurements, Comm. Pure Appl. Math. 37 (1984), 289–298.
- [KLOU14] Y. Kurylev, M. Lassas, L. Oksanen, G. Uhlmann, Inverse problems for Einstein-scalar field equations, arXiv:1406.4776.
- [KLU18] Y. Kurylev, M. Lassas, G. Uhlmann, Inverse problems for Lorentzian manifolds and non-linear hyperbolic equations, Invent. Math. 212 (2018), 781–857.
- [LSV94] A. Laptev, Y. Safarov, D. Vassiliev, On global representation of Lagrangian distributions and solutions of hyperbolic equations, Comm. Pure Appl. Math. 47 (1994), 1411–1456.
- [LLLS19] M. Lassas, T. Liimatainen, Y.-H. Lin, M. Salo, Inverse problems for elliptic equations with power type nonlinearities, arXiv:1903.12562.
- [LOSU19] M. Lassas, L. Oksanen, P. Stefanov, G. Uhlmann, The light ray transform on Lorentzian manifolds, arXiv:1907.02210.
- [LUW18] M. Lassas, G. Uhlmann, Y. Wang, Inverse problems for semilinear wave equations on Lorentzian manifolds, Comm. Math. Phys. 360 (2018), 555–609.
- [LU89] J.M. Lee, G. Uhlmann, Determining anisotropic real-analytic conductivities by boundary measurements, Comm. Pure Appl. Math. 42 (1989), 1097–1112.
- [MS74] A. Melin, J. Sjöstrand, Fourier integral operators with complex-valued phase functions, Springer Lecture Notes in Mathematics **459** (1974), 120–223.
- [Mo14] C. Montalto, Stable determination of a simple metric, a covector field and a potential from the hyperbolic Dirichlet-to-Neumann map, Comm. PDE **39** (2014), 120–145.
- [O'N83] B. O'Neill, Semi-Riemannian geometry, Academic Press Inc, 1983.
- [Pa99] G.P. Paternain, Geodesic flows. Progress in Mathematics 180, Birkhäuser, 1999.
- [PSU12] G.P. Paternain, M. Salo, G. Uhlmann, The attenuated ray transform for connections and Higgs fields, Geom. Funct. Anal. 22 (2012), 1460–1489.
- [PSU19] G.P. Paternain, M. Salo, G. Uhlmann, Geometric inverse problems in two dimensions. Textbook in preparation.
- [PU93] T. Paul, A. Uribe, A construction of quasi-modes using coherent states, Annales de l'I.H.P., section A 59 (1993), 357–381.
- [PU05] L. Pestov, G. Uhlmann, Two dimensional compact simple Riemannian manifolds are boundary distance rigid, Ann. of Math. 161 (2005), 1089–1106.
- [Ra90] Rakesh, Reconstruction for an inverse problem for the wave equation with constant velocity, Inverse Problems 6 (1990), 91–98.
- [RS88] Rakesh, W.W. Symes, Uniqueness for an inverse problem for the wave equation, Comm. PDE 13 (1988), 87–96.
- [Ra82] J. Ralston, Gaussian beams and the propagation of singularities, Studies in partial differential equations, 206–248, MAA Stud. Math. 23, Math. Assoc. America, Washington, DC, 1982.
- [Ra01] J. Ralston, Gaussian beams. Unpublished note, Institute for Pure and Applied Mathematics, April 18-19, 2001.

- [RR04] M. Renardy, R.C. Rogers, An introduction to partial differential equations. Texts in applied mathematics, Springer, 2004.
- [Ri09] H. Ringström, *The Cauchy Problem in General Relativity*, European Mathematical Society Publishing House, 2009.
- [SV97] Y. Safarov, D. Vassiliev, The asymptotic distribution of eigenvalues of partial differential operators, Amer. Math. Soc., Providence (RI), 1997.
- [Sa17] M. Salo, The Calderón problem and normal forms, arXiv:1702.02136.
- [St89] P. Stefanov, Uniqueness of the multi-dimensional inverse scattering problem for time dependent potentials, Math. Z. 201 (1989), 541–559.
- [St17] P. Stefanov, Support theorems for the light ray transform on analytic Lorentzian manifolds, Proc. Amer. Math. Soc. 145 (2017), 1259–1274.
- [SU05] P. Stefanov, G. Uhlmann, Boundary rigidity and stability for generic simple metrics, J. Amer. Math. Soc. 18 (2005), 975–1003.
- [SUV16] P. Stefanov, G. Uhlmann, A. Vasy, *Boundary rigidity with partial data*, J. Amer. Math. Soc. **29** (2016), 299–332.
- [SUV17] P. Stefanov, G. Uhlmann, A. Vasy, Local and global boundary rigidity and the geodesic X-ray transform in the normal gauge, arXiv:1702.03638.
- [SY18] P. Stefanov, Y. Yang, The inverse problem for the Dirichlet-to-Neumann map on Lorentzian manifolds, Analysis & PDE 11 (2018), 1381–1414.
- [SU88] J. Sylvester, G. Uhlmann, Inverse boundary value problems at the boundary continuous dependence, Comm. Pure Appl. Math. 41 (1988), 197–219.
- [SU90] J. Sylvester, G. Uhlmann, Inverse problems in anisotropic media, Contemp. Math. 122 (1991), Inverse scattering and applications (Amherst, MA, 1990), 105–117.
- [Ta04] D. Tataru, Unique continuation problems for partial differential equations, Geometric methods in inverse problems and PDE control, 239–255, IMA Vol. Math. Appl., 137, Springer, New York, 2004.
- [Ta81] M.E. Taylor, Pseudodifferential operators. Princeton Mathematical Series 34, Princeton University Press, 1981.
- [Tr80] F. Treves, Introduction to pseudodifferential and Fourier integral operators, vol. 2. The University series in mathematics, Plenum Press, 1980.
- [Uh04] G. Uhlmann, *The Cauchy data and the scattering relation*, Geometric methods in inverse problems and PDE control, 263–287, IMA Vol. Math. Appl., 137, Springer, New York, 2004.
- [VW19] A. Vasy, Y. Wang, On the light ray transform with wave constraints, arXiv:1912.02848.
- [Zw12] M. Zworski, Semiclassical analysis. AMS, 2012.

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON

 $E\text{-}mail \ address: \texttt{l.oksanen@ucl.ac.uk}$

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ

 $E\text{-}mail \ address: \texttt{mikko.j.salo@jyu.fi}$

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY

E-mail address: stefanop@purdue.edu

Department of Mathematics, University of Washington / Jockey Club Institute for Advanced Study, HKUST, Hong Kong

E-mail address: gunther@math.washington.edu