

**DIFFERENTIAL GEOMETRY/PDE/INVERSE  
PROBLEM SEMINAR**

MONDAY, NOVEMBER 16, 2009

RAI 116

4PM–5PM

Quantum decay rates for manifolds with hyperbolic ends

**Kiril Datchev**

(UC BERKELEY)

Mathematically, quantum decay rates appear as imaginary parts of poles of the meromorphic continuation of Green's functions. As energy grows, decay rates are related to properties of geodesic flow and to the structure at infinity. For a cusp, infinity is "small", which typically slows decay. However, I will present a class of examples for which decay rates go to infinity with energy even in the presence of a cusp. This is part of a more general investigation of resonances on manifolds with hyperbolic ends.

For more information about this seminar, visit the DG/PDE Seminar Web page (from the Math Department home page, [www.math.washington.edu](http://www.math.washington.edu), follow the link **Seminars, Colloquia, and Conferences**).

The University of Washington is committed to providing access, equal opportunity and reasonable accommodation in its services, programs, activities, education and employment for individuals with disabilities. To request disability accommodation contact the Disability Services Office at least ten days in advance at: 206-543-6450/V, 206-543-6452/TTY, 206-685-7264 (FAX), or [dso@u.washington.edu](mailto:dso@u.washington.edu).