## DIFFERENTIAL GEOMETRY/PDE SEMINAR

Wednesday, July 17, 2002 Smith 107 3:45 pm

## Electromagnetism, ellipticity and conformal Hodge theory

## ROD GOVER (UNIVERSITY OF AUCKLAND)

On a Riemannian signature manifold the Maxwell equations are elliptically deficient. This can be cured by adding extra equations to the system—so called gauge fixing equations. The celebrated Coulomb gauge is an example. However with this and with most other choices the extended system fails to be conformally invariant even though the Maxwell equations are. In fact there is an elliptic extension of the Maxwell system that is well defined on conformal 4-manifolds. Under mild restrictions, the conformally invariant null space of this operator is isomorphic to the first de Rham cohomology. Ideas from Lie representation theory show how to extend this conformally invariant de Rham Hodge theory to other dimensions and also to related BGG resolutions such as the (conformal) metric deformation complex.

For more information about this seminar, visit the DG/PDE Seminar Web page (from the Math Department home page, www.math.washington.edu, follow the link Seminars, Colloquia, and Conferences).

The University of Washington is committed to providing access, equal opportunity and reasonable accommodation in its services, programs, activities, education and employment for individuals with disabilities. To request disability accommodation contact the Disability Services Office at least ten days in advance at: 206-543-6450/V, 206-543-6452/TTY, 206-685-7264 (FAX), or dso@u.washington.edu.