

Closing Wed: HW_1A, 1B, 1C

Entry Task (You do): Approx. the area under $f(x) = x^3$ from $x = 0$ to $x = 1$ using $n = 4$ and *right-endpoints*.

Step 1: $\Delta x = \frac{b-a}{n} = \frac{1-0}{4} = \frac{1}{4}$

Step 2: $x_0 = a = 0$

$x_1 = a + \Delta x = 0 + \frac{1}{4}$

$x_2 = a + 2\Delta x = 0 + 2(\frac{1}{4})$

$x_3 = a + 3\Delta x = 0 + 3(\frac{1}{4})$

$x_4 = a + 4\Delta x = 0 + 4(\frac{1}{4})$

Step 3: Plug in right-endpoints to function to get rect. heights, then add up areas (height times width).

$$\text{Area} \approx \sum_{i=1}^4 f(x_i)\Delta x =$$

$$f(x_1)\Delta x + f(x_2)\Delta x + f(x_3)\Delta x + f(x_4)\Delta x$$

$$\left(\frac{1}{4}\right)^3 \frac{1}{4} + \left(\frac{2}{4}\right)^3 \frac{1}{4} + \left(\frac{3}{4}\right)^3 \frac{1}{4} + \left(\frac{4}{4}\right)^3 \frac{1}{4}$$


Right-endpoints

$$= 0.390625$$

$$\left(\frac{1}{4}\right)^3 \frac{1}{4} + \left(\frac{2}{4}\right)^3 \frac{1}{4} + \left(\frac{3}{4}\right)^3 \frac{1}{4} + \left(\frac{4}{4}\right)^3 \frac{1}{4}$$

PATTERN $\sum_{i=1}^4 \left(\frac{i}{4}\right)^3 \frac{1}{4} = \frac{1}{4^4} \sum_{i=1}^4 i^3$

ASIDE



I did this example again with 100 subdivisions, then 1000, then 10000. Here is a summary of my findings:

n	R_n	L_n
4	0.390625	0.140625
5	0.36	0.16
10	0.3025	0.2025
100	0.255025	0.245025
1000	0.25050025	0.24950025
10000	0.2499500025	0.2500500025

Pattern:

$$\Delta x = \frac{1-0}{n} = \frac{1}{n}, \quad x_i = 0 + i \frac{1}{n} = \frac{i}{n}$$

Adding up the area of each rectangle

$$\text{Sum} = \sum_{i=1}^n x_i^3 \Delta x = \sum_{i=1}^n \left(\frac{i}{n}\right)^3 \frac{1}{n}$$

$$\text{Area} = 0.25 = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(\frac{i}{n}\right)^3 \frac{1}{n}$$

$\lim_{n \rightarrow \infty} \left(\frac{1}{n^4} \sum_{i=1}^n i^3 \right)$
 ASIDE

Example: Approximate the area under $f(x) = 1 + x^2$ from $x = 2$ to $x = 3$ using Riemann sums with $n = 4$ and right endpoints.

$$\Delta x = \frac{3-2}{4} = \frac{1}{4}$$


$$x_0 = 2$$

$$x_1 = 2 + \frac{1}{4}$$

$$x_2 = 2 + 2\left(\frac{1}{4}\right)$$

$$x_3 = 2 + 3\left(\frac{1}{4}\right)$$

$$x_4 = 2 + 4\left(\frac{1}{4}\right)$$



$$(1 + (2.25)^2) \frac{1}{4} + (1 + (2.5)^2) \frac{1}{4} + (1 + (2.75)^2) \frac{1}{4} + (1 + (3)^2) \frac{1}{4} = 7.96875$$

What is the general pattern in terms of n ?

$$\Delta x = \frac{3-2}{n} = \frac{1}{n}$$

$$x_i = 2 + i \frac{1}{n} = 2 + \frac{i}{n}$$

$$\sum_{i=1}^n f(x_i) \Delta x = \sum_{i=1}^n (1 + x_i^2) \Delta x$$

$$= \sum_{i=1}^n \left(1 + \left(2 + \frac{i}{n} \right)^2 \right) \frac{1}{n}$$

\uparrow a \leftarrow $b-a$

Another Example:

Using sigma notation, write down the general Riemann sum definition of the area from $x = 5$ to $x = 7$ under

$$f(x) = 3x + \sqrt{x}$$

$$\Delta x = \frac{b-a}{n} = \frac{7-5}{n} = \frac{2}{n}$$

$$x_i = a + i\Delta x = 5 + i\left(\frac{2}{n}\right) = 5 + \frac{2i}{n}$$

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i)\Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n (3x_i + \sqrt{x_i}) \Delta x$$

$$= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(3\left(5 + \frac{2i}{n}\right) + \sqrt{5 + \frac{2i}{n}} \right) \frac{2}{n}$$

$a = 5$ $b - a = 2$

Velocity/Distance & Reimann Sums

When velocity is a **constant**:

$$\text{Distance} = \text{Velocity} \cdot \text{Time}$$

Example:

You are accelerating in a car. You get the following measurements:

t (sec)	0	0.5	1.0	1.5	2.0
v(t) (ft/s)	0	6.2	10.8	14.9	18.1

Estimate the distance traveled by the car traveled from 0 to 2 seconds.

HAVE TO BREAK IT UP!!

	LOW ESTIMATE	HI ESTIMATE
0 to 0.5	$0 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 0 \text{ ft}$	$6.2 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 3.1 \text{ ft}$
0.5 to 1	$6.2 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 3.1 \text{ ft}$	$10.8 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 5.4 \text{ ft}$
1 to 1.5	$10.8 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 5.4 \text{ ft}$	$14.9 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 7.45 \text{ ft}$
1.5 to 2	$14.9 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 7.45 \text{ ft}$	$18.1 \frac{\text{ft}}{\text{sec}} \cdot 0.5 \text{ sec} = 9.05 \text{ ft}$
TOTAL =	15.95 ft	25 ft

ASIDE: UNITS = $\frac{\text{ft}}{\text{sec}} \cdot \text{sec} = \text{ft}$

\swarrow "HEIGHT" UNITS \uparrow "WIDTH" UNITS

5.2 The Definite Integral

Def'n:

We define the **definite integral of $f(x)$ from $x = a$ to $x = b$** by

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x, \quad \leftarrow$$

where $\Delta x = \frac{b-a}{n}$ and $x_i = a + i\Delta x$.


NOTES:

" \int " = integral sign

a, b = bounds or limits of integration

$\int_a^b f(x) dx$ = "ADD UP" $f(x_i) \Delta x$
ACROSS THE INTERVAL

= A NUMBER.


Ex)  $y = 7$

$$\int_0^3 7 dx = 21$$

$$\int_0^3 -5 dx = -15$$

"SIGNED" or "NET" AREA

$$\int_{-\pi}^{\pi} \sin(x) dx = 0$$



Basic Integral Rules:

$$1. \int_a^b c \, dx = (b - a)c$$

$$2. \int_a^b f(x) \, dx + \int_b^c f(x) \, dx = \int_a^c f(x) \, dx$$

$$3. \int_a^b cf(x) \, dx = c \int_a^b f(x) \, dx$$

and

$$\int_a^b f(x) + g(x) \, dx = \int_a^b f(x) \, dx + \int_a^b g(x) \, dx$$


$$4. \int_b^a f(x) \, dx = - \int_a^b f(x) \, dx$$

Examples:

$$1. \int_4^{10} 5 \, dx = 5(10 - 4) = 30$$

$$2. \int_0^3 x^2 \, dx + \int_3^7 x^2 \, dx = \int_0^7 x^2 \, dx$$

$$3. \int_0^4 5x + 3 \, dx = \int_0^4 5x \, dx + \int_0^4 3 \, dx$$
$$= 5 \int_0^4 x \, dx + \int_0^4 3 \, dx$$



$$= 5 \cdot 8 + 12$$
$$= \boxed{52}$$

$$4. \int_3^1 x^3 \, dx = - \int_1^3 x^3 \, dx$$

Note on quick bounds (HW_1C: 9,10)

$$m(b - a) \leq \int_a^b f(x) dx \leq M(b - a)$$

Example: Consider the area under

$$f(x) = \sin(x) + 2$$


on the interval $x = 0$ to $x = 2\pi$.

(a) What is the max of $f(x)$? (label M)

(b) What is the min of $f(x)$? (label m)

(c) Draw **one** rectangle that contains all the shaded area? What can you conclude?

(d) Draw **one** rectangle that is completely inside the shaded area? Conclusion?



$$m \leq \sin(x) + 2 \leq M$$

so $\int_0^{2\pi} \sin(x) + 2 dx$

MUST BE BETWEEN THE AREA OF THESE TWO RECTANGLES

$$\underbrace{1 \cdot 2\pi}_{2\pi} \leq \int_0^{2\pi} \sin(x) + 2 dx \leq \underbrace{3 \cdot 2\pi}_{6\pi}$$