

Closing today: HW_8A, 8B (8.3, 9.1)

Closing Next Wed: HW_9A, 9B (9.3, 9.4)

Final: Sat, Dec 9th, 1:30-4:20 in KANE 130

9.3: Separable Differential Equations

Entry Task: (Motivation)

Implicitly differentiate $x^2 + y^3 = 8$

and solve for $\frac{dy}{dx}$.

$$\begin{aligned} & \frac{d}{dx} \left[x^2 + y^3 = 8 \right] \\ & \Rightarrow 2x + 3y^2 \frac{dy}{dx} = 0 \\ & \Rightarrow 3y^2 \frac{dy}{dx} = -2x \\ & \Rightarrow \boxed{\frac{dy}{dx} = -\frac{2x}{3y^2}} \end{aligned}$$

Idea: separate and integrate both sides.

Entry Task continued:

Find the *explicit* solution for $\frac{dy}{dx} = \frac{-2x}{3y^2}$
with $y(0) = 2$.

$$\begin{aligned} & \frac{dy}{dx} = -\frac{2x}{3y^2} \\ \Rightarrow & 3y^2 \frac{dy}{dx} = -2x \end{aligned}$$

$$\int 3y^2 dy = \int -2x dx$$

$$y^3 + C_1 = -x^2 + C_2$$

$$\Rightarrow x^2 + y^3 = C_2 - C_1 \leftarrow \text{A constant}$$

$$x^2 + y^3 = C$$

$$y(0) = 2 \Rightarrow 0^2 + 2^3 = C \Rightarrow C = 8$$

$$x^2 + y^3 = 8$$

$$\Rightarrow y^3 = 8 - x^2$$

$$\Rightarrow y = (8 - x^2)^{1/3} \leftarrow \text{Explicit}$$

9.3: Separable Differential Equations

A **separable** differential equation is one that can be written as:

$$\frac{dy}{dx} = f(x)g(y).$$

$$(\text{or } \frac{dy}{dx} = \frac{f(x)}{g(y)} \text{ or } \frac{dy}{dx} = \frac{g(y)}{f(x)}.)$$

Example: Find the explicit solution to

$$\frac{dy}{dx} = \frac{x}{y^4} \text{ with } y(0) = 1.$$

$$y^4 \frac{dy}{dx} = x$$

$$\int y^4 dy = \int x dx$$

$$\frac{1}{5} y^5 = \frac{1}{2} x^2 + C_1$$

$$y^5 = \frac{5}{2} x^2 + 5C_1$$

$$y^5 = \frac{5}{2} x^2 + C_2$$

$$y = \left(\frac{5}{2} x^2 + C_2 \right)^{1/5}$$

General Sol'n

$$y = \left(\frac{5}{2} x^2 + C \right)^{1/5}$$

$$y(0) = 1$$

$$\Rightarrow 1 = \left(\frac{5}{2} (0)^2 + C \right)^{1/5}$$

$$\Rightarrow 1^5 = C \Rightarrow C = 1$$

$$y = \left(\frac{5}{2} x^2 + 1 \right)^{1/5}$$

Q) What is C_1 ?
What is C_2 ?

$$\frac{1}{5} (0)^5 = \frac{1}{2} (0)^2 + C_1 \Rightarrow C_1 = 0$$

$$C_2 + C_1 = 1$$

Example: Find the explicit solution to

$$\frac{dy}{dx} = \frac{x \sin(2x)}{3y} \text{ with } y(0) = -1.$$

$$\int 3y \, dy = \int x \sin(2x) \, dx$$

$u = x \quad dv = \sin(2x) \, dx$
 $du = dx \quad v = -\frac{1}{2} \cos(2x)$

$$\frac{3}{2} y^2 = -\frac{1}{2} x \cos(2x) - \int -\frac{1}{2} \cos(2x) \, dx$$

$$\frac{3}{2} y^2 = -\frac{1}{2} x \cos(2x) + \frac{1}{4} \sin(2x) + C_1$$

$$C_2 = \frac{2}{3} C_1$$

$$y^2 = -\frac{1}{3} x \cos(2x) + \frac{1}{6} \sin(2x) + C_2$$

$$y = \pm \sqrt{-\frac{1}{3} x \cos(2x) + \frac{1}{6} \sin(2x) + C}$$

GENERAL
EXPLICIT SOLN.

$$y(0) = -1 \Rightarrow \textcircled{1} \text{ THIS IS } "-"$$

$$\Rightarrow \textcircled{2} -1 = -\sqrt{0+0+C}$$

$$\Rightarrow C = 1$$

Q] What is
 C_1 ?

$$C_1 = \frac{3}{2}$$

$$y = -\sqrt{-\frac{1}{3} x \cos(2x) + \frac{1}{6} \sin(2x) + 1}$$

Example: Find the explicit solution to

$$(x+1) \frac{dy}{dx} = \frac{x^2}{e^y} \text{ with } y(0) = 0.$$

$$e^y \frac{dy}{dx} = \frac{x^2}{x+1}$$

$$\begin{aligned} x+1 & \frac{x-1}{x^2} \\ & \underline{\underline{(x^2+x)}} \\ & \underline{\underline{-x}} \\ & \underline{\underline{-x=0}} \end{aligned}$$

$$\int e^y dy = \int \frac{x^2}{x+1} dx$$

$$\int e^y dy = \int x-1 + \frac{1}{x+1} dx$$

$$e^y = \frac{1}{2}x^2 - x + \ln|x+1| + C,$$

$$y = \ln\left(\frac{1}{2}x^2 - x + \ln|x+1| + C\right) \text{ general sol'n}$$

$$\begin{aligned} y(0) = 0 & \Rightarrow 0 = \ln(0-0+0+C) \\ & \Rightarrow \boxed{C=1} \end{aligned}$$

$$y = \ln\left(\frac{1}{2}x^2 - x + \ln|x+1| + 1\right)$$

Law of Natural Growth

Assumption: "The rate of growth of a population is proportional to the size of the population."

$P(t)$ = population at year t ,

$\frac{dP}{dt}$ = rate of change of the population

The law of natural growth assumes

$$\frac{dP}{dt} = kP,$$

for some constant k

(we call k the relative growth rate).

Find the explicit solution to

$$\frac{dP}{dt} = kP \text{ with } P(0) = P_0$$

$$\int \frac{1}{P} dP = \int k dt$$

$$\ln|P| = kt + C_1$$

$$\Rightarrow |P| = e^{kt+C_1}$$

$$\Rightarrow |P| = e^{C_1} e^{kt}$$

$$|P| = C_2 e^{kt}$$

$$\Rightarrow P = \pm C_2 e^{kt}$$

$$P(t) = C_3 e^{kt}$$

GENERAL SOLN

$$P(t) = C e^{kt}$$

$$P(0) = P_0 \Rightarrow P_0 = C e^0 \Rightarrow C = P_0$$

$P(t) = P_0 e^{kt}$

WHAT IS C_1 ?

$$\pm e^{C_1} = P_0$$

$$\Rightarrow C_1 = \ln |P_0|$$

Let
 $C_2 = e^{C_1}$

Let
 $C_3 = \pm C_2$

CLEANER THAN
WRITING

$$P(t) = \pm C e^{kt}$$

also correct

1. 500 bacteria are in a dish at $t=0$ hr.
 8000 bacteria are in the dish at $t=3$ hr.

Assume the population grows at a rate proportional to its size.

Find the function, $B(t)$, for the bacteria population with respect to time.

$$\frac{dB}{dt} = kB \quad B(0) = 500$$

USING THE GENERAL SOLN WE
 ALREADY FOUND

$$B(t) = B_0 e^{kt}$$

$$\bullet B(0) = 500 \Rightarrow B_0 = 500$$

$$B(t) = 500e^{kt}$$

$$\bullet B(3) = 8000 \Rightarrow$$

$$500e^{3k} = 8000$$

$$\Rightarrow e^{3k} = 16$$

$$3k = \ln(16) \Rightarrow k = \frac{\ln(16)}{3} \approx 0.924196$$

THUS,

$$\frac{1}{3} \ln(16) +$$

$$B(t) = 500e$$

$$B(t) = 500 (16)^{\frac{t}{3}}$$

NOTE:

$$\frac{t}{3} \ln(16) = \ln(16^{\frac{t}{3}})$$

$$\Rightarrow e^{\frac{t}{3} \ln(16)} = e^{\ln(16^{\frac{t}{3}})} = 16^{\frac{t}{3}}$$

2. The *half-life* of cesium-137 is 30 years. Suppose we start with a 100-mg sample. The mass decays at a rate proportional to its size.

Find the function, $m(t)$, for the mass with respect to time.

$$\frac{dm}{dt} = Km, \quad m(0) = 100$$

$$m(t) = m_0 e^{kt}$$

- $m(0) = 100 \Rightarrow m_0 = 100$

$$m(t) = 100 e^{kt}$$

- $m(30) = 50 \leftarrow \text{HALF!}$

$$50 = 100 e^{30k}$$

$$\Rightarrow \frac{1}{2} = e^{30k}$$

$$\Rightarrow \ln(\frac{1}{2}) = 30k$$

$$\Rightarrow k = \frac{1}{30} \ln(\frac{1}{2})$$

$$\approx -0.023104$$

NEGATIVE!
↓ DECAY!

lose about
2% per
year

$$m(t) = 100 e^{\frac{\ln(\frac{1}{2})}{30} t}$$

$$= 100 e^{\ln((\frac{1}{2})^{\frac{t}{30}})}$$

$$= 100 (\frac{1}{2})^{\frac{t}{30}}$$