

Closing Wed: HW_10A,10B (9.3,9.4)

Final Exam, Saturday, Dec. 9th

Kane 130, 1:30-4:20pm

Assigned seats, for your seat go to:

catalyst.uw.edu/gradebook/aloveles/102715

9.4 Diff. Eq. Apps (*continued*)

Example:

Assume a 12 Liter vat contains 7 kg of salt initially. A pipe pumps in salt water (brine) at 3 L/min with a concentration of 2 kg/L of salt.

The vat is well mixed.

The mixture leaves the vat at 3 L/min.

Let $y(t)$ = the amount of salt in the vat at time t .

(a) Find $y(t)$.

(b) Find the limit of $y(t)$ as $n \rightarrow \infty$.

Here is what these problems typically look like:

V = volume of vat (liters)

t = time (min)

$y(t)$ = amount in vat (kg)

$\frac{dy}{dt}$ = rate (kg/min)

Thus,

$$\begin{aligned} \frac{dy}{dt} &= \text{Rate In} - \text{Rate out} \\ &= \left(? \frac{kg}{L} \right) \left(? \frac{L}{min} \right) - \left(\frac{y}{V} \frac{kg}{L} \right) \left(? \frac{L}{min} \right) \end{aligned}$$

$$y(0) = ? \text{ kg}$$

Example:

Assume a 100 Liter vat contains 5kg of salt initially. Two pipes (A & B) pump in salt water (brine).

Pipe A: Enters at 3L/min with a concentration of 4kg/L of salt.

Pipe B: Enters at 5L/min with a concentration of 2kg/L of salt.

The vat is well mixed.

The mixture leaves the vat at 8L/min.

Let $y(t)$ = the amount of salt in the vat at time t .

(a) Find $y(t)$.

(b) Find the limit of $y(t)$ as $n \rightarrow \infty$.

4. Air Resistance:

A skydiver steps out of a plane that is 4,000 meters high with an initial downward velocity of 0 m/s.

The skydiver has a mass of 60 kg. (Treat downward as positive).

Let $y(t)$ = "height at time t "

Newton's 2nd Law says:

(mass)(acceleration) = Force

$$m \frac{d^2 y}{dt^2} = \text{sum of forces on the object}$$

The force due to gravity has constant magnitude (acting downward):

$$F_g = mg = 60 \cdot 9.8 = 588 \text{ N}$$

One model for air resistance

The force due to air resistance (*drag force*) is proportional to velocity and in the opposite direction of velocity. So

$$F_d = -k v \text{ Newtons}$$

Assume for this problem $k = 12$.

Spring 2011 Final:

$v(t)$ = velocity of an object

$$F = mg - kv$$

Recall:

$$F = ma = m \frac{dv}{dt}$$

You are given m , g , and k and asked for solve for $v(t)$.

Spring 2014 Final:

A lake has a volume of 1000 cubic meters and contains pure water. A farm near the lake begins using a pesticide. Runoff from the farmland into the lake is 10 cubic meters of water per day, with a concentration of 50 grams of pesticide per cubic meter of water. The lake drains to the ocean at a rate of 10 cubic meters per day.

Find the formula $p(t)$ for the amount of pesticide in the lake at time t days.

Winter 2011 Final:

Your friend wins the lottery, and gives you P_0 dollars to help you pay your college expenses. The money is invested in a savings account that earns 10% annual interest, compounded continuously, and you withdraw the money continuously at an average rate of \$3600 per year.

Find the formula $A(t)$ for the amount of money in the account after t years.

Fall 2009 Final:

The swine flu epidemic has been modeled by the Gompertz function, which is a solution of

$$\frac{dy}{dt} = 1.2 y (K - \ln(y)),$$

where $y(t)$ is the number of individuals (in thousands) in a large city that have been infected by time t , and K is a constant.

On July 9, 2009, 75 thousand individuals had been infected.

One month later, 190 thousand individuals had been infected.

Find the formula $y(t)$ for the number of people that are infected t months, July 9, 2009.

Side Note on Population Modeling

The Logistics Equation

Consider a population scenario where there is a limit (capacity) to the size of the population.

Let $P(t)$ = population size at time t .

M = maximum population size.

(capacity)

We sometimes want a model that

- a. ...is like natural growth when $P(t)$ is significantly smaller than M ;
- b. ...levels off (with a slope approaching zero), then the population approaches M .

One such model is the so-called logistics equation

$$\frac{dP}{dt} = kP \left(1 - \frac{P}{M} \right) \text{ with } P(0) = P_0$$