6. (8 points) A bag of sand is lifted from the ground to the top of a 30 foot high building at a constant speed with a cable that weighs 2 lb/ft. A small tear in the bag causes sand to slowly pour out. Initially the bag contains 100 pounds of sand, but the sand leaks out at a constant rate and the bag weighs 90 pounds just as it reaches the 30 foot height. How much work is done?

(Hint: Find a linear equation for the force (weight) of the bag of sand at a given height.)

work = work to lift cable

+ work to lift sandbag

work to lift cable

=
$$\int_{0}^{30} 2x \, dx = x^{2} \Big|_{0}^{30} = \boxed{900}$$

work to lift sandbag

= $\int_{0}^{30} force \, dx$
 $x = 0$ force = 0
 $x = 30$

force = 0
 $x = 30$
 $x = 30$

Note: you could also take x=0 at the bottom in which case the force equation becomes force = - \frac{1}{3} \times + 100. (But you get the same value) 5. (8 pts) After Dr. Loveless dries off, he continues his work out. He starts to lift a sandbag. The sandbag weighs 50 pounds when it is on the ground. As he lifts the bag it leaks out sand at a constant linear rate. When the sandbag is lifted 2 feet, it weighs 46 pounds. Before he passes out, Dr. Loveless does 145 foot-pounds of work in lifting the sandbag. How high did he lift the sandbag?

(Hint: Start by finding the linear function for weight (force) in terms of height.)

$$f(x) = mx + b \qquad m = \frac{46-50}{2-0} - 2$$

$$f(x) = -2x + 50 \quad \text{pounds} \qquad x = \text{dist. from}$$

$$\text{bottom.} \qquad x = 2 \quad \text{if } 46 \text{ lbs}$$

$$\text{only maker sense up to } x = 25$$

$$\text{Work} = \int_{0}^{a} -2x + 50 \, dx \qquad \text{if } 145 \quad \text{ft-lbs} \qquad \text{Fino } a = ?$$

$$-x^{2} + 50x \mid_{0}^{a} ?$$

$$-a^{2} + 50a \qquad = 145$$

$$0 = a^{2} - 50a + 145 \qquad \text{the othersol's list } 46.9063023$$

$$\text{take the ansure list than } 25$$

$$a = \frac{50 - \sqrt{50^{2} + 1.145}}{2} = \frac{50 - \sqrt{1920}}{2} \approx 3.091977 \quad \text{feet}$$

6. (6 points) Consider the region, R, bounded by $y = 4x^2$ and the x-axis between x = 0 and x = 1. Using both methods, cylindrical shells and cross-sectional slicing, set up two integrals for the volume of the solid obtained by rotating the region R about the **vertical line** x = 6. Only set up, DO NOT EVALUATE.

(a) Cross-sectional slicing:

$$\int_{0}^{4} \pi \left(6 - \frac{1}{2}\sqrt{y}\right)^{2} - \pi \cdot 5^{2} dy$$

= 14T = 439823

X=6


(b) Cylindrical Shells:

=
$$5^{\circ}$$
 40 + $50e^{-x/2}$ dx
= $40x + 50(-2)e^{-x/2}$ | 6° = $(40(8) - 100e^{-4}) - (40(6) - 100e^{-4})$
= $320 - 100e^{-4} + 100$
= $420 - 100e^{-4}$ | 5° = $420 - \frac{100}{64}$
= $420 - \frac{100}{64}$ | 5° = 418.168456111 | 5°

5. (10 points) Consider the region R in the first quadrant of the xy-plane bounded by $y=x^2$, y=4 and the y-axis for some positive number a. The water in a full tank is in the shape of the solid obtained by rotating R about the y-axis.

Assume all lengths are in meters, so the tank is 4 meters high. And remember the density of water is 1000 kg/m^3 and gravity is 9.8 m/s^2 .

Set up and evaluate an integral for the work required to pump all the water to the top of the tank and over the edge.



$$WORK = \int_{0}^{4} 9800 \pi (Jy)^{2} (4-y) dy$$

$$= 9800\pi \int_{0}^{4} 4y - y^{2} dy$$

$$= 9800\pi \left[2y^{2} - \frac{1}{3}y^{3}\right]_{0}^{4}$$

$$= 9800\pi \left[24^{2} - \frac{1}{3} + 7\right] - (0)$$

$$= 9800\pi \left[32 - \frac{64}{3}\right]$$

$$= 9800\pi \left[\frac{96}{3} - \frac{64}{3}\right]$$

5. (10 points) The portion of the graph $y = \frac{1}{9}x^2$ between x = 0 and x = 3 is rotated around the y-axis to form a container. The container is full of a liquid that has density 100 lbs/ft³. Find the work required to pump all of the liquid out over the side of the container. (Distance is measured in feet).

$$y = \frac{1}{4}x^{2}$$

$$\Rightarrow qy = x^{2}$$

$$\Rightarrow x = \sqrt{qy}$$

DIST TO = 1-yi
$$y$$
 $y = x^2/9$

BREAKINTO N SUBDIVISIONS
$$\Delta y = \frac{1-0}{n} = \frac{1}{n}$$

$$y = \frac{1}{n}$$

$$= 900\pi \left(\frac{1}{2} - \frac{1}{3}\right)$$

$$= 900\pi \frac{1}{6} = 150\pi + 105$$

 $=900\pi \left(\frac{1}{2}y^2 - \frac{1}{3}y^3\right)$

(a) A tank full of rainwater is outside your dorm. The shape of the tank is described as follows: Consider the region R in the first quadrant of the xy-plane bounded by $y=x^2$, y=1 and the y-axis (lengths are in meters). The full tank is in the shape of the solid obtained by rotating R about the y-axis.

You plan to pump all the water to the top of the tank and over the edge into your bucket.

(b) Once all the water is in your bucket. The bucket is lifted by cable to the roof of your dorm (where you will wait for your instructor to walk by). The cable weighs 5 Newtons per meter and the *empty* bucket weighs 100 Newtons. The top of the building is 20 meters high.

Recall the density of water is 1000 kg/m³ and gravity is 9.8 m/s².

Find the total about of work done in pumping out the water and lifting the full bucket to the roof of your dorm. (Give your final answer as a decimal in Joules).

PUMPING
WATER

FUNCE FOR A

FUNCE FOR A

SLICE

WORK TD

PUMP WATER

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 (1-y) dy = 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$

$$= 100 + 9800 \text{ Tr} (\sqrt{y})^2 dy = 2000 + 9800 \text{ Tr} (\sqrt{y})^2 dy$$