

Closing Wed night: HW_1A, 1B, 1C
Check out the first newsletter for
homework hints, review sheets and
old exam practice.

Entry Tasks:

(a) $f''(x) = 5\sqrt{x} + x$, $f(0) = 3$, $f(1) = 4$
Find $f(x)$.

(b) Ron steps off the 10 meter high
dive at his local pool. Find a
formula for his height above the
water.

(Assume his acceleration is a
constant 9.8 m/s^2 downward)

5.1 Defining Area

Calculus is based on limiting processes that “approach” the exact answer to some rate question.


In Calculus I, you defined

$$\begin{aligned} f'(x) &= \text{'slope of the tangent at } x\text{'} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \end{aligned}$$

In Calculus II, we will see that antiderivatives are related to the area ‘under’ a graph

$$= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$$

Visual:



Example:

Approximate the area under $f(x) = x^3$ from $x = 0$ to $x = 1$ using $n = 3$ subdivisions and *right-endpoints* to find the height.

Riemann sums set up:

We are going to build a procedure to get better and better approximations of the area “under” $f(x)$.

1. Break into n equal subintervals.

$$\Delta x = \frac{b-a}{n} \text{ and } x_i = a + i\Delta x$$

2. Draw n rectangles; use function.

Area of each rectangle =

$$(\text{height})(\text{width}) = f(x_i^*)\Delta x$$

3. Add up rectangle areas.

I did this again with 100 subdivisions, then 1000, then 10000. Here is the summary of my findings:

n	R_n	L_n
4	0.390625	0.140625
5	0.36	0.16
10	0.3025	0.2025
100	0.255025	0.245025
1000	0.25050025	0.24950025
10000	0.2499500025	0.2500500025

Right-Endpoint Pattern:

$$\Delta x = \frac{1-0}{n} = \frac{1}{n}, \quad x_i = 0 + i\frac{1}{n} = \frac{i}{n}$$

$$\text{Height of rect.} = f(x_i) = x_i^3 = \left(\frac{i}{n}\right)^3$$

$$\text{Area} = f(x_i)\Delta x = x_i^3\Delta x = \left(\frac{i}{n}\right)^3 \frac{1}{n}$$

$$\text{Sum} = \sum_{i=1}^n x_i^3 \Delta x = \sum_{i=1}^n \left(\frac{i}{n}\right)^3 \frac{1}{n}$$

$$\text{Exact Area} = \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(\frac{i}{n}\right)^3 \frac{1}{n}$$

Definition of the Definite Integral

We define the exact area “under” $f(x)$ from $x = a$ to $x = b$ curve to be

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x,$$

where $\Delta x = \frac{b-a}{n}$ and

$$x_i = a + i\Delta x.$$

We call this the definite integral of $f(x)$ from $x = a$ to $x = b$, and we write

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$