

Close Wed: HW_5A, 5B (7.1, 7.2)

Close Fri: HW_5C (7.3)

Office Hours: 1:30-3:30 in THO 335

See posting of several more integration by parts examples (with solutions).

Entry Task: Evaluate

$$\int_0^1 \sin^{-1}(x) dx = \int_0^1 \arcsin(x) dx$$

7.1 Integration by Parts (continued)

Summary: $\int u dv = uv - \int v du$

1. Pick $u = ??$. The rest is dv .
2. Differentiate to get du
Integrate to get v .

Here are all the most common examples:

- a) Products: $xe^x, x^2 \cos(3x), x \sin(5x)$
- b) Logarithms: $\ln(x), x^{10} \ln(x), \dots$
- c) Inv. Trig: $\sin^{-1}(x), \tan^{-1}(x), \dots$
- d) Products: $e^x \sin(x), e^x \cos(x)$

Example: (Never ending integration by parts and how to end it):

$$\int e^x \cos(x) dx$$

7.2 Trigonometric Integral Methods

Goal: Build up rules for integrating combinations of trig functions.

Basic motivating examples:

All of these could be done with substitution: What is u ?

$$\int \sin^4(x) \cos(x) dx$$

$$\int \sin(x) \cos^3(x) dx$$

$$\int \tan^5(x) \sec^2(x) dx$$

$$\int \sec^6(x) \sec(x) \tan(x) dx$$

Idea: Use trig identities to turn a problem into a substitution problem like those above.

Tools

Essential Trig Identities:

$$\tan(x) = \frac{\sin(x)}{\cos(x)}, \cot(x) = \frac{\cos(x)}{\sin(x)},$$
$$\sec(x) = \frac{1}{\cos(x)}, \csc(x) = \frac{1}{\sin(x)}.$$

$$\sin^2(x) + \cos^2(x) = 1$$
$$\tan^2(x) + 1 = \sec^2(x)$$

$$\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$$

$$\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$$

$$\sin(x) \cos(x) = \frac{1}{2} \sin(2x)$$

See my online posting (or Appendix D of your book) for a more general discussion of trig identities.

Add these to your table of integrals you know (an updated table is already on the website):

$$\int \tan(x) dx = \ln |\sec(x)| + C$$

$$\int \sec(x) dx = \ln |\sec(x) + \tan(x)| + C$$

$$\int \sec^3(x) dx =$$
$$\frac{1}{2} \sec(x) \tan(x) + \frac{1}{2} \ln |\sec(x) + \tan(x)| + C$$