

Close Today: HW_5A, 5B (7.1,7.2)

Close Mon: HW_5C (7.3)

Close Wed: HW_6A (7.4)

Close next Fri: HW_6B, 6C (7.5, 7.7)

Office Hours today: 1:30-3:30pm, Com B-006

Entry Tasks: Fill in the blanks

Square Identities
$\sin^2(x) =$
$\cos^2(x) =$
$\sec^2(x) =$
$\tan^2(x) =$
Half Angle Identities
$\sin^2(x) =$
$\cos^2(x) =$
$\sin(x)\cos(x) =$


What are these in terms of a, b, c?

$\sin(\theta) =$

$\cos(\theta) =$

$\tan(\theta) =$

$\sec(\theta) =$



What is the first step in each below?

$$\int \sin^3(x)\cos^4(x)dx$$

$$\int \sin^5(x)\cos^3(x)dx$$

$$\int \cos^4(x)dx$$

$$\int \tan^5(x)\sec^4(x)dx$$

$$\int \tan^5(x)\sec(x)dx$$

7.3 Trigonometric Substitution

Goal: Evaluation method for integrals

with $\sqrt{a^2 - x^2}$, $\sqrt{a^2 + x^2}$, or $\sqrt{x^2 - a^2}$

3 Types of Problems/Motivation

1. $a^2 - x^2$, *example:*

$$\int \frac{x^3}{\sqrt{4 - x^2}} dx$$

2. $a^2 + x^2$, *example:*

$$\int x^2 \sqrt{9 + x^2} dx$$

3. $x^2 - a^2$, *example:*

$$\int \frac{\sqrt{x^2 - 16}}{x} dx$$

CASE	SUBSTITUTION
$a^2 - x^2$	$x = a \sin(\theta), \quad -\pi/2 \leq \theta \leq \pi/2$
$a^2 + x^2$	$x = a \tan(\theta), \quad -\pi/2 < \theta < \pi/2$
$x^2 - a^2$	$x = a \sec(\theta), \quad 0 \leq \theta < \pi/2, \text{ (positive } x)$ $\pi \leq \theta < 3\pi/2 \text{ (negative } x)$

Trigonometric Substitution Method:

- A) Make the substitution.
 Don't forget $dx = ??? d\theta$.
 Use trig identity (eliminate root)
- B) Use 7.2 methods.
- C) Draw a triangle and return to x .

Completing the Square:

$$\sqrt{ax^2 + bx + c}$$

If you ever encounter a “**middle term**” (like **bx** above), then you need to complete the square.

Example: $\sqrt{64 - 24x - 4x^2}$

i) Factor out the “a”.

$$\sqrt{4(16 - 6x - x^2)} = 2\sqrt{16 - 6x - x^2}$$

ii) Add and subtract “half the middle squared”

$$\text{Half of middle} = (-6)/2 = -3$$

$$\text{Squared} = (-3)^2 = 9$$

$$2\sqrt{16 + 9 - 9 - 6x - x^2}$$

iii) Factor the perfect square

$$2\sqrt{25 - (x + 3)^2}$$

iv) Check your work!