

Close Wed: HW_5A,5B,5C (6.5,7.1,7.2)

Next Wed: HW_6A,6B,6C (7.3,7.4,7.5)

Work ahead!!! Be done with 7.2 today and start to look at 7.3.

7.2 Trig Integrals (continued)

Entry Task: Fill in the blanks

Square Identities	
$\sin^2(x)$	$= 1 -$
$\cos^2(x)$	$= 1 -$
$\tan^2(x)$	$=$
$\sec^2(x)$	$=$
Half Angle Identities	
$\sin^2(x)$	$= \frac{1}{2}$
$\cos^2(x)$	$= \frac{1}{2}$
$\sin(x) \cos(x)$	$= \frac{1}{2}$


What are these in terms of a, b, and c?

$$\sin(\theta) =$$

$$\cos(\theta) =$$

$$\tan(\theta) =$$

$$\sec(\theta) =$$



Integrate

$$\int \sin^3(x) \cos^2(x) dx$$

Integrals involving $\sin(x)$ and $\cos(x)$

Case 1 ($\cos(x)$ or $\sin(x)$ has odd power)

- Separate one from the odd power.
(i.e. pull out one $\sin(x)$ or $\cos(x)$)
- Use $\sin^2(x) = 1 - \cos^2(x)$
 $\cos^2(x) = 1 - \sin^2(x)$
(get rest in term of the other)
- Use u-substitution.

$$\int \sin^3(x)\cos^5(x)dx$$

Case 2 (both $\sin(x)$, $\cos(x)$ even powers)

- Use

$$\sin^2(x) = \frac{1}{2}(1 - \cos(2x))$$

$$\cos^2(x) = \frac{1}{2}(1 + \cos(2x))$$

over and over again until you can integrate.

Example:

$$\int \cos^4(x) dx$$

Integrals involving $\tan(x)$ and $\sec(x)$

Case 3 ($\sec(x)$ has an even power)

- Separate out $\sec^2(x)$
- Use $\sec^2(x) = 1 + \tan^2(x)$
(get rest in terms of $\tan(x)$)
- Use $u = \tan(x)$

Example:

$$\int \tan^5(x) \sec^4(x) dx$$

Case 4 ($\tan(x)$ has an odd power)

- Separate out $\sec(x) \tan(x)$
- Use $\tan^2(x) = \sec^2(x) - 1$
(get rest in terms of $\sec(x)$)
- Use $u = \sec(x)$

Example:

$$\int \tan^3(x) \sec(x) dx$$

Tried the four cases and still stuck?

Then here are things to try:

1. Rewrite in terms of $\sin(x)$ and $\cos(x)$.
2. Rewrite in terms of $\sec(x)$ and $\tan(x)$.
3. Try using trig identities.

There are still a few “holes”.

Namely, when there is one $\tan(x)$ or an odd power on $\sec(x)$. You can quote these (proof in the book):

$$\int \tan(x) dx = \ln|\sec(x)| + C$$

$$\int \sec(x) dx = \ln|\sec(x) + \tan(x)| + C$$

$$\int \sec^3(x) dx = \frac{1}{2} \sec(x) \tan(x) + \frac{1}{2} \ln|\sec(x) + \tan(x)| + C$$

7.3 Trigonometric Substitution

Goal: Develop a method to evaluate integrals involving expressions of the form $\sqrt{a^2 - x^2}$, $\sqrt{a^2 + x^2}$, or $\sqrt{x^2 - a^2}$

CASE	SUBSTITUTION
$a^2 - x^2$	$x = a \sin(\theta), \quad -\pi/2 \leq \theta \leq \pi/2$
$a^2 + x^2$	$x = a \tan(\theta), \quad -\pi/2 < \theta < \pi/2$
$x^2 - a^2$	$x = a \sec(\theta), \quad 0 \leq \theta < \pi/2,$ $\pi \leq \theta < 3\pi/2$

Example:

1.
$$\int \frac{x^3}{\sqrt{4 - x^2}} dx$$

Trigonometric Substitution Method:

- A) Substitute, don't forget $dx = ??d\theta$.
Simplify (eliminate root)
- B) Use 7.2 methods for trig integrals.
- C) Draw a triangle and return to x .

$$2. \int \sqrt{9 + x^2} dx$$

$$3. \int \frac{\sqrt{x^2 - 16}}{x} dx$$

Completing the Square:

$$\sqrt{ax^2 + bx + c}$$

If you encounter a “**middle term**” (like **bx** above), then complete the square.

Example: $\sqrt{64 - 24x - 4x^2}$

i) Factor out the “a”.

$$\sqrt{4(16 - 6x - x^2)} = 2\sqrt{16 - 6x - x^2}$$

ii) Add/subtract “half-middle squared”

$$\text{Half of middle} = (-6)/2 = -3$$

$$\text{Squared} = (-3)^2 = 9$$

$$2\sqrt{16 + 9 - 9 - 6x - x^2}$$

iii) Factor the perfect square

$$2\sqrt{25 - (x + 3)^2}$$

iv) Check your work!!!!

Example:

$$\int \frac{x}{\sqrt{64 - 24x - 4x^2}} dx$$