

Closing *Wed*: HW_7A,7B (7.5, 7.7, 7.8)

Note: Exam 2 is **Thursday!!!**

Covers 6.4, 6.5, 7.1-7.5, 7.7, 7.8

The exam will roughly look like this:

First 3 pages: 6 integrals (*ALL* types)

4th page: 6.5, 7.7 and/or 7.8

5th page: 6.4

(8.1 Arc Length is NOT on our midterm)

A Brief Exam 2 Review

Integration!

Work.

Average Value.

Trapezoid, Simpson Rules.

Improper Integrals.

1. The 4 special methods

By parts:

$$xe^{3x}, x^2 \cos(5x), \\ \frac{\ln(x-1)}{x^2}, x \tan^{-1}(x).$$

Trig:

$$\sin^3(x) \cos(x), \cos^4(x), \\ \sec^3(2x) \tan^3(2x)$$

Trig sub:

$$\frac{1}{x\sqrt{x^2-9}}, \frac{1}{(4-x^2)^{3/2}}, \\ \sqrt{x^2 + 6x + 10}.$$

Part Frac:

$$\frac{x+2}{(x-1)(x-3)}, \frac{4x}{(x-1)^2(x-3)}, \\ \frac{5}{x(x^2+4)}, \frac{x^2}{x+7}, \frac{x-3}{x^2+8x+20}$$

2. Substitution and Simplifying

Try $u = \sqrt{x}$, $u = \text{inside}$, $u = e^x$, $u = \text{trig}$.

Completing the square.

Trig facts.

Square identities, half-angle.

Triangle trick.

Random Integrals from Old Finals:

$$1. \int \frac{1-x}{\sqrt{1-x^2}} dx$$

$$2. \int \frac{x^2 - x + 8}{x^3 + 4x} dx$$

$$3. \int 2x \ln(x+5) dx$$

$$4. \int \cos^3(x) dx$$

$$5. \int_0^2 \frac{1}{\sqrt{x^2 + 2x + 4}} dx$$

$$6. \int_1^3 \frac{1}{x^2 + x^3} dx$$

$$7. \int \tan^2 x \sec^4(x) dx$$

$$8. \int \frac{1}{(1 + \sqrt{x})^3} dx$$

$$9. \int \sin(x) \sqrt{\cos(x)} dx$$

3. Improper Integrals:

- a) Rewrite as a limit!!
- b) Integrate
- c) Take limit

Random Improper Integrals:

1. $\int_1^2 \frac{x}{\sqrt{x-1}} dx$
2. $\int_{-3}^{\infty} x e^{-x} dx$
3. $\int_1^{\infty} \frac{1}{\sqrt{x}(1+x)} dx$

4. Trapezoid/Simpson Rules

- a) Set up integral, then compute width and label tickmarks.
- b) Use formula.

Approximation Example:

1. Use Simpson's Method with $n = 4$ subdivision to approximate the value of

$$\int_0^4 \sqrt{1+4x^4} dx$$

5. New Applications

a) **Average value** = $\frac{1}{b-a} \int_a^b f(x) dx$

b) **Work** = $\int_a^b (Force)(Dist)$

Step 1: Draw picture (start and end)

Label clearly.

Draw a typical subdivision.

Step 2: Find pattern for *Force* and *Dist*.

Step 3: Integrate.

Type 1: “Changing force”

Force changing as object is moved
(leaky bucket, springs, given force).

$f(x)$ = “force formula at x ”

$Force = f(x)$, $Dist = \Delta x$;

$$Work = \int_a^b f(x) dx$$

Type 2: “Stack of books”

(chain, pumping)

Chain/Cable:

Given $k = \text{density} = \text{force/length}$

if $x = 0$ is labeled at the top.

then for any subdivision that makes
it to the top:

Force = $k \Delta x$, *Dist* = x

$$\text{Work} = \int_a^b k x dx$$

Pumping:

Given $k = \text{density} = \text{force/volume}$

if top is $y = b$, then

Force = $k(\text{Area})\Delta y$, *Dist* = $b - y$;

$$\text{Work} = \int_a^b k(\text{Area})(b - y)dy$$

You need to have an basic understanding of
how we label and find pattern to be able to
adapt to changes in these problems!!

Applications from old tests:

1. Find the average value of $\cos^3(x)$ on the interval 0 to $\pi/2$.

3. A 1600 lb elevator is suspended by a 200 ft cable that weighs 10 lb/ft. How much work is required to raise the elevator from the basement to the third floor, a distance of 30 ft?

4. A rope is used to pull a bucket full of water up from a well that is 10 m deep. The rope has a total mass of 5 kg. The bucket has a mass of 11 kg. Find the total work done in lifting the bucket to the top (Recall: Accel. due to gravity is 9.8 m/s^2)

5. A well is in the shape of a cylinder of radius 1 meter and depth 8 meters. It is half full of water. Find the work to pump all the water to the top.
(Recall: Water weighs 9800 N/m^3)

6. The portion of the graph $y = x^2 / 9$ between $x = 0$ and $x = 3$ is rotated about the y -axis to form a container. The container is full of a liquid that has density 100 lbs/ft^3 . Find the work required to pump all the liquid to the top of the container.