

1. (13 pts)

(a) Find a vector that has length 7 and is orthogonal to both  $\mathbf{u} = \langle 1, 0, 2 \rangle$  and  $\mathbf{v} = \langle 3, -2, 1 \rangle$

$$\vec{u} \times \vec{v} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & 0 & 2 \\ 3 & -2 & 1 \end{vmatrix} = (0 - -4)\vec{i} - (1 - 6)\vec{j} + (-2 - 0)\vec{k} = \langle 4, 5, -2 \rangle$$

CHECK:  $4 + 0 - 2 = 0 \checkmark$   
 $12 - 10 - 2 = 0 \checkmark$

$$|\vec{u} \times \vec{v}| = \sqrt{16 + 25 + 4} = \sqrt{45} = 3\sqrt{5}$$

$$\boxed{\frac{7}{\sqrt{45}} \langle 4, 5, -2 \rangle}$$

OR


$$\boxed{\frac{-7}{\sqrt{45}} \langle 4, 5, -2 \rangle}$$

(b) Find the distance from point A to point B in the picture below (Hint: Use vector tools!)

$$\vec{AC} = \langle 3, 0, 2 \rangle$$

$$\vec{AB} = \langle 2, 1, -2 \rangle$$

$$\text{COMP}_{\vec{AB}} \vec{AC} = \frac{6 + 0 - 4}{\sqrt{4 + 1 + 4}} = \boxed{\frac{2}{3}}$$



(c) Consider the line through the points  $(0, 0, 1)$  and  $(3, 4, 5)$ . Find the  $(x, y, z)$  point(s) where the line intersects the cylinder  $x^2 + y^2 = 4$ .

$$\text{LINE: } x = 0 + 3t, y = 0 + 4t, z = 1 + 4t$$

$$\text{INTERSECTION: } (3t)^2 + (4t)^2 = 4 \Rightarrow 25t^2 = 4 \Rightarrow t^2 = \frac{4}{25}$$

$$\Rightarrow t = \pm \frac{2}{5}$$

$$t = -\frac{2}{5} \Rightarrow (x, y, z) = \left(-\frac{6}{5}, -\frac{8}{5}, 1 - \frac{8}{5}\right) = \left(-\frac{6}{5}, -\frac{8}{5}, -\frac{3}{5}\right)$$

$$t = \frac{2}{5} \Rightarrow (x, y, z) = \left(\frac{6}{5}, \frac{8}{5}, 1 + \frac{8}{5}\right) = \left(\frac{6}{5}, \frac{8}{5}, \frac{13}{5}\right)$$

2. (12 pts)

- (a) Find parametric equations for the line of intersection of the planes  $x + y + z = 10$  and  $x - 3y - 4z = -10$ .

TWO POINTS?

COMBINING  $\Rightarrow 4y + 5z = 20$   $z = 10 - x - y$

$y = 0 \Rightarrow z = 4 \Rightarrow x = 6$      $P(6, 0, 4)$     CHECK  $\checkmark$

$z = 0 \Rightarrow y = 5 \Rightarrow x = 5$      $Q(5, 5, 0)$

$$\begin{aligned} x &= 6 - t \\ y &= 0 + 5t \\ z &= 4 - 4t \end{aligned}$$

ANY  
POINT ON  
LINE

ANY VECTOR  
PARALLEL TO  
 $\langle -1, 5, -4 \rangle$

- (b) Consider the plane that passes thru  $(4, 4, 2)$  and contains the line  $x = 5t, y = 3 + t, z = 4 - t$ . Find the  $(x, y, z)$  point where this plane intersects the  $y$ -axis.

THREE POINTS?

$P(0, 3, 4)$  ,  $Q(5, 4, 3)$  ,  $R(4, 4, 2)$

$t=0$                        $t=1$

TWO VECTORS  
PARALLEL TO  
DESIREZ PLANE

$\left\{ \begin{aligned} \vec{PQ} &= \langle 5, 1, -1 \rangle = \text{SAME AS DIRECTION VECTOR FOR LINE} \\ \vec{PR} &= \langle 4, 1, -2 \rangle \end{aligned} \right.$

$\vec{PQ} \times \vec{PR} = (-2 - -1)\vec{i} - (-10 - -4)\vec{j} + (5 - 4)\vec{k}$   
 $= \langle -1, 6, 1 \rangle$

CHECK:  $-5 + 6 - 1 = 0 \checkmark$   
 $-4 + 6 - 2 = 0 \checkmark$

PLANE:  $-(x-4) + 6(y-4) + (z-2) = 0$

$y$ -AXIS  $\Leftrightarrow x=0$  AND  $z=0 \Leftrightarrow 4 + 6(y-4) - 2 = 0$

$6(y-4) = -2$

$y-4 = -\frac{1}{3}$

$y = 4 - \frac{1}{3} = \frac{11}{3} = 3.\bar{6}$

$(0, \frac{11}{3}, 0) = (0, 3.\bar{6}, 0)$

3. (12 pts)

(a) Give the precise 3D name for  $4x^2 = 5y^2 + z$ .  $z = 4x^2 - 5y^2$

HYPERBOLIC PARABOLOID

(b) Set up, but DO NOT EVALUATE, an integral that represents the arc length of the curve of intersection of the cylinder  $x^2 + y^2 = 25$  and  $x + 2y + z = 10$ .

$$\begin{aligned} x &= 5 \cos(t) & z &= 10 - x - 2y \\ y &= 5 \sin(t) \\ z &= 10 - 5 \cos(t) - 10 \sin(t) \end{aligned} \quad \left. \vphantom{\begin{aligned} x &= 5 \cos(t) \\ y &= 5 \sin(t) \\ z &= 10 - 5 \cos(t) - 10 \sin(t) \end{aligned}} \right\} \begin{array}{l} \text{MANY OTHER} \\ \text{PARAMETERIZATIONS} \\ \text{ARE VALID} \end{array}$$

$$\int_0^{2\pi} \sqrt{(-5 \sin(t))^2 + (5 \cos(t))^2 + (5 \sin(t) - 10 \cos(t))^2} dt$$

$$= \int_0^{2\pi} \sqrt{25 + (5 \sin(t) - 10 \cos(t))^2} dt$$

(c) Consider the curves  $r_1(t) = \langle 2t, 3t^2, t^3 \rangle$  and  $r_2(u) = \langle 2-2u, 3+3u, u^2+1 \rangle$ . The curves have one point of intersection. Find the angle of intersection to the nearest degree.

Ⓐ  $2t = 2 - 2u \Rightarrow t = 1 - u$

Ⓑ  $3t^2 = 3 + 3u \Rightarrow t^2 = 1 + u \Rightarrow (1-u)^2 = 1 + u$   
 $1 - 2u + u^2 = 1 + u$

Ⓒ  $u = 0, t = 1 \Rightarrow$  (i)  $t^3 = 1 \leftarrow \text{YES!}$   
 $u^2 + 1 = 1$

$u^2 - 3u = 0$   
 $u(u-3) = 0$

$u = 0$  or  $u = 3$   
 $\downarrow \qquad \downarrow$   
 $t = 1 \qquad t = -2$

$u = 3, t = -2 \Rightarrow$  (ii)  $t^3 = -8 \leftarrow \text{NO}$   
 $u^2 + 1 = 10$

$r_1'(t) = \langle 2, 6t, 3t^2 \rangle \quad r_1'(1) = \langle 2, 6, 3 \rangle$   
 $r_2'(u) = \langle -2, 3, 2u \rangle \quad r_2'(0) = \langle -2, 3, 0 \rangle$

$\cos(\theta) = \frac{u \cdot v}{|u||v|} = \frac{-4 + 18 + 0}{\sqrt{4 + 36 + 9} \sqrt{4 + 9 + 0}} = \frac{14}{7\sqrt{13}} = \frac{2}{\sqrt{13}}$

$\theta = \cos^{-1}\left(\frac{2}{\sqrt{13}}\right) \approx 56.31^\circ \quad \boxed{56^\circ}$

4. (13 pts)

(a) Give parametric equations for the tangent line to  $\mathbf{p}(t) = \langle t^2, 3 - 3t, 3 + 2t \rangle$  at  $t = 1$ .

$$\vec{p}'(t) = \langle 2t, -3, 2 \rangle$$

$$\vec{p}(1) = \langle 1, 0, 5 \rangle$$

$$\vec{p}'(1) = \langle 2, -3, 2 \rangle$$

$$\begin{cases} x = 1 + 2u \\ y = 0 - 3u \\ z = 5 + 2u \end{cases}$$

(b) Find the principal unit normal vector  $\mathbf{N}(t)$  for  $\mathbf{q}(t) = \langle 3t, \cos(4t), \sin(4t) \rangle$ .

$$\vec{q}'(t) = \langle 3, -4\sin(4t), 4\cos(4t) \rangle$$

$$|\vec{q}'(t)| = \sqrt{9 + 16\sin^2(4t) + 16\cos^2(4t)} = \sqrt{9 + 16} = 5$$

$$\vec{T}(t) = \left\langle \frac{3}{5}, -\frac{4}{5}\sin(4t), \frac{4}{5}\cos(4t) \right\rangle$$

$$\vec{T}'(t) = \left\langle 0, -\frac{16}{5}\cos(4t), -\frac{16}{5}\sin(4t) \right\rangle$$

$$|\vec{T}'(t)| = \sqrt{\left(\frac{16}{5}\right)^2 \cos^2(4t) + \left(\frac{16}{5}\right)^2 \sin^2(4t)} = \frac{16}{5}$$

$$\vec{N}(t) = \langle 0, -\cos(4t), -\sin(4t) \rangle$$

(c) An object is moving such that its velocity is given by  $\mathbf{r}'(t) = \langle t, \sin(t), t \cos(t) \rangle$  and its initial location is  $\mathbf{r}(0) = \langle 0, 0, 1 \rangle$ . Find the position function  $\mathbf{r}(t)$ .

$$\vec{r}(t) = \int \langle t, \sin(t), t \cos(t) \rangle dt$$

$$= \left\langle \frac{1}{2}t^2 + c_1, -\cos(t) + c_2, \right\rangle$$

$$\int t \cos(t) dt = t \sin(t) - \int \sin(t) dt = t \sin(t) + \cos(t) + c_3$$

$$\begin{aligned} u = t & \quad dv = \cos(t) \\ du = dt & \quad v = \sin(t) \end{aligned}$$

$$\vec{r}(0) = \langle 0, 0, 1 \rangle = \langle c_1, -1 + c_2, 1 + c_3 \rangle \Rightarrow \begin{cases} c_1 = 0 \\ c_2 = 1 \\ c_3 = 0 \end{cases}$$

$$\vec{r}(t) = \left\langle \frac{1}{2}t^2, -\cos(t) + 1, t \sin(t) + \cos(t) \right\rangle$$