

ON q -DERANGEMENT NUMBERS

MICHELLE L. WACHS

(Communicated by Thomas Brylawski)

ABSTRACT. We derive a q -analogue of the classical formula for the number of derangements of an n element set. Our derivation is entirely analogous to the classical derivation, but relies on a descent set preserving bijection between the set of permutations with a given derangement part and the set of shuffles of two permutations.

A classical application of binomial inversion (more generally the principle of inclusion–exclusion) is the derivation of the formula for the number of derangements (permutations with no fixed points) of an n element set:

$$d_n = n! \sum_{k=0}^n \frac{(-1)^k}{k!}.$$

This is obtained by counting permutations according to their number of fixed points and then inverting the resulting equation.

In this note we shall derive a formula of I. Gessel [G] for q -counting derangements by the major index statistic in a way entirely analogous to the classical $q = 1$ case. That is, we shall q -count permutations with k fixed points and then use Gauss inversion (q -binomial inversion or more generally Möbius inversion on the lattice of subspaces of a vector space) to derive the following formula for q -derangement numbers:

$$d_n(q) = [n]! \sum_{k=0}^n \frac{(-1)^k}{[k]!} q^{\binom{k}{2}}.$$

A key step in our derivation and an interesting result in its own right is a descent-preserving bijection between the set of permutations with a given derangement part and the set of shuffles of two permutations. This bijection enables us to use a formula of A. Garsia and I. Gessel for q -counting shuffles.

Gessel [G] obtained the formula for q -derangement numbers as a corollary of an Eulerian generating function formula for counting permutations by descents, major index, and cycle structure, which is proved via a correspondence

Received by the editors February 6, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 05A30; Secondary 05A19, 05A15, 05A05.

Research partially supported by NSF grant DMS:8503700.

between partitions and permutations. q -Derangement numbers have also been interpreted combinatorially on sets of permutations bijectively associated with derangements by A. Garsia and J. Remmel [GR] using the inversion index statistic and by J. Désarménien [D₂] (see [D₁] and [DW]) using the major index and inversion index statistics.

We shall briefly review some permutation statistic notation and terminology. For each integer $n \geq 1$, let $[n]$ denote the polynomial $1 + q + q^2 + \dots + q^{n-1}$ and let $[n]!$ denote the polynomial $[n][n-1] \cdots [1]$. Also $[0]!$ is taken to be 1. The q -binomial coefficients are given by

$$\begin{bmatrix} n \\ k \end{bmatrix} = \frac{[n]!}{[k]![n-k]!}$$

for $0 \leq k \leq n$.

For any positive integer n , let $\langle n \rangle$ denote the set $\{1, 2, \dots, n\}$. We shall think of permutations in the symmetric group \mathcal{S}_n as words with n distinct letters in $\langle n \rangle$. More generally, for a set A of n positive integers, \mathcal{S}_A denotes the set of permutations of A or words with n distinct letters in A . The *descent set* of a permutation $\sigma = \sigma_1, \sigma_2, \dots, \sigma_n$ is $\text{des}(\sigma) = \{i \in \langle n-1 \rangle \mid \sigma_i > \sigma_{i+1}\}$. The *major index* of σ is $\text{maj}(\sigma) = \sum_{i \in \text{des}(\sigma)} i$. Let us recall MacMahon's [M] formula for maj- q -counting permutations in \mathcal{S}_n :

$$\sum_{\sigma \in \mathcal{S}_n} q^{\text{maj}(\sigma)} = [n]!$$

A letter $i \in A$ is said to be a fixed point of $\sigma \in \mathcal{S}_A$ if $\sigma(i) = i$. A permutation with no fixed points is called a *derangement*. Let D_n denote the set of all derangements in \mathcal{S}_n . The q -derangement numbers are defined by

$$d_n(q) = \sum_{\sigma \in D_n} q^{\text{maj}(\sigma)}.$$

It will be convenient to view the empty word Λ as a derangement and to define D_0 to be the set $\{\Lambda\}$. We also let $\text{maj}(\Lambda) = 0$ and $d_0(q) = 1$.

For any permutation $\alpha \in \mathcal{S}_A$, where $A = \{a_1 < a_2 < \dots < a_k\}$, define the *reduction* of α to be the permutation in \mathcal{S}_k obtained from α by replacing each letter a_i by i , $i = 1, 2, \dots, k$. The *derangement part* of a permutation $\sigma \in \mathcal{S}_n$, denoted $dp(\sigma)$, is the reduction of the subword of nonfixed points of σ . For example, $dp(5, 3, 1, 4, 7, 6, 2) =$ reduction of $5, 3, 1, 7, 2 = 4, 3, 1, 5, 2$. We shall use the convention that the derangement part of the identity permutation is the empty word Λ . Note that the derangement part of a permutation is a derangement, and that conversely, any derangement in D_k and k element subset of $\langle n \rangle$ determines a permutation in \mathcal{S}_n with $n - k$ fixed points. Hence, the number of permutations in \mathcal{S}_n with a given derangement part in D_k is $\binom{n}{k}$. Our goal is to q -count permutations with a given derangement part.

Let $\alpha \in D_k$. There is an obvious bijection between the set $\{\sigma \in \mathcal{S}_n \mid dp(\sigma) = \alpha\}$ and the set $\text{Sh}(\alpha, \beta)$ of all *shuffles* of α and $\beta = k + 1, k + 2, \dots, n$,

i.e. permutations in \mathcal{S}_n which contain α and β as complementary subwords. Indeed, for each permutation σ in the former set, replace the subword of non-fixed points of σ by α and the complementary subword of fixed points by β . A very useful result of Garsia and Gessel [GG, Theorem 3.1] allows us to q -count the latter set. Unfortunately, since the above-mentioned bijection does not preserve the major index, it does not help us in q -counting the former set. However, we shall show that there is another bijection between these two sets of permutations which, in fact, preserves descent sets.

Define a letter σ_i of $\sigma = \sigma_1, \sigma_2, \dots, \sigma_n \in \mathcal{S}_n$ to be an *excedant* of σ if $\sigma_i > i$ and a *subcedant* of σ if $\sigma_i < i$. Let $s(\sigma)$ and $e(\sigma)$ be the number of subcedants and excedants, respectively, of σ . We now fix n and let $k \leq n$. For $\sigma \in \mathcal{S}_k$, let $\tilde{\sigma}$ be the permutation of k letters obtained from σ by replacing its i th smallest subcedant by i , $i = 1, 2, \dots, s(\sigma)$, its i th smallest fixed point by $s(\sigma) + i$, $i = 1, 2, \dots, k - s(\sigma) - e(\sigma)$, and its i th largest excedant by $n - i + 1$, $i = 1, 2, \dots, e(\sigma)$. Note that $\tilde{\sigma}$ depends on n as well as σ . For example, if $\sigma = \overline{326541}$ (with subcedants underlined and excedants overlined) and $n = 8$ then $\tilde{\sigma} = 638721$. If $k = n$ then $\tilde{\sigma} \in \mathcal{S}_n$. If σ is a derangement then $\tilde{\sigma} \in \mathcal{S}_A$, where $A = \{1, 2, \dots, s(\sigma)\} \cup \{n - e(\sigma) + 1, n - e(\sigma) + 2, \dots, n\}$.

Lemma 1. *Let $\sigma \in \mathcal{S}_k$, $k \leq n$. Then $\text{des}(\sigma) = \text{des}(\tilde{\sigma})$.*

Proof. Suppose $\sigma = \sigma_1, \sigma_2, \dots, \sigma_k$ and $\tilde{\sigma} = \tilde{\sigma}_1, \tilde{\sigma}_2, \dots, \tilde{\sigma}_k$. For each $i \in \langle k - 1 \rangle$, we shall show $i \in \text{des}(\sigma)$ if and only if $i \in \text{des}(\tilde{\sigma})$, by considering the nine possible designations of subcedant (s), excedant (e), and fixed point (f) to σ_i and σ_{i+1} . First note that if σ_i is a subcedant of σ then $\tilde{\sigma}_i \leq \sigma_i$ and if σ_i is an excedant of σ then $\tilde{\sigma}_i \geq \sigma_i$.

Cases 1–3. Suppose (σ_i, σ_{i+1}) is an (s, s) , (e, e) , or (f, f) pair. It is then clear that $\sigma_i < \sigma_{i+1}$ if and only if $\tilde{\sigma}_i < \tilde{\sigma}_{i+1}$.

Case 4. Suppose (σ_i, σ_{i+1}) is a (s, e) pair. Then we have

$$\tilde{\sigma}_i \leq \sigma_i < i < i + 1 < \sigma_{i+1} < \tilde{\sigma}_{i+1},$$

which shows that $i \notin \text{des}(\sigma)$ and $i \notin \text{des}(\tilde{\sigma})$.

Case 5. Suppose (σ_i, σ_{i+1}) is a (s, f) pair. Now we have

$$\sigma_i < i < i + 1 = \sigma_{i+1} \quad \text{and} \quad \tilde{\sigma}_i \leq s(\sigma) < \tilde{\sigma}_{i+1},$$

which shows that $i \notin \text{des}(\sigma)$ and $i \notin \text{des}(\tilde{\sigma})$.

Case 6. Suppose (σ_i, σ_{i+1}) is a (f, s) pair. Then since $\sigma_{i+1} < i + 1$ and $\sigma_i = i$, we have

$$\sigma_{i+1} < \sigma_i \quad \text{and} \quad \tilde{\sigma}_{i+1} \leq s(\sigma) < \tilde{\sigma}_i.$$

This shows that $i \in \text{des}(\sigma)$ and $i \in \text{des}(\tilde{\sigma})$.

Cases 7–9. The remaining three cases are that (σ_i, σ_{i+1}) is a (f, e) , (e, s) , or (e, f) pair. These cases are handled similarly to the previous three cases and are left to the reader. \square

Theorem 2. Let $\alpha \in D_k$, $k \leq n$, and $\gamma = s(\alpha) + 1, s(\alpha) + 2, \dots, n - e(\alpha)$. Then the map $\varphi: \{\sigma \in \mathcal{S}_n \mid dp(\sigma) = \alpha\} \rightarrow \text{Sh}(\tilde{\alpha}, \gamma)$ defined by $\varphi(\sigma) = \tilde{\sigma}$ is a bijection which preserves descent sets, i.e. $\text{des}(\sigma) = \text{des}(\varphi(\sigma))$. Consequently, for all $J \subseteq \langle n - 1 \rangle$,

$$|\{\sigma \in \mathcal{S}_n \mid dp(\sigma) = \alpha, \text{des}(\sigma) = J\}| = |\{\sigma \in \text{Sh}(\tilde{\alpha}, \gamma) \mid \text{des}(\sigma) = J\}|.$$

Proof. In view of Lemma 1, we need only show that φ is an invertible map whose image is $\text{Sh}(\tilde{\alpha}, \gamma)$. First, we claim that if $dp(\sigma) = \alpha$ then $\tilde{\sigma}$ is obtained from σ by replacing the subword of nonfixed points of σ by $\tilde{\alpha}$ and the subword of fixed points of σ by γ . Indeed, the subword of fixed points of σ is replaced by the word $s(\sigma) + 1, s(\sigma) + 2, \dots, n - e(\sigma)$, which is precisely γ since $s(\sigma) = s(\alpha)$ and $e(\sigma) = e(\alpha)$. Also since α is the reduction of the subword of nonfixed points of σ , the position of the i th smallest subcedant of α is the same as the position of the i th smallest subcedant of σ in the subword of nonfixed points. The same is true for the i th smallest excedant. Hence each subcedant and excedant of σ is replaced by the same letter that replaces the corresponding subcedant or excedant of α . This means that the subword of subcedants and excedants of σ is replaced by $\tilde{\alpha}$. We may now conclude that $\tilde{\sigma} \in \text{Sh}(\tilde{\alpha}, \gamma)$.

The above description of $\tilde{\sigma}$ as a shuffle of $\tilde{\alpha}$ and γ also implies that φ is invertible. Indeed, if we replace the $\tilde{\alpha}$ subword of any $\tau \in \text{Sh}(\tilde{\alpha}, \gamma)$ by the permutation, of the subword positions, whose reduction is α , and the letters of the γ subword by their positions, we obtain a unique permutation $\sigma \in \mathcal{S}_n$ such that $dp(\sigma) = \alpha$ and $\varphi(\sigma) = \tau$. \square

Remark. Although a descent set preserving bijection between $\{\sigma \in \mathcal{S}_n \mid dp(\sigma) = \alpha\}$ and $\text{Sh}(\alpha, \beta)$, where $\beta = k + 1, k + 2, \dots, n$, will not be needed in the sequel, we should point out here that one can be constructed by composing the bijection φ with a bijection between $\text{Sh}(\alpha, \beta)$ and $\text{Sh}(\tilde{\alpha}, \gamma)$ described in [BW, Proof of Proposition 4.1].

Corollary 3. Let $\alpha \in D_k$ and $k \leq n$. Then

$$\sum_{\substack{dp(\sigma)=\alpha \\ \sigma \in \mathcal{S}_n}} q^{\text{maj}(\sigma)} = q^{\text{maj}(\alpha)} \begin{bmatrix} n \\ k \end{bmatrix}.$$

Proof. Since $\text{maj}(\sigma)$ depends only on $\text{des}(\sigma)$, it follows from Theorem 2 that

$$\begin{aligned} \sum_{dp(\sigma)=\alpha} q^{\text{maj}(\sigma)} &= \sum_{\sigma \in \text{Sh}(\tilde{\alpha}, \gamma)} q^{\text{maj}(\sigma)} \\ &= q^{\text{maj}(\tilde{\alpha})} \begin{bmatrix} n \\ k \end{bmatrix}, \end{aligned}$$

with the last step following from Garsia–Gessel [GG, Theorem 3.1]. (For a bijective alternative proof and generalization of the Garsia–Gessel result, see [BW].) By Lemma 1, $\text{maj}(\tilde{\alpha}) = \text{maj}(\alpha)$, which completes the proof. \square

Theorem 4. For all $n \geq 0$,

$$d_n(q) = [n]! \sum_{k=0}^n \frac{(-1)^k}{[k]!} q^{\binom{k}{2}}.$$

Proof. By maj- q -counting the permutations in \mathcal{S}_n according to derangement part and applying Corollary 3, we obtain

$$\begin{aligned} [n]! &= \sum_{\sigma \in \mathcal{S}_n} q^{\text{maj}(\sigma)} \\ &= \sum_{k=0}^n \sum_{\alpha \in D_k} \sum_{dp(\sigma)=\alpha} q^{\text{maj}(\sigma)} \\ &= \sum_{k=0}^n \sum_{\alpha \in D_k} q^{\text{maj}(\alpha)} \begin{bmatrix} n \\ k \end{bmatrix} \\ &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} d_k(q). \end{aligned}$$

Gauss inversion [A, p. 96] on the resulting equation yields,

$$\begin{aligned} d_n(q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} (-1)^{n-k} q^{\binom{n-k}{2}} [k]! \\ &= \sum_{k=0}^n \frac{[n]!}{[n-k]!} (-1)^{n-k} q^{\binom{n-k}{2}}, \end{aligned}$$

which is equivalent to the desired formula. \square

ACKNOWLEDGMENT

I am grateful to Adriano Garsia for the stimulating conversations which lead me to consider q -derangement numbers.

REFERENCES

- [A] M. Aigner, *Combinatorial theory*, Springer-Verlag, New York, 1979.
- [BW] A. Björner and M. Wachs, q -Hook length formulas for forests, *J. Combin. Th. Ser A* (to appear).
- [D₁] J. Désarménien, Une autre interprétation du nombre de dérangements, *Actes 8^e Séminaire Lotharingien de Combinatoire*, I.R.M.A. Strasbourg, 1984, pp. 11–16.
- [D₂] J. Désarménien, personal communication.
- [G] I. Gessel, *Counting permutations by descents, greater index, and cycle structure*, unpublished work.
- [GG] A. M. Garsia and I. Gessel, *Permutation statistics and partitions*, *Adv. in Math.* **31** (1979), 288–305.
- [GR] A. M. Garsia and J. Remmel, *A combinatorial interpretation of q -derangement and q -Laguerre numbers*, *European J. Combin.* **1** (1980), 47–59.

- [M] P. A. MacMahon, *The indices of permutations and the derivation therefrom of functions of a single variable associated with permutations of any assemblage of objects*, Amer. J. Math. **35** (1913), 281–322; reprinted in *Percy Alexander MacMahon: Collected papers*, vol. 1 (G. E. Andrews, ed.), M.I.T. Press, Cambridge MA., 1978, pp. 508–549.
- [DW] J. Désarménien and M. Wachs, *Descentes des dérangements et mots circulaires*, Actes 19 Séminaire Lotharinjien de Combinatoire, I.R.M.A. Strasbourg, 1988, 13–21.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MIAMI, CORAL GABLES, FLORIDA 33124

Current address: Department of Mathematics, 201 Walker Hall, University of Florida, Gainesville, Florida 32611