

# A Pattern Avoidance Characterization for Smoothness of Positroid Varieties

Sara Billey  
University of Washington

Based on joint work with:  
Jordan Weaver and Christian Krattenthaler

arXiv:2207.06508

February 1, 2023

**Quote.** “Combinatorics is the nanotechnology of mathematics”

Sara Billey, 2005

# Outline

Motivation: Pattern avoidance on Permutations

Pattern avoidance on Decorated Permutations

Positroid varieties

Characterizing Smooth Positroid Varieties

Enumeration

Future Work

# Permutations

**Defn.** A *permutation*  $w$  in the symmetric group  $S_n$  is a bijection from  $[n] = \{1, 2, \dots, n\}$  to itself, denoted in one-line notation  $w = [w(1), w(2), \dots, w(n)]$ .

**Example.**  $[4, 2, 1, 5, 6, 3] \in S_6$

# Permutations

**Defn.** A *permutation*  $w$  in the symmetric group  $S_n$  is a bijection from  $[n] = \{1, 2, \dots, n\}$  to itself, denoted in one-line notation  $w = [w(1), w(2), \dots, w(n)]$ .

**Example.**  $[4, 2, 1, 5, 6, 3] \in S_6$

**Example.** A perfect shuffle cuts the deck in half and drops the bottom card from each side alternating between the two piles. If done correctly, eight perfect shuffles returns the deck to the starting order. What is the one line notation for a perfect shuffle?

# Permutations

**Defn.** A *permutation*  $w$  in the symmetric group  $S_n$  is a bijection from  $[n] = \{1, 2, \dots, n\}$  to itself, denoted in one-line notation  $w = [w(1), w(2), \dots, w(n)]$ .

**Example.**  $[4, 2, 1, 5, 6, 3] \in S_6$

**Example.** A perfect shuffle cuts the deck in half and drops the bottom card from each side alternating between the two piles. If done correctly, eight perfect shuffles returns the deck to the starting order. What is the one line notation for a perfect shuffle?

$[1, 27, 2, 28, 3, 29, 4, 30, 5, 31, 6, 32, 7, 33, 8, 34, 9, 35, 10, 36,$   
 $11, 37, 12, 38, 13, 39, 14, 40, 15, 41, 16, 42, 17, 43, 18, 44, 19, 45,$   
 $20, 46, 21, 47, 22, 48, 23, 49, 24, 50, 25, 51, 26, 52]$

## Pattern Avoidance on Permutations

**Defn.** A *permutation*  $w = [w(1), w(2), \dots, w(n)] \in S_n$  *contains* another permutation  $v = [v(1), v(2), \dots, v(k)] \in S_k$  for  $k \leq n$  provided there exists  $1 < i_1 < i_2 < \dots < i_k \leq n$  such that  $[w(i_1), w(i_2), \dots, w(i_k)]$  and  $[v(1), v(2), \dots, v(k)]$  have the same relative order.

If no length  $k$  subsequence of  $[w(1), w(2), \dots, w(n)]$  has the same relative order as  $[v(1), v(2), \dots, v(k)]$  then we say  $w$  *avoids*  $v$ .

**Example.** Take  $v = [4, 2, 3, 1]$ . Then,  
 $w = [6, 2, 5, 4, 3, 1]$  contains  $6241 \sim 4231$   
 $w = [6, 1, 2, 5, 4, 3]$  avoids  $4231$

## Pattern Avoidance on Permutations

**Defn.** A *permutation*  $w = [w(1), w(2), \dots, w(n)] \in S_n$  *contains* another permutation  $v = [v(1), v(2), \dots, v(k)] \in S_k$  for  $k \leq n$  provided there exists  $1 < i_1 < i_2 < \dots < i_k \leq n$  such that  $[w(i_1), w(i_2), \dots, w(i_k)]$  and  $[v(1), v(2), \dots, v(k)]$  have the same relative order.

If no length  $k$  subsequence of  $[w(1), w(2), \dots, w(n)]$  has the same relative order as  $[v(1), v(2), \dots, v(k)]$  then we say  $w$  *avoids*  $v$ .

**Example.** Take  $v = [4, 2, 3, 1]$ . Then,  
 $w = [6, 2, 5, 4, 3, 1]$  contains  $6241 \sim 4231$   
 $w = [6, 1, 2, 5, 4, 3]$  avoids  $4231$

**Question.** Does the perfect shuffle permutation avoid or contain  $[4, 2, 3, 1]$ ?

$[1, 27, 2, 28, 3, 29, 4, 30, 5, 31, 6, 32, 7, 33, 8, 34, 9, 35, 10, 36, 11, 37, 12, 38, 1$

# Pattern Avoidance on Permutations

**Thm.**(Guillemot-Marx 2014) For every permutation  $v \in S_k$  there exists an algorithm to test if  $w \in S_n$  contains  $v$  which runs in linear time,  $O(n)$ .

# Pattern Avoidance on Permutations

Many important families of permutations have a pattern avoidance characterization and/or enumerative formulas.

1. Stack sortable permutations: 231 avoiding
2. Vexillary permutations: 2143 avoiding.
3. Smooth permutations: 3412 and 4231 avoiding.
4. Permutation patterns determine the irreducible components of singular loci of Schubert varieties.
5. Schubert varieties “defined by inclusions”: 4231, 35142, 42513, 351624.
6. Deodhar permutations/ 321-hexagon avoiding: 321, 56781234, 56718234, 46781235, 46718235.
7. Boolean permutations: 321 and 3412 avoiding.
8. Factorial Schubert varieties: 4231 and 3412 avoiding.

## Pattern Avoidance on Permutations

**Def.** A variety  $X$  is *factorial* at a point  $\iff$  the local ring at that point is a unique factorization domain.

**Thm.** (Bousquet-Mélou+Butler, 2007, conj. by Woo-Yong)  
 $X_w$  is factorial at every point  $\iff w$  avoids 4231 and 3412. Here 3412 means the 4 and 1 must be adjacent.

**Thm.** (Bousquet-Mélou+Butler) There is an explicit formula for counting the number  $f_n$  of factorial Schubert varieties for  $w \in S_n$ :

$$F(t) = \frac{(1-t)(1-4t-2t^2) - (1-5t)\sqrt{(1-4t)}}{2(1-5t+2t^2-t^3)}$$
$$= x + 2x^2 + 6x^3 + 22x^4 + 89x^5 + 379x^6 + 1661x^7 + 7405x^8 + \dots$$

# Pattern Avoidance on Permutations

Bridget Tenner's Database of Permutation Patterns has 61 families!

← → ↻ math.depaul.edu/bridget/cgi-bin/dppa.cgi

My Drive - Google... Post Race Summary UW Grad Admissi... Faculty Portal Webapps Main Pa... Microsoft Word -...

## Database of Permutation Pattern Avoidance

**ID:** P0001  
**Patterns:** 2 3 1  
**Title:** Stack sortable permutations  
**References:** D.E. Knuth, The Art of Computer Programming, Volume 1, Addison-Wesley, Reading, Massachusetts, 1968.  
R. Tarjan, Sorting using networks of queues and stacks, Journal of the ACM 19 (1972), 341-346.  
**Enumeration:** Catalan numbers:  $C(n) = \text{binomial}(2n, n) / (n+1)$   
**OEIS:** A000108  
**Contributor:** Bridget Tenner - November 13, 2005

**ID:** P0002  
**Patterns:** 1 2 1  
**Title:** Fully commutative permutations  
Permutations for which depth equals length  
Permutations  $w$  for which the interval  $[1, w]$  in the weak Bruhat order on  $S_n$  is a distributive lattice  
**References:** S.C. Billey, W. Jockusch, and B.P. Stanley, Some combinatorial properties of Schubert polynomials, J. of Alg. Combin. 2 (1993), 345-374.  
T.K. Petersen and B.E. Tenner, The depth of a permutation, arxiv:1202.4765  
**Enumeration:** Catalan numbers:  $C(n) = \text{binomial}(2n, n) / (n+1)$   
**OEIS:** A000108  
**Contributor:** Bridget Tenner - November 13, 2005  
Richard Stanley - July 5, 2012

**ID:** P0003  
**Patterns:** 1 3 2  
**Title:** Dominant permutations  
The Schubert polynomial is a single monomial  
**References:** L. Manivel, Symmetric Functions, Schubert Polynomials, and Degeneracy Loci, DMS/AMS Texts and Monographs, 4, American Mathematical Society, Providence, RI.  
I.G. Macdonald, Notes on Schubert Polynomials, Publ. Laboratoire de Combinatoire et d'Informatique Mathématique 6, Université de Québec à Montréal, 1991.  
**Enumeration:** Catalan numbers:  $C(n) = \text{binomial}(2n, n) / (n+1)$   
**OEIS:** A000108  
**Contributor:** Bridget Tenner - November 13, 2005  
Richard Stanley - November 14, 2005


The International Conference on Permutation Patterns 2023 will take place at the University of Burgundy located in Dijon, France!

# Decorated Permutations

**Defn.** A *decorated permutation*  $w^\circ$  on  $n$  elements is a permutation  $w \in S_n$  together with an orientation clockwise or counterclockwise assigned to each fixed point of  $w$ , denoted  $\overrightarrow{i}$  or  $\overleftarrow{i}$  respectively.

$$w^\circ = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 5 & 4 & 1 & 2 & 7 & \overleftarrow{6} & 9 & \overrightarrow{8} & 3 \end{bmatrix} = 54127\overleftarrow{6}9\overrightarrow{8}3$$

*Chord Diagram*  $D(w^\circ)$



# Decorated Permutations

Sylvie Corteel studied the  $q$ -analogs of Eulerian numbers related to the number of alignments and weak exceedances of permutations, which come from the enumeration of totally positive Grassmann cells following work of Williams and Postnikov.

**Question.** (Corteel, 2006) Can we define generalized patterns for decorated permutations?

**Question.** (Weaver, 2018) Are there interesting families of decorated permutations avoiding certain generalized patterns?

# Decorated Permutations

Sylvie Corteel studied the  $q$ -analogs of Eulerian numbers related to the number of alignments and weak exceedances of permutations, which come from the enumeration of totally positive Grassmann cells following work of Williams and Postnikov.

**Question.** (Corteel, 2006) Can we define generalized patterns for decorated permutations?

**Question.** (Weaver, 2018) Are there interesting families of decorated permutations avoiding certain generalized patterns?

**Answer.** Yes! The ones indexing smooth positroid varieties.

# Grassmannian Varieties

## Notation.

- ▶ Fix  $0 \leq k \leq n$ .
- ▶  $\text{Mat}_{kn}$  = the set of full rank  $k \times n$  matrices over  $\mathbb{C}$ .
- ▶  $Gr(k, n) =$  *Grassmannian variety* of  $k$ -dim subspaces in  $\mathbb{C}^n$
- ▶  $Gr(k, n) \approx \text{GL}_k \backslash \text{Mat}_{kn}$ .
- ▶  $\Delta_J(M) =$  determinant of the  $[k] \times J$  minor of  $M$  for  $J \in \binom{[n]}{k}$ .

The Grassmannian varieties are smooth manifolds via the *Plücker coordinate* embedding  $M \mapsto (\Delta_J(M) : J \in \binom{[n]}{k})$  of  $Gr(k, n)$  into projective space, including the case  $k = n = 0$ .

## Plücker coordinates

$$\begin{bmatrix} 0 & 3 & 1 & 2 & 4 & 0 \\ 0 & 0 & 0 & 1 & 2 & 1 \end{bmatrix} \mapsto [0:0:0:0:0:0:3:6:3:1:2:1:0:2:4] \in \mathbb{P}^{15}$$

Nonvanishing coordinates at exactly the coordinates

$$\{\{2, 4\}, \{2, 5\}, \{2, 6\}, \{3, 4\}, \{3, 5\}, \{3, 6\}, \{4, 6\}, \{5, 6\}\} \subseteq \binom{[6]}{2}.$$

**Def.** The *matroid* of  $A \in \text{Mat}_{kn}$  is determined by the set of *bases*

$$\mathcal{M}_A = \left\{ J \in \binom{[n]}{k} \mid \Delta_J(A) \neq 0 \right\}.$$

or by the *non-bases*

$$\mathcal{Q}_A = \left\{ J \in \binom{[n]}{k} \mid \Delta_J(A) = 0 \right\}.$$

# Positroids

**Defn.** (Postnikov) A *positroid* of rank  $k$  on ground set  $[n]$  is a matroid of the form  $\mathcal{M}_A$  for a matrix  $A \in \text{Mat}_{kn}$  such that every nonzero Plücker coordinate  $\Delta_J(A)$  is positive.

**Thm.** (Postnikov 2006 + Oh 2011) There are bijections

1. positroids  $\mathcal{M}$  of rank  $k$  on a ground set of size  $n$ ,
2. decorated permutations  $w^\circ$  on  $n$  elements with  $k$  anti-exceedances,
3. Grassmann necklaces  $\mathcal{N} = (I_1, \dots, I_n) \in \binom{[n]}{k}^n$ , and
4. Grassmann intervals  $[u, v]$  in  $Gr(k, n)$ .

# Positroids

**Defn.** (Postnikov) A *positroid* of rank  $k$  on ground set  $[n]$  is a matroid of the form  $\mathcal{M}_A$  for a matrix  $A \in \text{Mat}_{kn}$  such that every nonzero Plücker coordinate  $\Delta_J(A)$  is positive.

**Thm.** (Postnikov 2006 + Oh 2011) There are bijections

1. positroids  $\mathcal{M}$  of rank  $k$  on a ground set of size  $n$ ,
2. decorated permutations  $w^\circ$  on  $n$  elements with  $k$  anti-exceedances,
3. Grassmann necklaces  $\mathcal{N} = (I_1, \dots, I_n) \in \binom{[n]}{k}^n$ , and
4. Grassmann intervals  $[u, v]$  in  $Gr(k, n)$ .

Important in the theory of the totally positive part of the Grassmannian variety, cluster algebras, and soliton solutions to the KP equations and have connections to statistical physics, integrable systems, and scattering amplitudes.

See Williams.ICM.2022, along with Lusztig.1994, Rietsch.1998, Fomin-Zelevinsky.2002, Kodama-Williams.2015, AHBCGPT.2016

# Positroid Varieties

**Defn.** (Rietsch, Postnikov, Knutson-Lam-Speyer)

Given a decorated permutation  $w^\circ \in S_{n,k}^\circ$  along with its associated Grassmann interval  $[u, v]$  and positroid  $\mathcal{M} = \mathcal{M}(w^\circ) \subseteq \binom{[n]}{k}$ , the *positroid variety*  $\Pi_{w^\circ} = \Pi_{[u,v]} = \Pi_{\mathcal{M}}$  is the subvariety of  $Gr(k, n)$  with vanishing ideal generated by the Plücker coordinates indexed by the nonbases of  $\mathcal{M}$ ,  $\{\Delta_J : J \notin \mathcal{M}\}$ .

**Thm.** (Knutson-Lam-Speyer) The positroid variety  $\Pi_{[u,v]}$  is the projection of the Richardson variety  $X_u^v \subseteq \mathcal{Fl}(n)$  to  $Gr(k, n)$ .

# Positroid Varieties

The vanishing coordinates of

$$A = \begin{bmatrix} 0 & 3 & 1 & 2 & 4 & 0 \\ 0 & 0 & 0 & 1 & 2 & 1 \end{bmatrix} \mapsto [0:0:0:0:0:0:3:6:3:1:2:1:0:2:4]$$

are exactly the coordinates indexed by the nonbases of  $\mathcal{M}_A$

$$\{\{1,2\}, \{1,3\}, \{1,4\}, \{1,5\}, \{2,3\}, \{4,5\}\} \subseteq \binom{[6]}{2}.$$

Therefore, the points in the positroid variety  $\Pi_{\mathcal{M}_A}$  are represented by the full rank complex matrices of the form

$$\begin{bmatrix} 0 & a_{12} & ca_{12} & a_{14} & da_{14} & a_{16} \\ 0 & a_{22} & ca_{22} & a_{24} & da_{24} & a_{26} \end{bmatrix}.$$

# Smooth versus singular varieties

**Def.** A variety  $X$  is *singular* at a point  $x \in X$ , if the dimension of the tangent space to  $X$  at  $x$  is strictly larger than the dimension of  $X$ . If no such point exists,  $X$  is *smooth*.

**Def.** If  $X = \text{Var}(f_1, \dots, f_s)$ , the *Jacobian matrix*,  $Jac$ , is the matrix of partial derivatives of the  $f_i$  at each of the possible variables.

**Def.** The  $\text{rank}(Jac|_x)$  is the codimension of the tangent space to  $X$  at the point  $x$ .

# Characterizing Smooth Positroid Varieties

**Thm.** (Billey-Weaver 2023) Let  $\mathcal{M}$  be a rank  $k$  positroid on  $[n]$  with associated decorated permutation  $w^\circ$ , and Grassmann interval  $[u, v]$ . Then, the following are equivalent.


1. The positroid variety  $\Pi_{w^\circ} = \Pi_{[u, v]} = \Pi_{\mathcal{M}}$  is smooth.
2. The decorated permutation  $w^\circ$  has no crossed alignments.
3. The chord diagram  $D(w^\circ)$  is a disjoint union of spirographs.
4. The positroid  $\mathcal{M}$  is a direct sum of uniform matroids.

# Some Patterns in Decorated Permutations

## Patterns using chord diagrams.

- ▶ *alignments*: two directed edges in same direction
- ▶ *misalignments*: two directed edges in opposite direction
- ▶ *crossings*: two edges that must have a common point
- ▶ *crossed alignment*: an alignment plus a third edge crossing both sides

Consider the decorated permutation  $w^\circ = 895 \overleftarrow{4} 7 \overrightarrow{6} 132$






The highlighted crossed alignment pattern implies  $\prod w^\circ$  is singular.

# Patterns in Decorated Permutations


## Patterns.

- ▶ *spirograph*: chord diagram of a connected decorated perm with  $w(i) \mapsto i + m$  for all  $i \in [n]$ .



Thx to "Spirographs" made by the Spirograph Maker app for the iphone.

# Chord diagram of a smooth positroid variety



# Decorated Permutations to Grassmann intervals

**Defn.** The *anti-exceedance set* of  $w^\circ$  is

$$I_1(w^\circ) = \{i \in [n] \mid i < w^{-1}(i) \text{ or if } w^\circ(i) = \overrightarrow{i}\}$$

If  $i \in [n]$  is not an anti-exceedance, it is an *exceedance*. Let  $S_{n,k}^\circ$  be the set of decorated permutations on  $n$  elements with  $k$  anti-exceedances.

$$w^\circ = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 5 & 4 & 1 & 2 & 7 & \overleftarrow{6} & 9 & \overleftarrow{8} & 3 \end{bmatrix} \implies I_1(w^\circ) = \{1, 2, 6, 3\}.$$

Shuffling the anti-exceedances to the front bijectively determines the *Grassmann interval*  $[u, v]$  associated to  $w^\circ \in S_{n,k}^\circ$ , where  $u \leq v$  in Bruhat order on  $S_n$  and  $v$  is *k-Grassmannian*, so  $v_1 < v_2 < \dots < v_k$  and  $v_{k+1} < \dots < v_n$ .

$$\begin{bmatrix} v \\ u \end{bmatrix} = \begin{bmatrix} 3 & 4 & 6 & 9 & 1 & 2 & 5 & 7 & 8 \\ 1 & 2 & 6 & 3 & 5 & 4 & 7 & 9 & 8 \end{bmatrix}.$$

## Decorated Permutations to Positroids

The *Grassmann necklace* for  $w^\circ = 54127 \overrightarrow{6} 9 \overleftarrow{8} 3$  is

$$(l_1, \dots, l_9) = (\{1236\}, \{2356\}, \{3456\}, \{1456\}, \{1256\}, \\ \{1267\}, \{1267\}, \{1269\}, \{1269\}).$$

The *positroid* for  $w^\circ$  is


$$\begin{array}{ccccccc} 1236 & 1246 & 1256 & 1267 & 1269 & 1346 & 1356 \\ 1367 & 1369 & 1456 & 1467 & 1469 & 2356 & 2367 \\ 2369 & 2456 & 2467 & 2469 & 3456 & 3467 & 3469 \end{array}$$

Here, for each  $r \in [n]$ , let  $r <_r <_r r + 1 < \dots n <_r 1 <_r < \dots <_r r - 1$ ,

$$l_r(w^\circ) = \{i \in [n] \mid i <_r w^{-1}(i) \text{ or if } w^\circ(i) = \overrightarrow{i}\},$$

$$\mathcal{M}(w^\circ) := \left\{ I \in \binom{[n]}{k} : l_r(w^\circ) \leq_r I \text{ for all } r \in [n] \right\}$$

# Commutative Diagram of Bijections



**Grassmann intervals to Positroids via initial sets.**

$$\mathcal{M} = \{y[k] : y \in [u, v]\}.$$

# Characterizing Smooth Positroid Varieties

**Thm.** (Billey-Weaver 2023) Let  $\mathcal{M}$  be a rank  $k$  positroid on  $[n]$  with associated decorated permutation  $w^\circ$ , and Grassmann interval  $[u, v]$ . Then, the following are equivalent.

1. The positroid variety  $\Pi_{w^\circ} = \Pi_{[u, v]} = \Pi_{\mathcal{M}}$  is smooth.
2. The decorated permutation  $w^\circ$  has no crossed alignments.
3. The chord diagram  $D(w^\circ)$  is a disjoint union of spirographs.
4. The positroid  $\mathcal{M}$  is a direct sum of uniform matroids.

# Outline of Proof

**Thm.** [Postnikov, KLS]

$$\text{codim}(\Pi_{w^\circ}) = \#Alignments(w^\circ) = k(n - k) - [\ell(v) - \ell(u)].$$

**Corollary.** A positroid variety  $\Pi_{w^\circ}$  is a singular at  $x$  if and only if

$$\text{rank}(Jac|_x) < \text{codim} \Pi_{w^\circ} = \#Alignments(w^\circ).$$

## Outline of Proof

**By Construction.:** For every  $A \in \Pi_{\mathcal{M}}$ , the first nonzero Plücker coordinate in lex order of  $A$  is in  $\mathcal{M}$ . For every  $J \in \mathcal{M}$ , the matrix which is the identity in columns  $J$ , denoted  $A_J \in \Pi_{\mathcal{M}}$ .

**Thm.** Assume  $A \in \Pi_{\mathcal{M}}$  and  $J \in \mathcal{M}$  indexes its first nonzero Plücker coordinate in lex order. Then the codimension of the tangent space to  $\Pi_{\mathcal{M}}$  at  $A$  is bounded below by


$$\text{rank}(\text{Jac}|_{A_J}) = \#\left\{I \in \binom{[n]}{k} \setminus \mathcal{M} : |I \cap J| = k - 1\right\}.$$

**Hard Step.:** Given a starboard tacking crossed alignment based at 1, the set on RHS maps injectively into the set of alignments of  $w^{\circ}$  and for  $J = I_1(w^{\circ}) \in \mathcal{M}$  there is at least one alignment that is not in the image.

# Johnson Graphs

The *Johnson graph*  $J(k, n)$  has vertices labeled by  $k$ -subsets of  $[n]$  and two  $k$ -subsets  $I, J$  are connected by an edge precisely if  $|I \cap J| = k - 1$ .

For a positroid  $\mathcal{M} \subseteq \binom{[n]}{k}$ , let  $J(\mathcal{M})$  denote the induced subgraph of the Johnson graph on the vertices in  $\mathcal{M}$ . We call  $J(\mathcal{M})$  the *Johnson graph of  $\mathcal{M}$* .



# Regular Johnson Graphs and Smooth Positroid Varieties

**Thm.** (Billey-Weaver 2022) Let  $\mathcal{M}$  be a rank  $k$  positroid on  $[n]$  with associated decorated permutation  $w^\circ$ , and Grassmann interval  $[u, v]$ . Then, the following are equivalent.

1. The positroid variety  $\Pi_{w^\circ} = \Pi_{[u, v]} = \Pi_{\mathcal{M}}$  is smooth.
2. The decorated permutation  $w^\circ$  has no crossed alignments.
3. The chord diagram  $D(w^\circ)$  is a disjoint union of spirographs.
4. The positroid  $\mathcal{M}$  is a direct sum of uniform matroids.
5. The graph  $J(\mathcal{M})$  is regular, and each vertex has degree  $\ell(v) - \ell(u)$ .

## Enumerative Results

Similar to Ardila-Rincón-Williams enumeration of positroids via connected positroids, we use a theorem of Speicher to enumerate all smooth positroids via connected smooth positroids (aka spirograph permutations).

**Thm.** The number of smooth positroids on ground set  $[n]$  is the coefficient

$$s(n) = [x^n] \frac{1}{n+1} \left( 1 + 2x + \sum_{i=2}^n (i-1)x^i \right)^{n+1}. \quad (1)$$

## Enumerative Results

Similar to Ardila-Rincón-Williams enumeration of positroids via connected positroids, we use a theorem of Speicher to enumerate all smooth positroids via connected smooth positroids (aka spirograph permutations).

**Thm.** The number of smooth positroids on ground set  $[n]$  is the coefficient

$$s(n) = [x^n] \frac{1}{n+1} \left( 1 + 2x + \sum_{i=2}^n (i-1)x^i \right)^{n+1}. \quad (1)$$


$$s(51)/s(50) \approx 5.4489775,$$

$$s(101)/s(100) \approx 5.528236,$$

$$s(151)/s(150) \approx 5.555362,$$

$$s(201)/s(200) \approx 5.569062,$$

$$s(251)/s(250) \approx 5.5773263.$$

Based on this data, we conjectured  $s(n) \approx O(c^n)$  for  $c \leq 6$ . 

## Enumerative Results

Let  $f(x) = x^{n+1}$  and let  $g(x) = 1 + 2x + \sum_{i=2}^n (i-1)x^i$ , then

$$s(n) = \frac{1}{n+1} [x^n] f(g(x)).$$

So take  $\frac{d^n}{dx^n} f(g(x))$ , set  $x = 0$  and divide by  $n + 1$ .

## Enumerative Results

Let  $f(x) = x^{n+1}$  and let  $g(x) = 1 + 2x + \sum_{i=2}^n (i-1)x^i$ , then

$$s(n) = \frac{1}{n+1} [x^n] f(g(x)).$$

So take  $\frac{d^n}{dx^n} f(g(x))$ , set  $x = 0$  and divide by  $n+1$ .

**Faá di Bruno Formula.** (1855) Generalizing the chain rule from Calculus,

$$\frac{d^n}{dx^n} f(g(x)) = \sum_{k=1}^n f^{(k)}(g(x)) \cdot B_{n,k}(g'(x), g''(x), \dots, g^{(n+1-k)}(x)) \quad (2)$$

where *partial Bell polynomial*  $B_{n,k}(x_1, \dots, x_{n-k+1})$  is defined as

$$B_{n,k}(x_1, \dots, x_{n-k+1}) = \sum_{B_1 \sqcup \dots \sqcup B_k} \prod_{i=1}^k x_{|B_i|}, \quad (3)$$

where the sum is taken over all set partitions of  $[n]$  into  $k$  blocks.

# Enumerative Results

**Notation.** For  $1 \leq k \leq n$ , let  $b_{0,0} = 1$ , and  $b_{0,k} = b_{n,0} = 0$  if  $n > 0$  or  $k > 0$ ,

- ▶  $B_{n,k}(x_1, \dots, x_{n-k})$  Bell polynomial
- ▶  $b_{n,k} = B_{n,k}(2 \cdot 1!, 1 \cdot 2!, 2 \cdot 3!, \dots, (n-k) \cdot (n-k+1)!)$

**Cor.** The number of smooth positroids on ground set  $[n]$  is

$$s(n) = \sum_{k=1}^n \frac{b_{n,k}}{(n-k+1)!}. \quad (4)$$

# Enumerative Results

**Notation.** For  $1 \leq k \leq n$ , let  $b_{0,0} = 1$ , and  $b_{0,k} = b_{n,0} = 0$  if  $n > 0$  or  $k > 0$ ,

- ▶  $B_{n,k}(x_1, \dots, x_{n-k})$  Bell polynomial
- ▶  $b_{n,k} = B_{n,k}(2 \cdot 1!, 1 \cdot 2!, 2 \cdot 3!, \dots, (n-k) \cdot (n-k+1)!)$

**Cor.** The number of smooth positroids on ground set  $[n]$  is

$$s(n) = \sum_{k=1}^n \frac{b_{n,k}}{(n-k+1)!}. \quad (4)$$

**Cor.** The number of smooth positroids on ground set  $[n]$  with  $k$  connected components is  $s_2(n, k) = \frac{b_{n,k}}{(n-k+1)!}$ .

## Enumerative Results

**Thm.**(Krattenthaler) For all non-negative integers  $n$ , the number of smooth positroids on the ground set  $[n]$  is given by

$$s(n) = \frac{1}{n+1} \sum_{r=0}^{\lfloor (n+1)/2 \rfloor} (-1)^r 2^r \binom{n+1}{r} \binom{3n-3r+1}{n-2r}. \quad (5)$$

**Thm.**(Krattenthaler) The sequence  $(s(n))_{n \geq 0}$  satisfies the recurrence relation

$$\begin{aligned} & 2(n-2)(n+1)(2n+1)(26n-33)s(n) \\ & - (1118n^4 - 3343n^3 + 2092n^2 + 367n - 66)s(n-1) \\ & + 2(n-1)(2002n^3 - 6181n^2 + 4435n + 18)s(n-2) \\ & - 4(n-2)(n-1)(1586n^2 - 3547n + 555)s(n-3) \\ & + 152(n-3)(n-2)(n-1)(26n-7)s(n-4) = 0, \quad (6) \end{aligned}$$

with initial conditions  $s(0) = 1$ ,  $s(1) = 2$ ,  $s(2) = 5$ , and  $s(3) = 16$ .

# Enumerative Results

**Thm.**(Krattenthaler) The asymptotic growth of the number of smooth positroids on the ground set  $[n]$  is given by

$$s(n) \sim \frac{\rho^{n+1}}{n^{3/2}\sqrt{2\pi\xi}} = \frac{(5.61071\dots)^{n+1}}{(8.74042\dots) \cdot n^{3/2}}, \quad \text{as } n \rightarrow \infty, \quad (7)$$

where  $\rho = 5.61071\dots$  is the real root of

$$4z^3 - 35z^2 + 84z - 76 = 0,$$

and  $\xi = 12.15864\dots$  is the real root of

$$38z^3 - 425z^2 - 416z - 416 = 0.$$

# Enumerative Results

**Thm.**(Krattenthaler) For all integers  $n$  and  $k$  with  $n \geq k > 0$ , the number of smooth decorated permutations in  $S_n^\circ$  with exactly  $k$  connected components is given by

$$s(n, k) = \frac{1}{n+1} \binom{n+1}{k} \sum_{l=0}^k 2^{k-l} \binom{k}{l} \binom{n-k+l-1}{n-k-l}. \quad (8)$$

# Future Work

## Open problems.

1. Study the geometry and equivariant cohomology of a positroid variety via the induced directed Johnson graph on  $\mathcal{M}$ .
2. Identify the singular locus of a positroid variety.
3. Study the enumeration and coset structure of the group operations of flip, inverse, rotation for permutations, or for derangements, or SIF perms.
4. Study the combinatorics of the directed Johnson graphs of positroids.

Thank you!

