

MODULI OF ELLIPTIC SURFACES OF KODAIRA DIMENSION ONE FIBERED OVER RATIONAL CURVES

DORI BEJLERI, JOSIAH FOSTER, ANDRES FERNANDEZ HERRERO,
GIOVANNI INCHIOSTRO, SVETLANA MAKAROVA, AND JUNYAN ZHAO

ABSTRACT. In this article, we construct an infinite sequence of irreducible components of Kollár–Shepherd-Barron (KSB-) moduli spaces of surfaces of arbitrarily large volumes, and describe the boundary of each component completely. Moreover, we describe the stable reduction steps in finding the KSB-limits in an explicit combinatorial way. Our main approach is to study the moduli spaces of elliptic surfaces with Kodaira dimension one, fibered over rational curves, using the techniques of wall-crossing for KSBA moduli and twisted stable maps.

CONTENTS

1. Introduction	1
2. Preliminaries on stable pairs	4
3. Elliptic surfaces	7
4. Moduli spaces of elliptic surfaces	9
5. Local study of the contraction morphism Ψ_n	16
6. The case when $n = 3$	19
7. Compactifications	22
References	33

1. INTRODUCTION

The moduli spaces of smooth surfaces of general type admit Kollár–Shepherd-Barron (abbr. KSB) compactifications, which allow smooth objects to degenerate to certain singular surfaces satisfying *KSB-stability* ([KSB88]). The KSB-stable surfaces of a fixed volume v are parametrized by a proper Deligne-Mumford stack $\mathcal{M}_v^{\text{KSB}}$, whose coarse space is known as the KSB moduli space. Classical questions about the classification of surfaces can be translated, via the moduli theory, to questions about the geometry of the corresponding KSB moduli spaces. It is therefore an interesting and important endeavor to gain an explicit understanding of irreducible components of this moduli problem.

Besides the classical case of the moduli of curves (see [DM69]), there are currently few instances in the literature where the geometry of an irreducible component of the KSB moduli spaces of varieties is completely understood. Even fewer examples are known to admit an explicit combinatorial description of the singular varieties parametrized by the boundary of their respective components of the moduli space. The first examples in dimension higher than one, where the boundary was fully understood, include quotients or special covers of a product of two curves (see [vO06, Liu12, Rol10]), followed by the Campedelli surfaces and Burniat surfaces; see [AP23]. These are surfaces which admit a cover to some toric del Pezzo surface X_0 branched along a line arrangement $\sum L_i$, so one can reduce the study to the description of moduli spaces of pairs $(X_0, c \sum L_i)$, where $c > 0$ is a (rational) coefficient which makes the covering map crepant.

In this paper, we provide a novel example. We identify an infinite sequence of irreducible components $\overline{\mathcal{P}}_n$ of the KSB-moduli spaces of stable surfaces of volume $\frac{(n-2)^2}{n}$, depending on a parameter $n \in \mathbb{Z}_{\geq 3}$. Every point in $\overline{\mathcal{P}}_n$ represents a singular(!) *pseudo-elliptic surface* over a rational curve (not necessarily irreducible), and a general such surface contains a unique quotient singularity of type $\frac{1}{n}(1, 1)$. Moreover, we develop an

explicit combinatorial description of the closed boundary, which parametrizes surfaces with worse singularities than an isolated $\frac{1}{n}(1, 1)$.

Moduli of Weierstrass fibrations. The objects we study in this paper are elliptic surfaces of Kodaira dimension 1, fibered over a rational curve with a section S such that $S^2 = -n$. They are projective surfaces X which admit a proper equidimensional morphism $f: X \rightarrow C$ to a rational curve C (not necessarily irreducible) such that each fiber is a connected curve of arithmetic genus 1. A general such surface $(f: X \rightarrow C, S)$ satisfies that X is smooth, $C \simeq \mathbb{P}^1$, and the fibers of f are either smooth or nodal. The number $n \geq 0$ is a fundamental invariant of the elliptic surface called the *height*. The moduli of elliptic surfaces was first constructed using GIT by Miranda [Mir89]. It parametrizes Weierstrass fibrations (see Definition 4.1), which are elliptic surfaces with integral fibers such that the total space has log canonical singularities. We denote by $\mathcal{W}_n^{\text{lc}}$ (resp. $\mathcal{W}_n^{\text{min}}$) the moduli stacks parametrizing such Weierstrass fibrations with at worst log canonical singularities (resp. Weierstrass fibrations with at worst canonical singularities).

The main observation is the following. Given a general elliptic surface of Kodaira dimension 1 over \mathbb{P}^1 ($f: X \rightarrow \mathbb{P}^1, S$) with a section S such that $S^2 = -n$, there are two natural constructions to perform:

- (*KSB moduli*). Contracting the negative section S results in a surface Y with klt singularities, and the canonical divisor K_Y is \mathbb{Q} -Cartier and ample of volume $v = \frac{(n-2)^2}{n}$. Such a surface Y , called a *pseudo-elliptic surface*, is represented by a point in the KSB-moduli stack $\mathcal{M}_v^{\text{KSB}}$ (see Definition 2.17).
- (*Twisted stable maps*). Since the fibers of f are connected curves of arithmetic genus 1 with at worst nodal singularities, and the section S intersects each fiber at a smooth point of the fiber, then by the universality of $\overline{\mathcal{M}}_{1,1}$, one obtains a natural morphism $C \rightarrow \overline{\mathcal{M}}_{1,1}$. Such a morphism is parametrized by the moduli stack of twisted stable maps $\mathcal{K}_n := \overline{\mathcal{K}}_{0,0}(\overline{\mathcal{M}}_{1,1}, n)$ (see Definition 7.1), which was first introduced in [AV02].

We establish a link between the KSB moduli stacks and the moduli of twisted stable maps, using the KSBA moduli stacks $\mathcal{M}_{c,v}^{\text{KSBA}}$ of log pairs (X, cS) (see Definition 2.14) as a bridge, where $0 < c < \frac{n-2}{n}$ is a rational coefficient. Now we state our first main theorem.

Theorem 1.1 (= Proposition 4.8 + Proposition 4.13 + Theorem 5.4 + Theorem 7.2 + Proposition 7.3). *Let $n \geq 3$ be an integer with $n \neq 4$, and let $0 < \epsilon \ll 1$ be a rational number. Set $v := \frac{(n-2)^2}{n}$, $c(\epsilon) := \frac{n-2}{n} - \epsilon$, and $v(\epsilon) := v - n\epsilon$. Then the following statements hold.*

- (1) *For any $0 < \epsilon < \frac{n-2}{n}$, there is a natural morphism*

$$\Phi_{n,\epsilon} : \mathcal{W}_n^{\text{lc}} \longrightarrow \mathcal{M}_{c(\epsilon),v(\epsilon)}^{\text{KSBA}}, \quad (f: X \rightarrow C, S) \mapsto (X, c(\epsilon)S),$$

which is an open immersion.

- (2) *Let \mathcal{E}_n (resp. $\mathcal{E}_n^{\text{lc}}$) be the image of $\mathcal{W}_n^{\text{min}}$ (resp. $\mathcal{W}_n^{\text{lc}}$) under $\Phi_{n,\epsilon}$. Then there is a natural morphism*

$$\Psi_n : \mathcal{E}_n^{\text{lc}} \longrightarrow \mathcal{M}_v^{\text{KSB}}, \quad (X, c(\epsilon)S) \mapsto Y,$$


which is an open immersion. Here, the surface Y is obtained from X by contracting the section S .

- (3) *The stack \mathcal{K}_n of twisted stable maps is irreducible and proper. Its normalization \mathcal{K}_n^ν admits a natural birational morphism $\Xi_n : \mathcal{K}_n^\nu \longrightarrow \overline{\mathcal{E}}_{n,\epsilon}$, where $\overline{\mathcal{E}}_{n,\epsilon}$ denotes the closure of $\mathcal{E}_n \subset \mathcal{M}_{c(\epsilon),v(\epsilon)}^{\text{KSBA}}$.*

In particular, the moduli stacks \mathcal{K}_n , $\mathcal{W}_n^{\text{lc}}$ and the irreducible components of $\mathcal{M}_{c(\epsilon),v(\epsilon)}^{\text{KSBA}}$ and $\mathcal{M}_v^{\text{KSB}}$ parametrizing (pseudo)elliptic surfaces of height n over a rational curve.

Let $\overline{\mathcal{P}}_n$ be the closure of the image \mathcal{P}_n of Ψ_n . Then each $\overline{\mathcal{P}}_n$ and $\overline{\mathcal{E}}_{n,\epsilon}$ are irreducible components of their respective KSB- and KSBA-moduli. The compactifications $\overline{\mathcal{E}}_{n,\epsilon}$ are related by wall-crossing morphisms as ϵ varies, and we can view $\overline{\mathcal{P}}_n$ as the moduli space at the wall $\epsilon = 0$ [AB21, Inc20, ABIP23, MZ23].

We now summarize all the moduli stacks that we have introduced in one diagram.



To understand the surfaces parametrized by the boundary of $\overline{\mathcal{P}}_n$, we take a morphism from $\text{Spec } R$, where R is a DVR, to the stack of twisted stable maps \mathcal{K}_n , and run the (relative) minimal model program (abbrev. MMP) for the family pulled back from the universal family over $\overline{\mathcal{M}}_{1,1}$. We can associate a graph Γ , called a *sliced tree*, to each elliptic surface $(f: X \rightarrow C, S)$ derived from a twisted stable map (i.e. *tsm-stable elliptic surface*) in such a way that each vertex of Γ represents an irreducible component of X and each edge between two vertices means that the two corresponding components intersect (see Definition 7.15). Furthermore, such a sliced tree includes the data of certain numerical decorations for its vertices (which encode the degree of the j -map of each component) and decorations for its edges (which encode the gluing data for any two given components). We show that the MMP steps (i.e. stable reduction) can be described as a pruning process of the associated graphs. The main operations of the pruning process include cutting down edges and adding *klt-markings* and *lc-markings*.

Theorem 1.2. *The stable reduction in the KSB moduli spaces $\overline{\mathcal{P}}_n$ is given by Algorithm 7.11, which computes the KSB-stable limit. Moreover, the stable reduction corresponds to a pruning process of the graph associated to the tsm-stable elliptic surfaces.*

An immediate corollary of the above two theorems is the following.

Corollary 1.3. *Let $n \geq 3$ be an integer with $n \neq 4$, and $v := \frac{(n-2)^2}{n}$. Then $\overline{\mathcal{P}}_n$ is an irreducible component of $\mathcal{M}_v^{\text{KSB}}$, is proper and of dimension $10n - 2$. The interior of $\overline{\mathcal{P}}_n$ parametrizes pseudo-elliptic surfaces with an isolated $\frac{1}{n}(1, 1)$ -singularity, and the boundary of $\overline{\mathcal{P}}_n$ parametrizes pseudo-elliptic surfaces whose associated graph is a pruned tree of height n .*

Finally, we state a more explicit result in the special case when $S^2 = -3$: then the cohomology groups encoding deformation theory can be computed explicitly (Section 6), and the combinatorics is relatively simple.

Theorem 1.4. *Let $n = 3$ and X be a surface in $\overline{\mathcal{P}}_3$.*

- (1) *The pruned tree Π associated to X is a chain with at most six vertices.*
- (2) *If X is in the interior \mathcal{P}_3 , then one has $h^1(X, T_X) = 28$ and $h^2(X, T_X) = 0$. In particular, \mathcal{P}_3 is an irreducible smooth stack of dimension 28.*

Remark 1.5. We remark that for $n = 4$, we still obtain an irreducible component $\overline{\mathcal{P}}_4$ of the KSB-moduli space and the combinatorics of the set of boundary objects is still captured by pruned trees. However, the deformation theory fails and so the map $\mathcal{W}_n^{\text{min}} \rightarrow \overline{\mathcal{P}}_4$ may not be birational. In particular, it is an open question whether $\overline{\mathcal{P}}_4$ has generically non-reduced structure. See Remark 5.6.

Prior and related works. Beyond the relevant work that was mentioned before, this paper uses extensively the results of [LN02, AB17, AB21, Inc20] on moduli of elliptic surfaces. In particular, the existence of the morphism $\mathcal{K}_n^\nu \rightarrow \overline{\mathcal{E}}_n$ follows from general wall-crossing phenomena studied in [AB21, Inc20, ABIP23, MZ23], whereas the understanding of the limits in $\overline{\mathcal{P}}_n$ follows from [LN02, AB21, Inc20]. More precisely, the morphism Ξ_n was studied in [AB21], as a special case of general wall-crossing phenomena of [ABIP23, MZ23]. The explicit combinatorial gadget of Section 7.2 is instead an improvement of the refined numerical data defined in [Inc20], for elliptic surfaces of Kodaira dimension one fibered over \mathbb{P}^1 .

Outline of the paper. In Sections 2 and 3, we compile some preliminaries on elliptic surfaces, KSB(A)-stability and KSB(A)-moduli spaces. In Sections 4 and 5, we study different moduli spaces of elliptic surfaces with sections, and prove Theorem 1.1(1-2). In Section 6, we give a more direct proof in the case when $n = 3$ by computing explicitly the deformation theory of such surfaces. Finally, in Section 7, we introduce twisted stable maps, prove Theorem 1.1(3), and develop a combinatorial algorithm for computing the KSB-stable limits.

Acknowledgement. This project was initiated during the AGNES Summer School on Higher Dimensional Moduli at Brown University in 2022. We are grateful to the organizers as well as all the speakers. During the preparation of this article, AFH and GI were partially supported by an AMS-Simons travel grant. DB is partially supported by NSF grant DMS-2401483. SM was a visiting researcher at Universität Duisburg-Essen in 2023–2024 and would like to thank it and Marc Levine for the hospitality.

2. PRELIMINARIES ON STABLE PAIRS

In this article, we work over an algebraically closed field k of characteristic 0. The reader may assume that $k = \mathbb{C}$.

2.1. Background definitions. We recall a few definitions that will be useful for the rest of the paper.

Notation 2.1. Consider a flat and finite type morphism $g: X \rightarrow Y$ of relative dimension n . If the fibers of g are Gorenstein, then the relative dualizing complex is isomorphic to a complex of the form $\omega_g[n]$ where ω_g is a line bundle. If the fibers of $g: X \rightarrow Y$ are normal, then we have an explicit description:

$$\omega_g = \left(\Lambda^n \Omega_{X/Y}^1 \right)^{**},$$

where $\Omega_{X/Y}^1$ is the sheaf of relative differentials. Indeed, both are reflexive sheaves: ω_g is a line bundle so it is reflexive, while $(\Lambda^n \Omega_{X/Y}^1)^{**}$ is reflexive by virtue of being a dual. By flat base change for dualizing complexes, they agree on the smooth locus of g , which has codimension at least 2 along each fiber, so they are isomorphic by Hartogs's lemma.

We now introduce a few standard definitions from birational geometry [KM98, Notation 0.4]. We refer the reader to [KM98] for a more extensive treatment of what follows.

Notation 2.2. Given a normal variety X with two Weil divisors D_1, D_2 , we write $D_1 \sim D_2$ to denote that D_1 and D_2 are linearly equivalent.

Definition 2.3. Let X be a normal variety. A \mathbb{Q} -divisor $D = \sum a_i D_i$ on X is a formal linear combination of integral Weil divisors $D_i \subseteq X$ with rational coefficients $a_i \in \mathbb{Q}$.

Definition 2.4. Let X be a normal variety, D a Weil divisor on X , and $\pi: X' \rightarrow X$ a birational morphism. We use the same notation as [KM98, Notation 0.4 (11)] and write $\pi_*^{-1}(D)$ for the *proper transform* of D . When π^{-1} is defined on a dense open $D_0 \subset D$, Then $\pi_*^{-1}(D)$ is the closure of $\pi^{-1}(D_0)$. The definition is extended to \mathbb{Q} -divisors by linearity.

Definition 2.5 ([KM98, Def. 2.11]). A normal variety X is said to have *canonical singularities* if

- (1) there exists $n > 0$ such that nK_X is a Cartier divisor, in other words, K_X is \mathbb{Q} -Cartier, and
- (2) for some resolution of singularities $\pi: X' \rightarrow X$, we have $\pi_*(\mathcal{O}_{X'}(nK_{X'})) \cong \mathcal{O}_X(nK_X)$.

Definition 2.6 ([Kol23, Def. 11.5]). Let X be a normal projective variety, and $D = \sum a_i D_i$ be an effective \mathbb{Q} -divisor. Let $i: X^{\text{sm}} \hookrightarrow X$ be the inclusion of the smooth locus of X . Then (X, D) is called a *log pair* if there is an integer $n > 0$ such that the sheaf

$$i_* \left(\omega_{X^{\text{sm}}}^{\otimes n}(nD|_{X^{\text{sm}}}) \right)$$

is a line bundle on X . We will denote the corresponding Cartier divisor by $n(K_X + D)$ and call $K_X + D$ \mathbb{Q} -Cartier. We say that a log pair (X, D) is *Kawamata log terminal* or *klt* (resp. *log-canonical* or *lc*) if the following conditions are satisfied:

- (1) the coefficients a_i satisfy $0 \leq a_i < 1$ (resp. $0 \leq a_i \leq 1$), and
- (2) given a log-resolution $\pi: X' \rightarrow X$, denote by E_j the integral components of the exceptional divisor. Assume that π is such that the union of the supports of $\pi_*^{-1}D$ and $\sum E_j$ is simple normal crossing. Then we require that $b_j > -n$ (resp. $b_j \geq -n$) for every j , where the $b_j \in \mathbb{Z}$ are defined by

$$(1) \quad n(K_{X'} + \pi_*^{-1}D) \sim \pi^*(n(K_X + D)) + \sum b_j E_j.$$

Remark 2.7. Often one divides Eq. (1) in Definition 2.6 by n and deals with the \mathbb{Q} -Cartier divisor $K_X + D$ instead. Moreover, if k has positive characteristic, resolution of singularities is not known, and so the definition of klt or lc singularities is slightly different from the one given above. We refer the reader to [KM98, Def. 2.28, 2.34] for more precise definitions.

Definition 2.8. Let X be a projective variety and D be an effective \mathbb{Q} -divisor. Then (X, D) is called a *semi-log-canonical pair* (or an *slc pair*) if X is equidimensional and S_2 , all points of codimension 1 are either smooth or nodal, the irreducible components of D intersect the smooth locus of X , and the following hold.

- (1) Denote by $U \subseteq X$ the locus where X is Gorenstein and D is Cartier. We then require that, for some $n > 0$, the sheaf $i_* (\omega_U^{\otimes n} \otimes \mathcal{O}_U(nD|_U))$ is a line bundle on X . We denote the corresponding Cartier divisor by $n(K_X + D)$ and say that $K_X + D$ is \mathbb{Q} -Cartier.
- (2) If $\nu: X^\nu \rightarrow X$ is the normalization of X , $\Delta \subseteq X^\nu$ is the preimage of the nodal locus, and D^ν is the proper transform of D in X^ν , then the pair $(X^\nu, D^\nu + \Delta)$ is log canonical.

Definition 2.9. Let (X, D) be a pair such that K_X and D are both \mathbb{Q} -Cartier and X is log-canonical. We define the *log-canonical threshold* of (X, D) , denoted by $\text{lct}(X; D)$, to be

$$\sup\{\alpha \in \mathbb{R} : (X, \alpha D) \text{ is log-canonical}\}.$$

2.2. Canonical models.

Definition 2.10. Let (X, D) be an lc pair and assume that X is a surface. A *canonical model* of (X, D) is a birational map $\pi: X \rightarrow X'$ to a variety X' such that, if we write $D' := \pi_* D$, then:

- (1) (X', D') is an lc pair,
- (2) there is $n > 0$ such that $n(K_{X'} + D')$ is ample, and
- (3) given m such that both $m(K_X + D)$ and $m(K_{X'} + D')$ are Cartier, if we write

$$m(K_X + D) - \pi^*(m(K_{X'} + D')) \sim \sum a_i E_i,$$

where E_i are the components of the exceptional divisor of π , then $a_i \geq 0$.

Remark 2.11. The definition of canonical model extends to higher dimensional varieties. However, one can no longer assume that π is a morphism, rather it has to be a birational rational map whose inverse does not contract divisors. In this manuscript, we will only need the surface version, so we refer the reader to [KM98, §3.8] for more details on the higher dimensional case.

Theorem 2.12 (cf. [KM98, Thm. 3.52]). *Let (X, D) be an lc pair with a proper X . Fix m such that $m(K_X + D)$ is Cartier. Then a canonical model (X', D') , if exists, is unique up to isomorphism and is given by*

$$X' = \text{Proj} \left(\bigoplus_{r \geq 0} H^0 \left(X, \mathcal{O}_X(rm(K_X + D)) \right) \right).$$

Definition 2.13. More generally, if (X, D) is a normal but not necessarily log canonical pair, we can define the *canonical model* of (X, D) as follows. Let $\mu: X' \rightarrow X$ be a log resolution with reduced exceptional divisor E . By assumption $\text{Supp}(\mu_*^{-1}D) \cup E$ is normal crossings, so the pair $(X, \mu_*^{-1}D + E)$ is log canonical and has a well-defined canonical model. We define this to be the canonical model of (X, D) . One can check by the above theorem that this is independent of a choice of μ .

Definition 2.14. A pair (X, D) is called *KSBA-stable* if

- (1) it is slc and X is connected;
- (2) $K_X + D$ is an ample \mathbb{Q} -Cartier \mathbb{Q} -divisor.

The *volume* of a pair (X, D) is $\text{vol}(X, D) = (K_X + D)^{\dim X}$.

Theorem 2.15 (cf. [Kol23, Theorem 8.15]). *Fix $v \in \mathbb{Q}_{>0}$ and $c \in [0, 1]_{\mathbb{Q}}$. Then there is a proper Deligne-Mumford stack $\mathcal{M}_{c,v}^{\text{KSBA}}$, whose closed points parametrize KSBA-stable surface pairs (X, cD) with volume $(K_X + D)^2 = v$, where D is an integral effective divisor.*

The complete definition of a family of KSBA-stable pairs is more subtle and we omit it. For this paper, the following remark suffices, and it follows from Kollár's definition [Kol23, 8.13] and [Kol23, Theorem 5.4], noting that relative ampleness can be defined fiberwise for finite type morphisms [Gro66, 9.6.4].

Remark 2.16. Assume that $\mathcal{X} \rightarrow B$ is a flat, pure dimensional, Gorenstein and projective morphism with B reduced, and $\mathcal{D} \subseteq \mathcal{X}$ is a Cartier divisor, flat over B , and such that for every $b \in B$ the fiber $(\mathcal{X}_b, c\mathcal{D}_b)$ is a stable pair with $(K_{\mathcal{X}_b} + c\mathcal{D}_b)^{\dim \mathcal{X}_b} = v$ for some rational coefficient c . Then $\mathcal{X} \rightarrow B$ is a family of KSBA-stable pairs: namely, it corresponds to a morphism $B \rightarrow \mathcal{M}_{c,v}^{\text{KSBA}}$.

Definition 2.17. A connected projective variety X is *KSB-stable* if $(X, 0)$ is KSBA-stable, i.e., X is slc and K_X is ample. Its volume is defined as $\text{vol}(X) := (K_X)^{\dim X}$.

Theorem 2.15 is still applicable here as the special case when $D = 0$, and we get a moduli space which we denote $\mathcal{M}_v^{\text{KSB}}$. Through out this paper, when we refer to $\mathcal{M}_v^{\text{KSB}}$, we take $v = \frac{(n-2)^2}{n}$ to be the volume of a pseudo-elliptic surface of with a $\frac{1}{n}(1, 1)$ -singularity, where $n \geq 3$.

2.3. KSB-stable families. In this subsection, we introduce the notion of KSB-stable family.

Notation 2.18. Let $X \rightarrow B$ be a flat family with S_2 fibers. For any reflexive sheaf \mathcal{L} on X which is a line bundle over a dense open subset, we set $\mathcal{L}^{[m]} := (\mathcal{L}^{\otimes m})^{**}$ to be the reflexive hull of the m^{th} tensor power.

Definition 2.19. Let $\pi: X \rightarrow B$ be a flat and proper morphism. We say that π is *KSB-stable* if for every $b \in B$, the fiber X_b is KSB-stable, and π satisfies *Kollár's condition*: For every $B' \rightarrow B$ and for every $n \in \mathbb{Z}$, if we denote by $p_1: X \times_B B' \rightarrow X$ the first projection, then the natural map

$$p_1^*(\omega_{X/B}^{[n]}) \longrightarrow \omega_{X \times_B B'/B'}^{[n]}$$

is an isomorphism.

It was proven in [Kol09a] that Kollár's condition is an algebraic condition, i.e, the corresponding subfunctor is represented by a monomorphism of schemes. When the base B is reduced, then Kollár's condition can be checked numerically as follows.

Theorem 2.20 (cf. [Kol23, Thm. 5.1]). *Let $\pi: X \rightarrow B$ be a flat, pure dimensional, proper morphism over a reduced base scheme B . Suppose that for all $b \in B$, the fiber X_b is KSB-stable. If the volume $(K_{X_b})^{\dim X_b}$ is the same constant value for all $b \in B$, then π is KSB-stable.*

We now introduce an auxiliary tool, developed in [Hac04, AH11], which we will use to study the infinitesimal structure of \mathcal{M}^{KSB} .

Definition 2.21. The *covering stack* of a KSB-stable family $X \rightarrow B$ is the quotient stack

$$\mathcal{X} := \left[\text{Spec}_X \left(\bigoplus_{m \in \mathbb{Z}} \omega_{X/B}^{[m]} \right) / \mathbb{G}_m \right].$$

Note that the natural map $\pi: \mathcal{X} \rightarrow X$ is an isomorphism on the open set $U \subset X$ where $\omega_{X/B}$ is a line bundle.

The main relevance of Definition 2.21 lies in the following result.

Theorem 2.22 (cf. [AH11, Thm. 5.3.6]). *Let $X \rightarrow B$ be a KSB-stable family. Then its covering stack $\mathcal{X} \rightarrow B$ is a flat and proper family of Deligne-Mumford stacks, and \mathcal{X}_b is the covering stack of X_b for every $b \in B$. Conversely, if $\mathcal{X} \rightarrow B$ is a flat and proper morphism, whose fibers \mathcal{X}_b are the covering stacks of a KSB-stable variety, then there is a KSB-stable family $X \rightarrow B$ whose covering stack is $\mathcal{X} \rightarrow B$.*

In particular, if X is KSB-stable, to study the local structure of $\mathcal{M}_b^{\text{KSB}}$ around the point p corresponding to X , one can either study flat deformations of X which satisfy Kollár's condition, or, equivalently, study flat deformations of the covering stack of \mathcal{X} .

3. ELLIPTIC SURFACES

In this part, we will recall a few facts about elliptic fibrations from [Mir89] that are needed in the rest of the paper. Throughout this section, let C be a smooth curve.

Definition 3.1. An *elliptic surface* over C is a pair $(g: X \rightarrow C, S)$ where

- (1) $g: X \rightarrow C$ is a flat proper morphism with connected fibers of arithmetic genus 1 such that the generic fiber of g is smooth, and
- (2) $S \subset X$ is a section.

The elliptic surface is *standard* if $S \subset X$ does not pass through a singular point of any fiber. A standard elliptic surface $(f: X \rightarrow C, S)$ is *minimal* if X is smooth and there are no (-1) -curves contracted by g .

Given a standard elliptic surface $(f: Y \rightarrow C, S_Y)$, one can always contract every fiber component not meeting S_Y to obtain a standard elliptic surface $(g: X \rightarrow C, S)$ with integral fibers, called the *Weierstrass model*.

Definition 3.2 ([Mir89, II.3.2]). A *Weierstrass fibration* is a standard elliptic surface whose geometric fibers are integral.

The condition of integrality on the fiber implies that X admits a *global Weierstrass equation*. That is, X is isomorphic to

$$(2) \quad \{y^2z - z^3 - Axz^2 - Bz^3 = 0\} \subseteq \mathbb{P}(\mathcal{O}_C \oplus \mathcal{L}^{-2} \oplus \mathcal{L}^{-3}),$$

where \mathcal{L} is an effective line bundle on C and $A \in H^0(C, \mathcal{L}^4)$, $B \in H^0(C, \mathcal{L}^6)$ (see [Mir89, II.5.1, II.5.2]). In particular, the fibers of g are either a smooth elliptic curve, a rational curve with a node, or a rational curve with a cusp. The line bundle \mathcal{L} is called the *fundamental line bundle* of the fibration, and determines the canonical bundle of X as follows.

Theorem 3.3 (Kodaira's canonical bundle formula [BPVdV84, V.12.1], [Mir89, Prop. III.1.1]). *Let $(g: X \rightarrow C, S)$ be a Weierstrass fibration, and let \mathcal{L} be the fundamental line bundle. Then we have*

$$\omega_X \cong g^*(\omega_C \otimes \mathcal{L}).$$

3.1. Minimal Weierstrass fibrations. We will be particularly interested in the case of canonical surface singularities (also called *du Val* or *ADE* singularities). See [KM98, §4.2], especially [KM98, Thm. 4.20], for a description.

Definition 3.4. Let $(X \rightarrow C, S)$ be a Weierstrass fibration. We say it is a *minimal Weierstrass fibration* if X has canonical singularities. We say it is an *lc Weierstrass fibration* if X has log-canonical singularities, i.e. $(X, 0)$ is lc.

The naming comes from the well-known fact that $(X \rightarrow C, S)$ is a minimal Weierstrass fibration if and only if the minimal resolution of X is a minimal elliptic surface [Mir89, Def. III.3.1, Prop. III.3.2]. In particular, the minimal Weierstrass model is unique among birational models of the elliptic fibration since the minimal model of surfaces is unique.

Remark 3.5 (cf. [Mir89, III.3.2]). In terms of the Weierstrass data (\mathcal{L}, A, B) , the Weierstrass fibration is minimal (resp. lc) if and only if for all $p \in C$:

$$\min \{3\text{ord}_p(A), 2\text{ord}_p(B)\} \underset{(-)}{<} 12.$$

Construction 3.6. We now describe a birational transformation that, given a minimal Weierstrass fibration $(g: X \rightarrow C, S)$, produces a strictly lc Weierstrass fibration $(g': X' \rightarrow C, S')$, i.e. one which is not minimal. First, consider a fiber F of g over $p \in C$, and let X' be the blow-up of X at the intersection point $F \cap S$. Let F', S' be the proper transforms of F, S , respectively, and let E denote the exceptional divisor, then the fiber of $X' \rightarrow C$ over p is $F' \cup E$. Assume that $(X, S + F)$ is an lc pair, that is, F was chosen either smooth or nodal. One can check that $K_{X'} + S' + F'$ is nef over C by checking its intersections with F' and E' are nonnegative (0 and 1, respectively). The canonical model of $(X', S' + F')$ over C contracts only F' , and the resulting surface pair (X^c, S^c) is strictly log-canonical, as F' is an exceptional divisor with discrepancy -1 . In particular, this process generates strictly lc singularities.

Recall that \mathcal{L} is the dual of the normal bundle of the section, so this process corresponds to replacing \mathcal{L} with $\mathcal{L} \otimes \mathcal{O}_C(p)$. Similarly, we are replacing (A, B) with $(z^4 A, z^6 B)$, where z is a nonzero section of $\mathcal{O}_C(p)$ vanishing at p .

Remark 3.7. The singular fibers of a minimal Weierstrass fibration are classified in terms of the dual graph of a minimal resolution, which can be determined from the vanishing order of the Weierstrass data via Tate's algorithm. The classification of singular fibers is due to Kodaira and Nerón, and we use Kodaira's notation here. We refer the reader to [Mir89, Section I.4] and [SS10, Page 66] for more details.

Remark 3.8. Let $g: X \rightarrow \mathbb{P}^1$ be a Weierstrass fibration with section $S \subseteq X$. It follows from [Mir89, II.5.6] that $S^2 \leq 0$.

When S is rational and $S^2 = -n$, it follows from the adjunction formula that $\omega_X|_S \simeq \mathcal{O}_{\mathbb{P}^1}(n-2)$. Moreover, ω_g is trivial on the fibers, so by the see-saw lemma we can write $\omega_g = g^* \mathcal{O}_{\mathbb{P}^1}(k)$ for some k , and we have

$$\omega_X = g^* \omega_{\mathbb{P}^1} \otimes \omega_g = g^* \mathcal{O}_{\mathbb{P}^1}(-2) \otimes g^* \mathcal{O}_{\mathbb{P}^1}(k).$$

We can combine these two observations to deduce that $\omega_X \simeq g^* \mathcal{O}_{\mathbb{P}^1}(n-2)$ and $\omega_g \simeq g^* \mathcal{O}_{\mathbb{P}^1}(n)$. The canonical bundle formula (Theorem 3.3) implies that $\mathcal{L} \cong \mathcal{O}_{\mathbb{P}^1}(n)$ and $\omega_g \cong g^* \mathcal{L}$.

Definition 3.9. Given an irreducible elliptic surface $(X, S) \rightarrow \mathbb{P}^1$ with fibers that are irreducible genus one curves, we define the *height* of $(X, S) \rightarrow \mathbb{P}^1$ to be $-(S)^2$.

Remark 3.10. The height $n = 0$ if and only if $X = E \times C$ for an elliptic curve E [Mir89, Lemma III.1.4].

3.2. Twisted fibers. Not all elliptic surfaces are standard. For example, one can start with a Weierstrass fibration $X \rightarrow C$ as in Construction 3.6, but then fix a cuspidal fiber F instead of a smooth or nodal one, and section S . One can then consider the minimal log-resolution $(X', S' + F' + E) \rightarrow (X, S + F)$, where S', F' are proper transforms of S, F , respectively, and E is the reduced exceptional divisor. Now we may assume the pair $(X', S' + F' + E)$ is SNC, so one can construct its relative canonical model $(X^c, S^c + F^c)$ over C as in [AB17]. It is proven in *loc. cit.* that the new pair (X^c, S^c) is not a standard elliptic surface, namely, X^c is singular at $F^c \cap S^c$. However, the singular fibers arising in this process are controlled.

Definition 3.11 ([AB17, Def. 4.9]). Let $g: (X, S) \rightarrow C$ be a relative canonical model of an elliptic surface with section. A *twisted fiber* of g is an irreducible but non-reduced fiber.

Twisted fibers also appear as follows. Consider a Weierstrass fibration $(X, S) \rightarrow C$. Let $U \subseteq C$ be the locus where the fibers of g are either smooth or nodal. Assume that $U \neq C$. There is a map $\phi: U \rightarrow \overline{\mathcal{M}}_{1,1}$. Up to replacing C by some *root stack* $\gamma: \mathcal{C} \rightarrow C$ of C , with $U \subseteq C$ the open locus over which γ is an isomorphism, we can extend ϕ to $\Phi: \mathcal{C} \rightarrow \overline{\mathcal{M}}_{1,1}$ (cf. [AB19, BPS22, BV23]). Consider the pull-back of the universal family via Φ , denoted by $(\mathcal{X}, \mathcal{S}) \rightarrow \mathcal{C}$, and let $g': (X', S') \rightarrow C$ be the corresponding map on coarse

moduli spaces. From [AB17] all the fibers of g' over $C \setminus U$ are twisted, so this procedure replaces the fibers of g away from U with twisted fibers.

In fact, these two constructions of twisted fibers are equivalent.

Proposition 3.12 ([AB19, Prop. 4.12]). *Let $(g: X \rightarrow C, S)$ be a Weierstrass fibration with a cuspidal fiber F . Then the relative canonical model $(X^c, S^c + F^c)$ of $(X, S + F)$ over C is the twisted model (X', S') obtained from the root stack construction above.*

3.3. Stability for Weierstrass fibrations. For the remainder of this section, we assume that $g: X \rightarrow \mathbb{P}^1$ is an lc Weierstrass fibration with section S , fiber class f , and height $-S^2 = n$.

Lemma 3.13. *Let $(X \rightarrow \mathbb{P}^1, S)$ be an lc Weierstrass fibration as above with height n . Then the following holds for the pair (X, cS) .*

- (1) *When $n \geq 3$, the pair is KSBA-stable if and only if $0 < c < \frac{n-2}{n}$.*
- (2) *Its volume is $\text{vol}(X, cS) = 2c(n-2) - nc^2$.*

Proof. Assume first that $0 < c < \frac{n-2}{n}$. Since X has lc singularities, S is smooth and contained in the smooth locus of X , and $c < 1$, we conclude that (X, cS) is automatically lc, hence slc. As its volume is

$$(K_X + cS)^2 = ((n-2)f + cS)^2 = 2c(n-2) - c^2n,$$

and the inequalities on c imply that $(K_X + cS)^2 > 0$. Now to check that $K_X + cS$ is ample, it suffices to show that $K_X + cS$ intersects every irreducible curve D positively. Since $g: X \rightarrow \mathbb{P}^1$ has irreducible fibers and $(K_X + cS).f = c > 0$, we may assume that D is not a fiber. Then in particular we have $(D.f) \geq 1$. Thus, it follows that

$$(K_X + cS).D \geq n - 2 + c \cdot (S.D) \geq n - 2 - cn > 0,$$

for $0 < c < \frac{n-2}{n}$. Here we have used that $(S.D) \geq 0$ if $D \neq S$ and $(S.D) = -n$ if $S = D$. The converse statement in part (1) is clear from the formula for $(K_X + cS)^2$. \square

Observation 3.14. *Consider a pair (X, cS) as in Lemma 3.13. When $c = \frac{n-2}{n}$, we have that $(K_X + cS).S = 0$. One can check that the canonical model of $(X, \frac{n-2}{n}S)$ is the surface Y obtained by contracting the section S to a point. If we denote by $\pi: X \rightarrow Y$ such a contraction, then*

$$\pi^*K_Y = K_X + \frac{n-2}{n}S,$$

and thus the volume of Y is

$$\text{vol}(K_Y) = 2 \cdot \frac{(n-2)}{n} \cdot (n-2) - n \cdot \frac{(n-2)^2}{n^2} = \frac{(n-2)^2}{n}.$$

Remark 3.15. In [LN02, AB21], the surfaces obtained from a Weierstrass fibration by contracting the section are also called *pseudo-elliptic surfaces*.

4. MODULI SPACES OF ELLIPTIC SURFACES

In this section, we introduce the three moduli spaces of elliptic surfaces we will be dealing with, and we will present some of the relations between them.

4.1. Weierstrass fibrations. We begin by recalling a description for the moduli stack of Weierstrass fibrations.

Definition 4.1. Fix $n \geq 0$. We define the stack \mathcal{W}_n^{\min} as the pseudofunctor

$$\mathcal{W}_n^{\min}(B) := \left\{ \begin{array}{c} \begin{array}{ccc} S \subset X & \xrightarrow{f} & C \\ & \searrow & \swarrow \\ & B & \end{array} \\ \left. \begin{array}{l} (1) f: X \rightarrow C \text{ is flat projective of relative dimension 1;} \\ (2) C \rightarrow B \text{ is flat projective with geometric fibers } \mathbb{P}^1; \\ (3) S \subset X \text{ is a section of } f; \\ (4) \text{ for each } b \in B, \text{ the fiber } f_b: (X_b, S_b) \rightarrow C_b \text{ is a} \\ \text{minimal Weierstrass fibration of height } -(S_b)^2 = n. \end{array} \right\}.$$

Similarly, we let $\mathcal{W}_n^{\text{lc}}$ be the pseudofunctor of families as above, where we allow $(f_b: X_b \rightarrow C_b, S_b)$ to be an lc Weierstrass fibration for all $b \in B$.

In the definition above, $C \rightarrow B$ is a family of one-dimensional Brauer-Severi varieties and, in particular, corresponds to a PGL_2 -torsor over B .

Theorem 4.2. (cf. [PS21, Main Theorem 1.2] & [CDLI24, Thm. 2.8]) *Keep the notations as above.*

- (1) *The pseudofunctors $\mathcal{W}_n^{\text{min}}$ and $\mathcal{W}_n^{\text{lc}}$ are representable by smooth and irreducible Deligne-Mumford stacks of finite type. Moreover, $\mathcal{W}_n^{\text{min}}$ is an open substack of $\mathcal{W}_n^{\text{lc}}$*
- (2) *There are nonempty open loci $\mathcal{W}_n^{\text{reg}}, \mathcal{W}_n^{\text{nod}} \subset \mathcal{W}_n^{\text{min}}$ parametrizing those Weierstrass fibrations such that X is smooth or that have at worst nodal fibers respectively.*

Proof. The construction of $\mathcal{W}_n^{\text{min}}$ as a coarse moduli space is classical (cf. [Mir81, Sei87]) and the modern stacky perspective is given in [PS21, §4] and [CDLI24, §2]. We sketch here the irreducibility, smoothness and quasicompactness of the stacks. We refer the interested reader to the references (e.g. [PS21, Thm. 1.2(b)]) for the fact that the stacks are Deligne-Mumford.

Consider the pseudofunctor $\widetilde{\mathcal{W}}_n^{\text{lc}}$ whose B -points consist of tuples

$$(f: X \rightarrow C, S, \alpha: C \cong \mathbb{P}_B^1),$$

where $(f: X \rightarrow C, S)$ is as above and α is an isomorphism of C with \mathbb{P}_B^1 . We call α a *framing*. There is a PGL_2 -action on $\widetilde{\mathcal{W}}_n^{\text{lc}}$, which precomposes the framing with an automorphism of \mathbb{P}_B^1 , such that

$$\mathcal{W}_n^{\text{lc}} \cong [\widetilde{\mathcal{W}}_n^{\text{lc}} / \text{PGL}_2].$$

Thus, it suffices to show that $\widetilde{\mathcal{W}}_n^{\text{lc}}$ is representable by a smooth Deligne-Mumford stack and that the corresponding subfunctors

$$\widetilde{\mathcal{W}}_n^{\text{reg}} \subseteq \widetilde{\mathcal{W}}_n^{\text{nod}} \subseteq \widetilde{\mathcal{W}}_n^{\text{min}} \subseteq \widetilde{\mathcal{W}}_n^{\text{lc}}$$

are all open. Let $V_m := H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(m))$ be the space of degree m homogeneous polynomials and define

$$\mathbb{V}_n := V_{4n} \oplus V_{6n}.$$

Two Weierstrass equations give isomorphic surfaces over \mathbb{P}^1 if and only if they differ by the \mathbb{G}_m -action with weights $(4, 6)$ on \mathbb{V}_n . Thus there is a natural morphism

$$\varphi: \widetilde{\mathcal{W}}_n^{\text{lc}} \rightarrow [\mathbb{V}_n / \mathbb{G}_m]$$

which sends a framed lc Weierstrass fibration $(f: X \rightarrow \mathbb{P}^1, S)$ of height n to the coefficients of its Weierstrass equation $(A, B) \in [\mathbb{V}_n / \mathbb{G}_m]$ as in Eq. (2). See [PS21, §4] for a description of this construction in families. The image of φ is the open substack \mathcal{U}^{lc} defined by the condition

$$(3) \quad \min \{3 \cdot \text{ord}_x(A), 2 \cdot \text{ord}_x(B)\} \leq 12$$

for any $x \in \mathbb{P}^1$. Similarly, one has that

- (1) the image of $\widetilde{\mathcal{W}}_n^{\text{min}}$ is the open substack $\mathcal{U} \subset \mathcal{U}^{\text{lc}}$ defined by a strict inequality in (3),
- (2) the image of $\widetilde{\mathcal{W}}_n^{\text{nod}}$ is the open locus where the vanishing of A and the vanishing of B are disjoint,
- (3) and the image of $\widetilde{\mathcal{W}}_n^{\text{reg}}$ is open because smoothness is open in flat families (cf. [Har13, Chap. III.10]).

The construction, which sends a family of Weierstrass data (A, B) to the surface

$$X = \mathbb{V}(Y^2Z - (X^3 + AXZ^2 + BZ^3)) \subseteq \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{L}^{-2} \oplus \mathcal{L}^{-3}) \rightarrow \mathbb{P}^1$$

with the section $S = \mathbb{V}(X, Z)$, defines an inverse $\mathcal{U}^{\text{lc}} \rightarrow \widetilde{\mathcal{W}}_n^{\text{lc}}$, which maps \mathcal{U} isomorphically onto $\widetilde{\mathcal{W}}_n^{\text{min}}$.

Finally, it follows from this construction that $\mathcal{W}_n^{\text{min}}$ and $\mathcal{W}_n^{\text{lc}}$ are smooth, since they are given as quotients of open substacks of \mathbb{V}_n . \square

Corollary 4.3. *Let $(f: X \rightarrow C, S) \rightarrow B$ be a family of lc Weierstrass fibrations. Then there exists a closed subscheme*

$$\Delta(f)^{\text{cusp}} \subseteq C$$

such that for each $b \in B$, $x \in \Delta(f)_b^{\text{cusp}}$ if and only if the fiber $f_b^{-1}(x)$ is cuspidal. Moreover, the formation of $\Delta(f)^{\text{cusp}}$ commutes with base change.

Proof. It suffices to define the closed substack $\Delta^{\text{cusp}} \subseteq \mathcal{C}^{\text{univ}} \rightarrow \mathcal{W}_n^{\text{lc}}$ on the universal family and define $\Delta(f)^{\text{cusp}}$ via pulling back. Moreover, we can do this smooth locally over $\mathcal{W}_n^{\text{lc}}$, so it suffices to work with framed Weierstrass fibrations. Given a Weierstrass fibration $(f: X \rightarrow \mathbb{P}^1, S)$ over $\text{Spec } k$, the cuspidal fibers are given by the vanishing

$$\{A = B = 0\}$$

where A and B are the Weierstrass data. Let \mathcal{A} and \mathcal{B} be the universal Weierstrass data on $\mathbb{P}_{\mathcal{W}_n^{\text{lc}}}^1$. Then the closed substack

$$\tilde{\Delta}^{\text{cusp}} = \{\mathcal{A} = \mathcal{B} = 0\} \subseteq \mathbb{P}_{\mathcal{W}_n^{\text{lc}}}^1$$

descends to a closed substack $\Delta^{\text{cusp}} \subseteq \mathcal{C}^{\text{univ}}$, which does the job. \square

Each point in $\partial\mathcal{W}_n^{\text{lc}} := \mathcal{W}_n^{\text{lc}} \setminus \mathcal{W}_n^{\text{min}}$ corresponds to a Weierstrass fibration $(f: X \rightarrow C, S)$ with some finite number $m > 0$ of cuspidal fibers, where X has elliptic singularities. The associated minimal Weierstrass model

$$(f^{\text{min}}: X^{\text{min}} \rightarrow C, S^{\text{min}})$$

has height $n - m$, and $(f: X \rightarrow C, S)$ can be recovered from X^{min} by choosing m smooth or nodal fibers F_1, \dots, F_m , blowing up $F_i \cap S^{\text{min}}$ and contracting the strict transform of F_i to an elliptic singularity as in Construction 3.6. This yields a stratification of $\mathcal{W}_n^{\text{lc}}$ which we extend to a compactification in Section 7.

Definition 4.4. Define the pseudofunctor $\mathcal{W}_{n-m,m}$ by sending a base scheme T to

$$\mathcal{W}_{n-m,m}(T) = \left\{ \begin{array}{c} \begin{array}{ccc} S \subset X & \xrightarrow{f} & C \supset D \\ & \searrow & \swarrow \\ & T & \end{array} \\ \left. \begin{array}{l} (1) (f: X \rightarrow C, S) \rightarrow T \text{ is a family of minimal Weierstrass fibrations of height } n - m; \\ (2) D \subset C \rightarrow T \text{ is a relative effective Cartier divisor of degree } m \text{ which is étale over } T; \\ (3) D \cap \Delta(f)^{\text{cusp}} = \emptyset. \end{array} \right\} \end{array} \right.$$

Note in particular that $\mathcal{W}_{n,0} = \mathcal{W}_n^{\text{min}}$.

The following result will not be used in the rest of the paper. However, it is of independent interest as it describe the structure of $\mathcal{W}_n^{\text{lc}}$.

Theorem 4.5. *The stacks $\mathcal{W}_{n-m,m}$ are smooth, separated Deligne-Mumford stacks of finite type. There exists a locally closed stratification*

$$\bigsqcup_{m=0}^n \mathcal{W}_{n-m,m} \rightarrow \mathcal{W}_n^{\text{lc}}$$

such that the image of $\mathcal{W}_{n-m,m}$ is the locus of Weierstrass fibrations of height n with exactly m strictly log canonical elliptic singularities.

Proof. Analogous to the construction of $\mathcal{W}_n^{\text{min}}$, let $\widetilde{\mathcal{W}}_{n-m,m}$ be the PGL_2 cover of $\mathcal{W}_{n-m,m}$ given by adding the data of a framing $\alpha: C \cong \mathbb{P}_T^1$ to the definition of $\mathcal{W}_{n-m,m}$. Then $\widetilde{\mathcal{W}}_{n-m,m}$ can be identified with the substack of $\widetilde{\mathcal{W}}_{n-m} \times \text{Sym}^m \mathbb{P}^1$ sending any base scheme T to the groupoid of pairs $(f: X \rightarrow \mathbb{P}_T^1, S)$ in $\widetilde{\mathcal{W}}_{m,n}^{\text{min}}(T)$ along with a degree m relative Cartier divisor $D \in \text{Sym}^m \mathbb{P}^1(T)$ that is étale over T and satisfies the condition that $\Delta(f)^{\text{cusp}} \cap D = \emptyset$. Note that the étaleness of $D \rightarrow T$ is an open condition. On the other hand, since

$$\Delta(f)^{\text{cusp}} \subseteq \mathbb{P}_T^1 \quad \text{and} \quad D \subseteq \mathbb{P}_T^1$$

are both closed and $\mathbb{P}_T^1 \rightarrow T$ is proper, then the condition that $\Delta(f)^{\text{cusp}} \cap D = \emptyset$ is open on T . Therefore, $\widetilde{\mathcal{W}}_{n-m,m}$ is an open substack of $\widetilde{\mathcal{W}}_{n-m} \times \text{Sym}^m \mathbb{P}^1$. By Theorem 4.2 for $\mathcal{W}_{n-m}^{\text{min}}$, this concludes the proof of the first claim.

Now we define natural morphisms $\widetilde{\mathcal{W}}_{n-m,m} \longrightarrow \widetilde{\mathcal{W}}_n^{\text{lc}}$ as follows. For any T -point of $\widetilde{\mathcal{W}}_{n-m,m}$, let

$$A_0 \in H^0(\mathbb{P}_T^1, L_0^{\otimes 4} \boxtimes \mathcal{O}_{\mathbb{P}^1}(4n-4m)) \quad \text{and} \quad B_0 \in H^0(\mathbb{P}_T^1, L_0^{\otimes 6} \boxtimes \mathcal{O}_{\mathbb{P}^1}(6n-6m))$$

be the minimal Weierstrass data associated to the height $n-m$ fibration $(f: X \rightarrow \mathbb{P}_T^1, S)$, where L_0 is a line bundle on T , and let $z \in H^0(\mathbb{P}_T^1, \mathcal{O}_{\mathbb{P}^1}(D))$ be a defining polynomial of D . Note that

$$\mathcal{O}_{\mathbb{P}_T^1}(D) \cong M \boxtimes \mathcal{O}_{\mathbb{P}^1}(m)$$

for some line bundle $M \in \text{Pic}(T)$ via the isomorphism $\text{Pic}(\mathbb{P}_T^1) \cong \text{Pic}(T) \times \text{Pic}(\mathbb{P}^1)$. Then we can define new Weierstrass data

$$A = A_0 z^4 \in H^0(\mathbb{P}_T^1, L^{\otimes 4} \boxtimes \mathcal{O}_{\mathbb{P}^1}(4n)) \quad \text{and} \quad B = B_0 z^6 \in H^0(\mathbb{P}_T^1, L^{\otimes 6} \boxtimes \mathcal{O}_{\mathbb{P}^1}(6n))$$

where $L = L_0 \otimes M$. By assumption z has distinct roots and $z \neq 0$ whenever $A_0 = B_0 = 0$. Thus (A, B) is minimal away from D , and strictly log canonical along D . Therefore, the Weierstrass pair (A, B) defines a family of framed lc Weierstrass fibrations over T with elliptic singularities over D by Construction 3.6.

This construction is functorial and PGL_2 -equivariant and hence defines natural morphisms

$$\varphi_m : \mathcal{W}_{n-m,m} \longrightarrow \mathcal{W}_n^{\text{lc}}$$

for each $m = 0, \dots, n$, whose image is the locus of lc Weierstrass fibrations with exactly m elliptic singularities. We claim that

$$\bigsqcup_{m=0}^n \mathcal{W}_{n-m,m} \longrightarrow \mathcal{W}_n^{\text{lc}}$$

is a locally closed stratification.

After passing to $\widetilde{\mathcal{W}}_n^{\text{lc}}$, we are reduced to checking that

$$\bigsqcup_{m=0}^n \varphi_m : \bigsqcup_{m=0}^n \widetilde{\mathcal{W}}_{n-m,m} \longrightarrow \widetilde{\mathcal{W}}_n^{\text{lc}}$$

is a locally closed stratification. It is surjective on closed points, because any lc Weierstrass data over \mathbb{P}_k^1 can be factored as $A = A_0 z^4$ and $B = B_0 z^6$, where (A_0, B_0) is minimal. Notice also that, at the level of geometric points, we have that φ_m is stabilizer preserving (the stabilizer consist of μ_2 generically, μ_4 if $B = 0$ and μ_6 if $A = 0$ for both the source and target stacks), so each φ_m is representable. Furthermore, since the factorization $A = A_0 z^4$ and $B = B_0 z^6$, where (A_0, B_0) is minimal, is unique (up to scaling) at the level of field-valued points, φ_m is injective on geometric points. We claim that the morphism of smooth stacks φ_m is unramified. To see this, it suffices to check that φ_m induces injections for tangent spaces for each geometric point.

Set $\mathbb{V}_d := H^0(\mathcal{O}_{\mathbb{P}^1}(d))$ for any given d . We note that $\widetilde{\mathcal{W}}_{n-m,m}$ may be viewed as an open substack of

$$\mathcal{N}_1 = [(\mathbb{V}_{4n-4m} \times \mathbb{V}_{6n-6m})/\mathbb{G}_m] \times [\mathbb{V}_m/\mathbb{G}_m],$$

where the first copy of \mathbb{G}_m acts with weights 4 and 6 on the vector spaces \mathbb{V}_{4n-4m} and \mathbb{V}_{6n-6m} , and the second copy of \mathbb{G}_m acts with weight 1 on the vector space \mathbb{V}_m . A k -point of \mathcal{N}_1 consists of a triple of elements (α, β, z) in $H^0(\mathcal{O}_{\mathbb{P}^1}(4n-4m)) \times H^0(\mathcal{O}_{\mathbb{P}^1}(6n-6m)) \times H^0(\mathcal{O}_{\mathbb{P}^1}(m))$, and the tangent space of the stack at that point is the cokernel of the linear morphism of k -vector spaces

$$\psi_1 : k^{\oplus 2} \longrightarrow H^0(\mathcal{O}_{\mathbb{P}^1}(4n-4m)) \oplus H^0(\mathcal{O}_{\mathbb{P}^1}(6n-6m)) \oplus H^0(\mathcal{O}_{\mathbb{P}^1}(m))$$

given by $(x, y) \mapsto (4x\alpha, 6x\beta, yz)$. On the other hand, the stack $\widetilde{\mathcal{W}}_n^{\text{lc}}$ can be viewed as an open substack of

$$\mathcal{N}_2 = [(\mathbb{V}_{4n} \times \mathbb{V}_{6n})/\mathbb{G}_m],$$

where \mathbb{G}_m acts with weights 4 and 6 on the vector spaces \mathbb{V}_{4n} and \mathbb{V}_{6n} . A k -point of \mathcal{N}_2 is given by a pair $(\delta, \gamma) \in (H^0(\mathcal{O}_{\mathbb{P}^1}(4n)) \times H^0(\mathcal{O}_{\mathbb{P}^1}(6n)))$, and the corresponding tangent space is the cokernel of

$$\psi_2 : k \rightarrow H^0(\mathcal{O}_{\mathbb{P}^1}(4n)) \oplus H^0(\mathcal{O}_{\mathbb{P}^1}(6n))$$

given by $x \mapsto (4x\delta, 6x\gamma)$. We note that the morphism φ_m extends to a morphism $\tilde{\varphi}_m$ that sends a k -point (α, β, z) to $(\alpha z^4, \beta z^6)$. A direct computation shows that the induced morphism of tangent complexes at the k -point (α, β, z) is given by

$$\begin{array}{ccc} k^{\oplus 2} & \xrightarrow{\psi_1} & \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(4n-4m)) \oplus \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(6n-6m)) \oplus \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(m)) \\ \downarrow \mu & & \downarrow \nu \\ k & \xrightarrow{\psi_2} & \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(4n)) \oplus \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^1}(6n)) \end{array}$$

where μ is the surjective morphism $\mu(x, y) = x + y$, and we have $\nu(a, b, c) = (z^4 \cdot a + 4\alpha z^3 \cdot c, z^6 \cdot b + 6\beta z^5 \cdot c)$. Notice that, if the k -point (α, β, z) is in the open substack $\tilde{\mathcal{W}}_{n-m, m}$, then we must have $(\alpha, \beta) \neq (0, 0)$, $z \neq 0$, and furthermore the three sections α, β, z don't have a joint simultaneous zero in \mathbb{P}^1 (because the divisor cut out by $z = 0$ does not intersect the locus of cusps, which is the locus cut out by $\alpha, \beta = 0$). It follows then from the description above that the kernel of ν is given by the subspace of (a, b, c) satisfying the linear equations $z \cdot a + 4\alpha \cdot c = 0$ and $z \cdot b + 6\beta \cdot c = 0$.

Claim: If (a, b, c) is in the kernel of ν , then it is of the form $(4x\alpha, 6x\beta, -xz)$ for some constant $x \in k$.

Let us prove the claim. We may change coordinates in \mathbb{P}^1 so that none of α, β, z vanish at ∞ , and then we view α, β, z as elements of $k[t]$. We may assume furthermore for the sake of the following argument that α, β, z are monic. Set $g_1 := \gcd(\alpha, z)$ and $g_2 = \gcd(\beta, z)$. The solutions to the equation $z \cdot a + 4\alpha \cdot c = 0$ are of the form $(a, c) = (xp \cdot 4\alpha/g_1, -xp \cdot z/g_1)$, where $x \in k$ and p is a monic polynomial. Similarly, any solution of $z \cdot b + 6\beta \cdot c = 0$ is of the form $(b, c) = (yq(6\beta/g_2), -yq(z/g_2))$, where $y \in k$ and q is a monic polynomial. If we want (a, b, c) to be a solution to both equations, then we have $-xp(z/g_1) = c = -yq(z/g_2)$, which, in view of the polynomials being monic, implies $x = y$. Therefore, we have $-xp(z/g_1) = c = -xq(z/g_2)$, which implies that $pg_2 = qg_1$. Now, the polynomials g_1 and g_2 cannot have a common root, because that would yield a simultaneous root of α, β and z , contradicting our assumptions. Therefore g_1 and g_2 are coprime, and the equation $pg_2 = qg_1$ forces $p = g_1$ and $q = g_2$. We conclude that $(a, b, c) = (4x\alpha, 6x\beta, -xz)$, as claimed.

Now, observe that any element of the kernel $(4x\alpha, 6x\beta, -xz)$ is of the form $\psi_1(x, -x)$. From this we conclude that the induced morphism on tangent spaces $\mathrm{Coker}(\psi_1) \rightarrow \mathrm{Coker}(\psi_2)$ is injective, and therefore φ_m is unramified. By [Sta24, Tag 05VH], we conclude that φ_m is a monomorphism. To complete the proof, we will apply the valuative criterion for locally closed embeddings (cf. [Kol09a, Prop. 42]).

Let $T = \mathrm{Spec} R$ be the spectrum of a DVR and let $T \rightarrow \tilde{\mathcal{W}}_n^{\mathrm{lc}}$ a map whose image is contained in $\varphi_m(\tilde{\mathcal{W}}_{n-m, m})$. This is equivalent to a family of lc Weierstrass data (A, B) over T such that for each $t \in T$, the pair (A_t, B_t) defines a fibration with exactly m elliptic singularities. Let $\eta \in T$ denote the generic point. There is a canonical subscheme D_A of the generic fiber \mathbb{P}_η^1 whose support consists of the locus of points where the section A_η has a zero of multiplicity ≥ 4 (Zariski locally around every vanishing point of A_η , it is given by the vanishing of the 3^{rd} differential of the section). Similarly, there is a canonical subscheme D_B where B_η has a zero of multiplicity ≥ 6 . Consider the reduced subscheme $D_\eta := (D_A \cap D_B)^{\mathrm{red}}$ of the intersection $D_A \cap D_B$, which is a Cartier divisor on \mathbb{P}_η^1 . We denote by $D \subset \mathbb{P}_T^1$ the flat closure of the reduced subscheme D_η . By construction, there exist factorizations $A_\eta = (A_0)_\eta + 4D_\eta$ and $B_\eta = (B_0)_\eta + 6D_\eta$ for some Cartier divisors $(A_0)_\eta$ and $(B_0)_\eta$ on \mathbb{P}_η^1 . By taking flat closures in \mathbb{P}_T^1 , we get factorizations of relative Cartier divisors $A = A_0 + 4D$ and $B = B_0 + 6D$. To conclude the proof of the valuative criterion, it suffices to show that the tuple (A_0, B_0, D) yields a T -point of $\tilde{\mathcal{W}}_{n-m, m}$. This amounts to showing that D is étale over T , and that the fibers of Weierstrass fibration defined by (A_0, B_0) are minimal. By the assumption that the original Weierstrass fibration $T \rightarrow \tilde{\mathcal{W}}_n^{\mathrm{lc}}$ was log canonical, it follows that for each geometric T -fiber there is no point where the section A has a zero of order ≥ 5 and simultaneously the section B has a zero of order ≥ 7 . Therefore, from the factorizations $A = A_0 + 4D$ and $B = B_0 + 6D$ it follows that the geometric T -fibers of D are forced to be reduced, and hence D is étale over T . On the other hand, by construction we have

arranged so that at the generic point the pair $((A_0)_\eta, (B_0)_\eta)$ yields a minimal Weierstrass fibration. By our assumption that the image of η is contained in $\varphi_m(\widetilde{\mathcal{W}}_{n-m,m})$, it follows that the order of the Cartier divisor D_η is m . If we denote by $s \in T$ the special point, then the order of the special fiber D_s is also m . Since the image of s is also contained in $\varphi_k(\widetilde{\mathcal{W}}_{n-m,m})$, the factorizations $A_s = (A_0)_s + 4D_s$ and $B = (B_0)_s + 6D_s$ with D_s of degree m force $((A_0)_s, (B_0)_s)$ to define a minimal Weierstrass fibration (otherwise we would be able to factor out a further divisor $D'_s \supset D_s$, which would mean that (A_s, B_s) is actually in the image of $\varphi_{m'}$ for some $m' > m$; this would be disjoint from $\varphi_m(\widetilde{\mathcal{W}}_{n-m,m})$). We conclude that the tuple (A_0, B_0, D) yields the desired T -point of $\widetilde{\mathcal{W}}_{n-m,m}$. \square

4.2. Locus of elliptic surfaces in the KSBA moduli. Let us start by fixing some notations.

Notation 4.6. Write $c(\epsilon) := \frac{n-2}{n} - \epsilon$ and $v(\epsilon) = \frac{(n-2)^2 - (n\epsilon)^2}{n}$. Then $v(\epsilon)$ is the volume of the pair $(X, c(\epsilon)S)$ as in Lemma 3.13.

By Lemma 3.13 and Remark 2.16, for any $0 < \epsilon < \frac{n-2}{n}$, there is a morphism

$$(4) \quad \Phi_{n,\epsilon} : \mathcal{W}_n^{\text{lc}} \longrightarrow \mathcal{M}_{c(\epsilon),v(\epsilon)}^{\text{KSBA}}, \quad (X \rightarrow C, S) \mapsto (X, c(\epsilon)S),$$

which forgets the fibration structure. Note that the moduli stacks $\mathcal{M}_{c(\epsilon),v(\epsilon)}^{\text{KSBA}}$ are not isomorphic for different $c(\epsilon)$. However, we will show now that $\Phi_{n,\epsilon}$ is an open immersion for any choice of $\epsilon \in (0, \frac{n-2}{n})$. The parameter ϵ determines the target of the morphism $\Phi_{n,\epsilon}$.

Definition 4.7. We denote by $\overline{\mathcal{E}}_{n,\epsilon}$ the seminormalization of the scheme-theoretic image (cf. [Sta24, Tag 0CMH]) of $\Phi_{n,\epsilon}$.

Proposition 4.8. For any $n > 2$ and $0 < \epsilon < \frac{n-2}{n}$, the morphism $\Phi_{n,\epsilon} : \mathcal{W}_n^{\text{lc}} \rightarrow \mathcal{M}_{c(\epsilon),v(\epsilon)}^{\text{KSBA}}$ is an open immersion.

Proof. It suffices to prove that $\Phi_{n,\epsilon}$ is smooth, induces a bijection on automorphisms of geometric points, and is injective on isomorphism classes of geometric points, since this would jointly imply that it is an étale monomorphism, hence an open immersion.

$\Phi_{n,\epsilon}$ is injective. This follows from [Inc20, Lem. 3.6]. This lemma is stated only for minimal Weierstrass fibrations, however the argument works for our case of lc Weierstrass fibrations of Kodaira dimension 1.

$\Phi_{n,\epsilon}$ is representable. It suffices to check on geometric points. We need to show that if σ is an automorphism of the Weierstrass fibration $(f : X \rightarrow \mathbb{P}^1, S)$ which induces the identity on (X, S) , then it is the identity. Indeed, from the proof of injectivity above, the map f is uniquely determined from (X, S) , and since f is surjective, the only (set-theoretic) function σ which makes the following diagram commutative is the identity, as desired

$$\begin{array}{ccc} X & \xrightarrow{\text{Id}} & X \\ \downarrow f & & \downarrow f \\ \mathbb{P}^1 & \xrightarrow{\sigma} & \mathbb{P}^1. \end{array}$$

$\Phi_{n,\epsilon}$ is surjective on automorphisms of geometric points. We need to check that any automorphism of (X, S) extends to an automorphism of $(f : X \rightarrow \mathbb{P}^1, S)$, namely, to one of the fibration. But this follows since $\omega_X = f^* \mathcal{O}_{\mathbb{P}^1}(n-2)$ so the global sections of ω_X induce the morphism f , and any automorphism of X extends (by functoriality) to an automorphism of ω_X and therefore to its global sections.

$\Phi_{n,\epsilon}$ is smooth. Smoothness is smooth local on the source so it suffices to check that the composition $\widetilde{\mathcal{W}}_n^{\text{lc}} \rightarrow \mathcal{M}^{\text{KSBA}}$ is smooth. We use the infinitesimal lifting property (cf. [Sta24, Tag 0DP0], [Sta24, Tag 02HT]), which applies to our case since the source and the target are stacks locally of finite type over a field. Let $A' \rightarrow A$ be a surjection of local Artinian rings with residue field k such that the kernel I satisfies $I^2 = 0$.

Given a diagram of solid arrows as below, we will show that we can find a lifting, marked as a dotted arrow.

$$\begin{array}{ccc} \mathrm{Spec}(A) & \longrightarrow & \widetilde{\mathcal{W}}_n^{\mathrm{lc}} \\ \downarrow & \nearrow \text{dotted} & \downarrow \\ \mathrm{Spec}(A') & \longrightarrow & \mathcal{M}_{c(\epsilon), v(\epsilon)}^{\mathrm{KSBA}} \end{array}$$

In other words, we need to show that for any family

$$(\mathcal{X}', c(\epsilon)S') \longrightarrow \mathrm{Spec} A'$$

in $\mathcal{M}_{c(\epsilon), v(\epsilon)}^{\mathrm{KSBA}}$ whose restriction to $\mathrm{Spec}(A)$ is a Weierstrass fibration, there is a morphism $\mathcal{X}' \rightarrow \mathbb{P}_{A'}^1$ such that $(\mathcal{X}' \rightarrow \mathbb{P}_{A'}^1, S')$ is a Weierstrass fibration. Namely, the deformation problem is the following:

$$\begin{array}{ccc} (\mathcal{X}, S) & \longrightarrow & (\mathcal{X}', S') \\ \downarrow & & \downarrow \text{dotted} \\ \mathbb{P}_A^1 & \dashrightarrow & \mathbb{P}_{A'}^1 \\ \downarrow & & \downarrow \\ \mathrm{Spec}(A) & \longrightarrow & \mathrm{Spec}(A') \end{array}$$

From [BHPS13, Prop. 3.10], it suffices to prove that $\mathrm{Hom}(\Omega_{\mathbb{P}^1}^1, R^1 f_* \mathcal{O}_X) = 0$. We have $R^1 f_* \mathcal{O}_X = \mathcal{O}_{\mathbb{P}^1}(-n)$ by Lemma 4.9, so

$$\mathrm{Hom}(\Omega_{\mathbb{P}^1}^1, R^1 f_* \mathcal{O}_X) \simeq H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-n+2)) = 0.$$

□

Lemma 4.9. *Let $(f : X \rightarrow \mathbb{P}^1, S)$ be an lc Weierstrass elliptic fibration with $S^2 = -n$. Then*

$$R^1 f_* \mathcal{O}_X = \mathcal{O}_{\mathbb{P}^1}(-n).$$

Proof. This follows because $h^1(\mathbb{P}^1, R^1 f_* (\mathcal{O}_X)) = n - 1$, as seen from the Leray spectral sequence for f and the computations $h^2(X, \mathcal{O}_X) = h^0(X, \omega_X) = n - 1$, and $h^0(\mathbb{P}^1, R^2 f_* \mathcal{O}_X) = h^2(\mathbb{P}^1, f_* \mathcal{O}_X) = 0$. □

Notation 4.10. We denote by $\mathcal{E}_n^{\mathrm{lc}}$ (resp. \mathcal{E}_n) the image of $\mathcal{W}_n^{\mathrm{lc}}$ (resp. $\mathcal{W}_n^{\mathrm{min}}$) under $\Phi_{n, \epsilon}$. We denote by $\Phi_n : \mathcal{W}_n^{\mathrm{lc}} \rightarrow \mathcal{E}_n^{\mathrm{lc}}$ the corresponding isomorphism.

Corollary 4.11. *The moduli space $\mathcal{E}_n^{\mathrm{lc}}$ is smooth and irreducible.* □

Proof. This follows from Theorem 4.2 and Proposition 4.8. □

Remark 4.12. It follows from [ABIP23, MZ23] that the compactification $\mathcal{E}_n \subseteq \overline{\mathcal{E}}_{n, \epsilon}$ of the image does not depend on ϵ for any ϵ close enough to either 0 or $\frac{n-2}{n}$. More generally, there are finitely many critical values

$$0 < t_1 < \cdots < t_m < \frac{n-2}{n}$$

such that $\overline{\mathcal{E}}_{n, t}$ is independent of t for $t_i < t < t_{i+1}$.

4.3. Locus of pseudo-elliptic surfaces in the KSB moduli. Consider the moduli stack \mathcal{E}_n constructed in the previous subsection. The closed points of \mathcal{E}_n parametrize KSBA-stable pairs

$$\left(X, \left(\frac{n-2}{n} - \epsilon \right) S \right)$$

such that there is a map $X \rightarrow \mathbb{P}^1$ which makes $(X \rightarrow \mathbb{P}^1, S)$ a Weierstrass fibration. Moreover, we proved that these pairs above do not depend on ϵ , as long $0 < \epsilon < \frac{n-2}{n}$. Therefore, it is natural to ask what happens if $\epsilon = 0$. In this case, there is a morphism

$$\Psi_n : \mathcal{E}_n \longrightarrow \mathcal{M}_v^{\mathrm{KSB}}$$

for $v = \frac{(n-2)^2}{n}$ which, on the level of points, sends a pair $(X, (\frac{n-2}{n} - \epsilon)S)$ to the log canonical model of $(X, \frac{n-2}{n}S)$, which is the surface obtained by contracting the negative section S (cf. Observation 3.14). The construction of such a morphism follows from some of the results [ABIP23, MZ23], which we recall here for the convenience of the reader.

Proposition 4.13. *Let $n \geq 3$ and $v = \frac{(n-2)^2}{n}$. Then there is a natural morphism $\Psi_n: \mathcal{E}_n^{\text{lc}} \rightarrow \mathcal{M}_v^{\text{KSB}}$.*

Proof. Let $(X, \frac{n-2}{n}S)$ be a pair parameterized by a closed point of $\mathcal{E}_n^{\text{lc}}$. This is an lc pair whose lc centers are isolated points, namely the cusps of any strictly lc fibers. Since $K_X + \frac{n-2}{n}S$ is big and nef, then by Kawamata-Viehweg vanishing for lc pairs [Fuj14, Thm. 1.10] we have that $H^i(X, m(K_X + \frac{n-2}{n}S)) = 0$ for any $m > 0$ such that $m(K_X + \frac{n-2}{n}S)$ is an integral divisor and for any $i > 0$. Then cohomology and base change applied to the universal family $\pi: (\mathcal{X}, \mathcal{S}) \rightarrow \mathcal{E}_n^{\text{lc}}$ implies that for any positive integer m such that $m(K_X + \frac{n-2}{n}S)$ is an integral divisor, the formation of

$$\mathcal{Y} = \text{Proj}_{\mathcal{E}_n^{\text{lc}}} \left(\bigoplus_{d \geq 0} \pi_* \mathcal{O}_{\mathcal{X}} \left(dm(K_{\mathcal{X}/\mathcal{E}_n^{\text{lc}}} + \frac{n-2}{n}\mathcal{S}) \right) \right)$$

commutes with base change, and the push-forwards $\pi_* \mathcal{O}_{\mathcal{X}}(dm(K_{\mathcal{X}/\mathcal{E}_n^{\text{lc}}} + \frac{n-2}{n}\mathcal{S}))$ are vector bundles. It gives rise to a projective morphism $\mathcal{Y} \rightarrow \mathcal{E}_n^{\text{lc}}$ which fiberwise is the canonical model of $(X, \frac{n-2}{n}S)$, and which is flat since we are taking Proj of a $\mathcal{E}_n^{\text{lc}}$ -flat algebra. Moreover, since the base is smooth (Corollary 4.11) and the volume of every fiber is $v = \frac{(n-2)^2}{n}$ (Observation 3.14), then the family $\mathcal{Y} \rightarrow \mathcal{E}_n^{\text{lc}}$ is a KSB stable family by Theorem 2.20. Therefore, it induces a morphism $\mathcal{E}_n^{\text{lc}} \rightarrow \mathcal{M}_v^{\text{KSB}}$ by the universality of $\mathcal{M}_v^{\text{KSB}}$. \square

Definition 4.14. We denote by \mathcal{P}_n the image of Ψ_n , and by $\overline{\mathcal{P}}_n$ its scheme-theoretic closure.

5. LOCAL STUDY OF THE CONTRACTION MORPHISM Ψ_n

In this section, we will prove that $\Psi_n: \mathcal{E}_n^{\text{lc}} \rightarrow \mathcal{M}_v^{\text{KSB}}$ is an open immersion when $n = 3$ or $n > 4$. To this end, we need a few observations on deformations of local models of the covering stack at the contracted point. We record the necessary lemmas first.

Lemma 5.1. *Let n be an even positive integer. Consider the action of $G = \mu_n/\mu_2$ on $Z = \text{Spec}(k[x, y, z]/(xy - z^2))$ defined by $\xi \cdot x = \xi^2 x$, $\xi \cdot y = \xi^2 y$ and $\xi \cdot z = \xi^2 z$.*

(1) *If $n > 4$ then*

$$\text{Ext}_{\mu_n/\mu_2}^1(\Omega_Z^1, \mathcal{O}_Z) = 0$$

and the only infinitesimal deformation of $[Z/G]$ is the trivial one.

(2) *If $n = 4$, then*

$$\text{Ext}_{\mu_n/\mu_2}^1(\Omega_Z^1, \mathcal{O}_Z) = k$$

and $[Z/G]$ admits a unique formal smoothing direction.

Proof. Let $R := k[x, y, z]/(xy - z^2)$ and consider the G -equivariant presentation of Ω_Z^1 given by

$$R^{\oplus 1} = Re_f \xrightarrow{1 \rightarrow (y, x, -2z)} R^{\oplus 3} = Rdx \oplus Rdy \oplus Rdz \longrightarrow \Omega_Z^1 \longrightarrow 0.$$

A generator $\xi \in \mu_n$ acts with weight 2 on dx , dy and dz and with weight 4 on e_f . To compute $\text{Ext}_{\mu_n/\mu_2}^1(\Omega_Z^1, \mathcal{O}_Z)$, we dualize the sequence above

$$\psi: R(dx)^\vee \oplus R(dy)^\vee \oplus R(dz)^\vee \longrightarrow Re_f^\vee,$$

and the cokernel of ψ is k with the action with weight $n - 4$. So there are no invariants if $n \neq 4$. If $n = 4$, then the invariants are one-dimensional and equal to the group $\text{Ext}^1(\Omega_Z^1, \mathcal{O}_Z)$ whose nonzero element corresponds to the unique smoothing direction of the A_1 singularity of Z . \square

Lemma 5.2. *Let R be an Artinian local ring over k , and consider the action of μ_n on $\mathbb{A}_R^2 = \text{Spec}(R[x, y])$ defined by $\xi \cdot x = \xi x$ and $\xi \cdot y = \xi y$. Let $f: \mathcal{I} \rightarrow \mathbb{A}_R^2$ be the pullback of the inertia stack of $[\mathbb{A}_R^2/\mu_n]$ along the smooth cover $\mathbb{A}_R^2 \rightarrow [\mathbb{A}_R^2/\mu_n]$. Then the scheme-theoretic support of the cokernel of $\mathcal{O}_{\mathbb{A}_R^2} \rightarrow f_* \mathcal{O}_{\mathcal{I}}$ is defined by the ideal (x, y) .*

Proof. The group scheme $\mathcal{I} \rightarrow \mathbb{A}_R^2$ fits into the following fiber product:

$$\begin{array}{ccc} \mathcal{I} & \longrightarrow & \mathbb{A}_R^2 \times \mu_n \\ f \downarrow & & \downarrow \\ \mathbb{A}_R^2 & \xrightarrow{\text{diag}} & \mathbb{A}_R^2 \times \mathbb{A}_R^2. \end{array}$$

If we write $\mu_n = \text{Spec}(k[t]/(t^n - 1))$, then $\mathcal{I} = \text{Spec}(R[x, y, t]/(t^n - 1, tx - x, ty - y))$, and the morphism f induces the inclusion $R[x, y] \rightarrow R[x, y, t]/(t^n - 1, tx - x, ty - y)$ on global sections. In particular, we have

$$f_* \mathcal{O}_{\mathcal{I}} / \mathcal{O}_{\mathbb{A}_R^2} = \bigoplus_{i=0}^{n-1} R \cdot t^i \cong (R[x, y]/(x, y))^{\oplus n}$$

as an $R[x, y]$ -module, which has scheme-theoretic support defined by the ideal (x, y) as desired. \square

Lemma 5.3. *Let R be an Artinian local ring, and consider the action of μ_n on $Z := \text{Spec}(R[x, y, z]/(xy - z^2))$ defined by $\xi \cdot x = \xi x$, $\xi \cdot y = \xi y$ and $\xi \cdot z = \xi z$. Let $f: \mathcal{I} \rightarrow Z$ be the pullback of the inertia stack of $[Z/\mu_n]$ along $Z \rightarrow [Z/\mu_n]$. Then the support of the cokernel of $\mathcal{O}_Z \rightarrow f_* \mathcal{O}_{\mathcal{I}}$ is the vanishing of the ideal (x, y, z) .*

Proof. The argument is the same as in Lemma 5.2: in this case we have that

$$\mathcal{I} = \text{Spec}(R[x, y, z, t]/(xy - z^2, t^n - 1, tx - x, ty - y, tz - z))$$

and f is induced by the following inclusion on global sections

$$R[x, y, z]/(xy - z^2) \rightarrow R[x, y, z, t]/(xy - z^2, t^n - 1, tx - x, ty - y, tz - z).$$

We can conclude similarly as before. \square

Theorem 5.4. *Suppose that $n > 2$ and $n \neq 4$ is an integer. Then the morphism*

$$\Psi_n : \mathcal{E}_n^{\text{lc}} \longrightarrow \mathcal{M}_v^{\text{KSB}}$$

defined in Proposition 4.13 is an open immersion.

Proof. In this proof, we will denote by Y the pseudo-elliptic surface obtained by contracting the section S of an elliptic surface X with $S^2 = -n$, and by p be the point to which S is contracted. From [KM98, Remark 4.9 (2)] the singularity is determined from the dual graph associated to its minimal resolution, and by the local analysis of Lemma 3.13 we have that Y has a $\frac{1}{n}(1, 1)$ -singularity at p .

Ψ_n is injective on isomorphism classes of geometric points. It suffices to check that there is an inverse on sets of geometric points. As all the singularities of Y away from p are either Du Val or strictly log-canonical (see Remark 3.5), we can identify X with the minimal resolution of Y around its unique klt but not canonical singularity, and S with its exceptional divisor.

Ψ_n is injective on automorphism groups. Let $(f: X \rightarrow \mathbb{P}^1, S)$ be a Weierstrass fibration associated to a geometric point of $\mathcal{E}_n^{\text{lc}}$ and let Y be its image via Ψ_n . It suffices to observe that any automorphism on X which induces the identity on Y agrees with the identity on the dense open subset $X \setminus S$, so it must be the identity.

Ψ_n is surjective on automorphism groups. This is because any automorphism of Y must fix p as it is the unique strictly klt singularity. Thus, any such automorphism lifts to the blowup of Y around p , which is exactly (X, S) .

Ψ_n is smooth: we apply the infinitesimal lifting property of smoothness (cf. [Sta24, Tag 0DP0], [Sta24, Tag 02HT]). We need to show that if $A' \rightarrow A$ is a quotient of an Artinian local rings with residue field k and with square-zero ideal, and if we have a diagram of solid arrows as below, then one can find the dotted arrow:

$$\begin{array}{ccc} \text{Spec}(A) & \longrightarrow & \mathcal{E}_n^{\text{lc}} \\ \downarrow & \searrow \text{dotted} & \downarrow \\ \text{Spec}(A') & \longrightarrow & \mathcal{M}_v^{\text{KSB}}. \end{array}$$

The morphism $\mathrm{Spec}(A') \rightarrow \mathcal{M}_v^{\mathrm{KSB}}$ induces a KSB-stable family $Y' \rightarrow \mathrm{Spec}(A')$, and let $\mathcal{Y}' \rightarrow \mathrm{Spec}(A')$ its covering stack (cf. Definition 2.21). As mentioned at the beginning of the proof, the singularity at p is formally locally isomorphic to

$$(0 \in \mathrm{Spec} k[[x, y]]/\mu_n)$$

with the action $\xi \cdot x = \xi x$ and $\xi \cdot y = \xi y$.

The canonical covering stack $\mathcal{Y}_k \rightarrow Y_k$, on a neighborhood of p , is formally locally isomorphic to:

- (1) $[\mathrm{Spec}(k[[x, y]]/\mu_n)]$ for n odd with the action $\xi \cdot x = \xi x$ and $\xi \cdot y = \xi y$; and
- (2) $[\mathrm{Spec}(k[[x, y, z]]/xy - z^2)/(\mu_n/\mu_2)]$ for n even, with the action $\xi \cdot x = \xi^2 x$, $\xi \cdot y = \xi^2 y$ and $\xi \cdot z = \xi^4 z$.

Indeed, formally locally around p , the covering stack \mathcal{Y}_k is the relative coarse moduli space of the map $[\mathrm{Spec} k[[x, y]]/\mu_n \rightarrow \mathbf{B}\mathbb{G}_m$ given by the line bundle with section $dx \wedge dy$. More explicitly, it is the stacky quotient of $\mathrm{Spec} k[[x, y]]$ by the kernel of the representation of μ_n on $dx \wedge dy$. As ξ acts on $dx \wedge dy$ as

$$\xi \cdot (dx \wedge dy) = \xi^2 dx \wedge dy,$$

the kernel is trivial if n is odd and μ_2 if it is even.

Since $[\mathrm{Spec}(k[[x, y]]/\mu_n)$ is smooth, by Lemma 5.1(1), the small deformations of the analytic local singularity of \mathcal{Y}_k at the preimage of p are trivial for $n > 2$ and $n \neq 4$. In particular, for n odd, there is roof diagram of pointed stacks as follows, with all the arrows étale and inducing an isomorphism on automorphisms groups:

$$([\mathrm{Spec}(A'[x, y]]/\mu_n), 0) \longleftarrow (U, u) \longrightarrow (\mathcal{Y}', y).$$

Similarly, for n even, we have a diagram as follows

$$([\mathrm{Spec}(A'[x, y, z]/xy - z^2)/(\mu_n/\mu_2)], 0) \longleftarrow (W, w) \longrightarrow (\mathcal{Y}', y).$$

From Lemma 5.2 and Lemma 5.3, the support of the inertia is the closed substack which on U is the pull-back of the vanishing of (x, y) and on W is the vanishing of (x, y, z) . Hence, we can perform the blow-up along the closed substack given by the support of the inertia stack, which étale locally corresponds to performing the blow up of (x, y) in $[\mathrm{Spec}(A'[x, y]]/\mu_n$ and (x, y, z) in $[\mathrm{Spec}(A'[x, y, z]/xy - z^2)/(\mu_n/\mu_2)]$. In particular, from the analogous computation on the local charts, this blow-up, denoted by $\mathcal{X}' \rightarrow \mathcal{Y}'$, is flat and commutes with base change. Taking the coarse moduli space commutes with base change, so if $\mathcal{X}' \rightarrow X'$ is the coarse moduli space of \mathcal{X}' , then $\mathcal{X}_A := \mathcal{X}'_A \rightarrow X_A := X'_A$ is the coarse moduli space of \mathcal{X}_A . Then it follows from [Kol09b, Section 2.4 page 86] that the surface X_A is a minimal resolution of Y_A around the strictly klt singularity: we have that X_A is the elliptic surface associated to $\mathrm{Spec}(A) \rightarrow \mathcal{E}_n^{\mathrm{lc}}$. The desired family giving the morphism $\mathrm{Spec}(A') \rightarrow \mathcal{E}_n^{\mathrm{lc}}$ is the pair $(X', S') \rightarrow \mathrm{Spec}(A')$ where S' is the coarse moduli space of the exceptional divisor of $\mathcal{X}' \rightarrow \mathcal{Y}'$.

In particular, $\mathcal{M}_v^{\mathrm{KSB}}$ is smooth along the image of Ψ_n , which is open since Ψ_n is smooth. Therefore, it follows that Ψ_n is an étale monomorphism, and hence an open immersion. \square

Corollary 5.5. *The scheme-theoretic image $\overline{\mathcal{P}}_n$ is an irreducible component of $\mathcal{M}_v^{\mathrm{KSB}}$ for $n = 3$ and $n > 4$.*

Remark 5.6. In fact, the conclusion of the previous corollary holds even for $n = 4$. Indeed, the proof of Theorem 5.4 shows that any deformation of Y which is locally trivial around p is in the image of Ψ_n for any n . Thus, if $\overline{\mathcal{P}}_4$ is not an irreducible component of $\mathcal{M}_v^{\mathrm{KSB}}$, then there is a large deformation of Y which is not locally trivial around p . By 5.1(2), this deformation must be a \mathbb{Q} -Gorenstein smoothing of Y , but it is well known that Y cannot admit a projective \mathbb{Q} -Gorenstein smoothing since such a smoothing would violate the Noether inequality: indeed, one has

$$K^2 = 1, \text{ and } p_g = h^0(X, K_X) = 3.$$

On the other hand, it is an interesting question whether this irreducible component of $\mathcal{M}_v^{\mathrm{KSB}}$ has some non-trivial generic non-reduced structure for $n = 4$.

6. THE CASE WHEN $n = 3$

In this section, we present a more explicit proof of some of our results in the case when $n = 3$. Throughout most of this section, we will adopt the following assumptions.

Context 6.1. Let $(g: X \rightarrow \mathbb{P}^1, S)$ be a Weierstrass fibration with X smooth, with $S^2 = -3$ and with 36 singular nodal fibers. Let $\pi: X \rightarrow Y$ be the contraction of S , and p_i be the nodal points on the singular fibers of g .

Lemma 6.2. *The locus in \mathcal{W}_3^{\min} where the conditions of Context 6.1 are satisfied is open.*

Proof. A Weierstrass fibration $(g: X \rightarrow \mathbb{P}^1, S)$ of height 3 has 36 nodal singular fibers if and only if all the fibers are of Kodaira type I_1 . The condition of g having only I_1 fibers is equivalent to g having only nodal fibers and X being smooth. Thus, the locus satisfying the required condition is exactly the intersection $\mathcal{W}_3^{\text{nod}} \cap \mathcal{W}_3^{\text{reg}} \subset \mathcal{W}_3^{\min}$ which is open by Theorem 4.2(2). \square

The following two exact sequences will be useful:

$$(5) \quad 0 \longrightarrow \Omega_{X/\mathbb{P}^1}^1 \longrightarrow \omega_g \longrightarrow \bigoplus_{i=1}^{36} k_{p_i} \longrightarrow 0,$$

$$(6) \quad 0 \longrightarrow g^* \omega_{\mathbb{P}^1} \longrightarrow \Omega_X^1 \longrightarrow \Omega_{X/\mathbb{P}^1}^1 \longrightarrow 0.$$

We begin with the following preliminary computations.

Lemma 6.3. *In the situation of Context 6.1, we have the following:*

$$(7) \quad \text{Ext}^1(g^* \omega_{\mathbb{P}^1}, \mathcal{O}_X) = H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2)) \oplus H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) = 0$$

$$(8) \quad \text{Ext}^1(\omega_g, \mathcal{O}_X) = H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-3)) \oplus H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-6)) \simeq k^{\oplus 2}$$

$$(9) \quad H^1(X, g^* \omega_{\mathbb{P}^1} \otimes g^* \mathcal{O}_{\mathbb{P}^1}(1)) = H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) \oplus H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-4)) = 0$$

Proof. These follow from the Leray spectral sequence for $g: X \rightarrow \mathbb{P}^1$, the fact that $\omega_g = g^* \mathcal{O}_{\mathbb{P}^1}(3)$, the fact that $R^1 g_* \mathcal{O}_X = \mathcal{O}_{\mathbb{P}^1}(-3)$, and the projection formula. We will compute the first one to illustrate this. First note that

$$\text{Ext}^1(g^* \omega_{\mathbb{P}^1}, \mathcal{O}_X) = H^1(X, g^* \omega_{\mathbb{P}^1}^\vee) = H^1(X, g^* \mathcal{O}_{\mathbb{P}^1}(2)).$$

Then by the projection formula, $g_* g^* \mathcal{O}_{\mathbb{P}^1}(2) = \mathcal{O}_{\mathbb{P}^1}(2)$ and $R^1 g_* g^* \mathcal{O}_{\mathbb{P}^1}(2) = \mathcal{O}_{\mathbb{P}^1}(-1)$ and the Leray spectral sequence yields the following exact sequence and the claim.

$$0 \rightarrow H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2)) \rightarrow H^1(X, g^* \mathcal{O}_{\mathbb{P}^1}(2)) \rightarrow H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) \rightarrow 0$$

\square

Lemma 6.4. *In the situation of Context 6.1, we have*

$$h^2(X, T_X(-S)) = 0 \quad \text{and} \quad h^1(X, T_X) = 30.$$

Proof.

Proof of $h^2(X, T_X(-S)) = 0$. From Serre duality, it suffices to show that $h^0(X, \Omega_X^1 \otimes \omega_X(S)) = 0$. Twisting (6) by $\omega_X(S)$ and taking the induced long exact sequence, we get the following.

$$0 \rightarrow H^0(X, g^* \omega_{\mathbb{P}^1} \otimes \omega_X(S)) \rightarrow H^0(X, \Omega_X^1 \otimes \omega_X(S)) \rightarrow H^0(X, \Omega_{X/\mathbb{P}^1}^1 \otimes \omega_X(S)) \rightarrow H^1(X, g^* \omega_{\mathbb{P}^1} \otimes \omega_X(S))$$

Note that $g_* \mathcal{O}_X(S) = g_* \mathcal{O}_X = \mathcal{O}_{\mathbb{P}^1}$ [CDLI24, Lem. 2.5], $R^1 g_* \mathcal{O}_X(S) = 0$ by cohomology and base change, and $\omega_X = g^* \mathcal{O}_{\mathbb{P}^1}(1)$ by the canonical bundle formula (Theorem 3.3). By the projection formula, $g_*(g^* \omega_{\mathbb{P}^1} \otimes \omega_X(S)) = \mathcal{O}_{\mathbb{P}^1}(-1)$ so

$$H^0(X, g^* \omega_{\mathbb{P}^1} \otimes \omega_X(S)) = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) = 0.$$

Similarly, from the Leray spectral sequence, we have $H^1(X, g^* \omega_{\mathbb{P}^1} \otimes \omega_X(S)) = H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-1)) = 0$. Therefore

$$H^0(X, \Omega_X^1 \otimes \omega_X(S)) = H^0(X, \Omega_{X/\mathbb{P}^1}^1 \otimes \omega_X(S)).$$

So it suffices to show that $H^0(X, \Omega_{X/\mathbb{P}^1}^1 \otimes \omega_X(S)) = 0$. Twisting (5) by $\omega_X(S)$ and taking the associated long exact sequence, leads to

$$0 \rightarrow H^0(X, \Omega_{X/\mathbb{P}^1}^1 \otimes \omega_X(S)) \rightarrow H^0(X, \omega_g \otimes \omega_X(S)) \xrightarrow{\alpha} \bigoplus_{i=1}^{36} k_{p_i}.$$

But

$$H^0(X, \omega_g \otimes \omega_X(S)) = H^0(X, g^* \mathcal{O}_{\mathbb{P}^1}(3) \otimes g^* \mathcal{O}_{\mathbb{P}^1}(1)(S)) = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(4) \otimes g_* \mathcal{O}_X(S)) = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(4)).$$

We can identify the map α as the evaluation of a quartic polynomial on \mathbb{P}^1 on the 36 points on \mathbb{P}^1 whose fibers via g are singular. Then α is injective, as a quartic polynomial on \mathbb{P}^1 that vanishes at 36 points has to be zero. Hence, we have the desired vanishing.

Proof of $h^1(X, T_X) = 30$. Observe first that $h^2(X, T_X) = 0$. Indeed, we have an exact sequence

$$0 \rightarrow T_X(-S) \rightarrow T_X \rightarrow (T_X)|_S \rightarrow 0.$$

As S has dimension 1, we have that $h^2(S, (T_X)|_S) = 0$. The desired vanishing follows from the previous point, and the long exact sequence in cohomology. Moreover, we have

- $\text{Ext}^1(\bigoplus_{i=1}^{36} k_{p_i}, \mathcal{O}_X) = 0$,
- $\text{Ext}^2(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) = 0$, and
- $H^0(X, \Omega_X^1) = 0$.

Indeed, the first bullet point follows from the local-to-global spectral sequence for Ext , and since the points p_i are smooth points of the surface X . The second one follows by applying $\text{Hom}(\bullet, \mathcal{O}_X)$ to the short exact sequence (6), Lemma 6.3(7), and the fact that $\text{Ext}^2(\Omega_X^1, \mathcal{O}_X) = H^2(X, T_X) = 0$ we just proved. The third bullet point follows since there is an injection $H^0(X, \Omega_X^1) \rightarrow H^0(X, \Omega_X^1 \otimes \omega_X(S))$ and above we prove that the latter is 0.

Then, if we apply $\text{Hom}(\bullet, \mathcal{O}_X)$ to the short exact sequence (5), using the previous two vanishings we get

$$0 \longrightarrow \text{Ext}^1(\omega_g, \mathcal{O}_X) \longrightarrow \text{Ext}^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) \longrightarrow \text{Ext}^2(\bigoplus_{i=1}^{36} k_{p_i}, \mathcal{O}_X) \longrightarrow \text{Ext}^2(\omega_g, \mathcal{O}_X) \longrightarrow 0.$$

From Lemma 6.3(8), $\text{ext}^1(\omega_g, \mathcal{O}_X) = 2$ and $\text{ext}^2(\omega_g, \mathcal{O}_X) = 5$. By Serre duality, $\text{ext}^2(\bigoplus k_{p_i}, \mathcal{O}_X) = h^0(X, \bigoplus k_{p_i}) = 36$, therefore

$$\text{ext}^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) = 33.$$

Applying $\text{Hom}(\bullet, \mathcal{O}_X)$ to the sequence (6), using $H^0(X, \Omega_X) = 0$ and Lemma 6.3(7), we obtain

$$0 \longrightarrow \text{Hom}(g^* \omega_{\mathbb{P}^1}, \mathcal{O}_X) \longrightarrow \text{Ext}^1(\Omega_{X/\mathbb{P}^1}^1, \mathcal{O}_X) \longrightarrow \text{Ext}^1(\Omega_X^1, \mathcal{O}_X) \longrightarrow 0.$$

As we have

$$\text{Hom}(g^* \omega_{\mathbb{P}^1}, \mathcal{O}_X) = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2) \otimes g_* \mathcal{O}_X) = H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(2)) \simeq k^{\oplus 3},$$

then $h^1(X, T_X) = 33 - 3 = 30$. □

Corollary 6.5. *In the situation of Context 6.1, we have*

$$h^1(Y, T_Y) = 28, \quad \text{and} \quad h^2(Y, T_Y) = 0,$$

where we denote $T_Y := \pi_* T_X$.

Proof. First, we prove that $R^1 \pi_* T_X \cong k_p^{\oplus 2}$, where $p \in Y$ is the point to which S is contracted. As π is an isomorphism away from p , the sheaf $R^1 \pi_* T_X$ is a skyscraper sheaf supported at p . We now compute its length. Let S_m be the m -th thickened neighborhood of S , and let \mathcal{I} be the ideal sheaf of $S = S_1$ in X . Notice that we have

$$\mathcal{I}/\mathcal{I}^2 \simeq \mathcal{N}_{S/X}^* \simeq \mathcal{O}_{\mathbb{P}^1}(3), \quad \text{and} \quad \mathcal{I}^m/\mathcal{I}^{m+1} \simeq (\mathcal{I}/\mathcal{I}^2)^m \simeq \mathcal{O}_{\mathbb{P}^1}(3m).$$

Also, notice that $T_S \simeq \mathcal{O}_{\mathbb{P}^1}(2)$ and we have the exact sequence

$$0 \longrightarrow T_S \longrightarrow T_X|_S \longrightarrow N_{S/X} \longrightarrow 0,$$

then $T_X|_S \simeq \mathcal{O}_{\mathbb{P}^1}(2) \oplus \mathcal{O}_{\mathbb{P}^1}(-3)$ since the extension has to be trivial (as $\text{Ext}^1(\mathcal{O}_{\mathbb{P}^1}(-3), \mathcal{O}_{\mathbb{P}^1}(2)) = 0$). Taking cohomology of the exact sequence

$$0 \longrightarrow (\mathcal{I}^m/\mathcal{I}^{m+1}) \otimes T_X \longrightarrow T_X|_{S_{m+1}} \longrightarrow T_X|_{S_m} \longrightarrow 0,$$

we see that $H^1(T_X|_{S_{m+1}})$ is canonically isomorphic to $H^1(T_X|_{S_m})$, which is in turn isomorphic to $H^1(T_X|_S) = k^{\oplus 2}$. It follows from the theorem on formal functions that

$$(\widehat{R^1\pi_*T_X})_p = \varprojlim_m H^1(S_m, T_X|_{S_m}) \simeq k^{\oplus 2}.$$

Taking the five-term exact sequence associated to the Leray spectral sequence

$$E_2^{p,q} := H^q(Y, R^p\pi_*T_X) \Rightarrow H^{p+q}(X, T_X),$$

we get an exact sequence

$$0 \rightarrow H^1(Y, T_Y) \rightarrow H^1(X, T_X) \rightarrow H^0(Y, R^1\pi_*T_X) \rightarrow H^2(Y, T_Y) \rightarrow H^2(X, T_X).$$

Since $H^2(X, T_X(-S)) = 0$ by Lemma 6.4, the map $H^1(X, T_X) \rightarrow H^1(X, T_X|_S)$ is surjective. As a consequence, the morphism

$$H^1(X, T_X) \longrightarrow H^0(Y, R^1\pi_*T_X)$$

is surjective. The desired statement now follows from Lemma 6.4. \square

Proposition 6.6. *Set $\mathcal{U} := X \setminus S$ and denote by $i : \mathcal{U} \rightarrow X$ the corresponding inclusion. Then we have $\pi_*T_X = \pi_*i_*T_{\mathcal{U}}$.*

Proof. Since we have $\omega_X = g^*\mathcal{O}_{\mathbb{P}^1}(1)$, then there is an exact sequence

$$0 \longrightarrow \omega_X^{-1} \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_F \longrightarrow 0$$

where F is a (general) fiber of g . Twisting the previous sequence by Ω_X^1 and applying $T_X \simeq \Omega_X^1 \otimes \omega_X^{-1}$, we get that

$$0 \longrightarrow T_X \longrightarrow \Omega_X^1 \longrightarrow \mathcal{O}_F \otimes \Omega_X^1 \longrightarrow 0.$$

Consider now the following diagram, where V is a Zariski open neighborhood of S , and the vertical arrows are restrictions to $U := V \setminus S$

$$\begin{array}{ccccc} 0 & \longrightarrow & \Gamma(V, T_X) & \longrightarrow & \Gamma(V, \Omega_X^1) & \longrightarrow & \Gamma(V \cap F, \Omega_X^1|_F) \\ & & \downarrow & & \alpha \downarrow & & \beta \downarrow \\ 0 & \longrightarrow & \Gamma(U, T_U) & \longrightarrow & \Gamma(U, \Omega_U^1) & \longrightarrow & \Gamma(U \cap F, \Omega_X^1|_{F \cap U}) \end{array}$$

The map β is injective, as Ω_X^1 is a vector bundle on an integral scheme, so the restriction to the generic point is an injective morphism. The morphism α is an isomorphism by [GKKP11, Obs. 1.3, Thm. 1.4], so from diagram chasing the first map is an isomorphism. In particular, $\pi_*T_X = \pi_*i_*T_{\mathcal{U}}$ as desired. \square

Proposition 6.7. *The moduli stack \mathcal{P}_3 is smooth of dimension 28 at the points Y arising from Context 6.1 as above.*

Proof. From Theorem 2.22, if we denote by \mathcal{Y} the covering stack of Y , it suffices to check that

- (1) \mathcal{Y} is smooth,
- (2) $h^2(\mathcal{Y}, T_{\mathcal{Y}}) = 0$, and
- (3) $h^1(\mathcal{Y}, T_{\mathcal{Y}}) = 28$.

Indeed, (1) follows from the proof of Theorem 5.4. The second one holds true since, if we denote by $q : \mathcal{Y} \rightarrow Y$ the coarse space map, then

$$H^2(\mathcal{Y}, T_{\mathcal{Y}}) = H^2(Y, q_*T_{\mathcal{Y}}), \quad \text{and} \quad H^1(\mathcal{Y}, T_{\mathcal{Y}}) = H^1(Y, q_*T_{\mathcal{Y}})$$

as q_* is exact (cf. [AV02, Lem. 2.3.4]). But as \mathcal{Y} is smooth, if we denote by $j : \mathcal{U} \rightarrow \mathcal{Y}$ the inclusion of the schematic locus of \mathcal{Y} (namely, \mathcal{Y} without a single point), then

$$T_{\mathcal{Y}} = j_*T_{\mathcal{U}} \quad \text{and} \quad q_*T_{\mathcal{Y}} = q_*j_*T_{\mathcal{U}}.$$

From Proposition 6.6, $q_*T_Y = \pi_*T_X$, and the desired statements now follow from Corollary 6.5. \square

Theorem 6.8. *The morphism $\Psi_3 : \mathcal{E}_3^{\text{lc}} \rightarrow \mathcal{M}_{\frac{1}{3}}^{\text{KSB}}$ is an open embedding at the points (X, S) as in Context 6.1.*

Proof. We apply Zariski's main theorem, which requires that Ψ_3 is representable, injective, and birational.

To check that Ψ_3 is representable, we need to check that if σ is an automorphism of (X, S) which induces the identity on Y , then it is the identity. This is clear, as if σ is the identity on a dense open subset (namely, the complement of S), then it has to be the identity.

To check that Ψ_3 is injective: we can construct X as the minimal resolution of Y , and S is the exceptional divisor. Since the minimal resolution is unique, Ψ_3 is injective.

To check that Ψ_3 is birational: from Proposition 6.7 the dimension of \mathcal{P}_3 is 28. From the isomorphism between \mathcal{E}_3 and \mathcal{W}_3 , we have $\dim(\mathcal{E}_3) = \dim(\mathcal{W}_3)$, and the latter is 28 from the explicit description of \mathcal{W}_3 as a quotient stack given in [CDLI24, Section 2].

As both \mathcal{E}_3 and \mathcal{P}_3 are smooth (hence normal), from Zariski's main theorem the map Ψ_3 is an open embedding at the points (X, S) as above (which is an open locus in \mathcal{E}_3 from Lemma 6.2). \square

6.1. **Case $n > 3$.** It is natural to wonder if the (naive) purely cohomological methods of this section can be applied even when $n > 3$. Unfortunately, the answer is no.

For this subsection X will be a smooth Weierstrass fibration with $12n$ singular fibers, and $n \geq 4$. As before, we have the exact sequence

$$(10) \quad 0 \longrightarrow g^*\omega_{\mathbb{P}^1} \longrightarrow \Omega_X^1 \longrightarrow \Omega_{X/\mathbb{P}^1}^1 \longrightarrow 0,$$

where the points p_i are the nodes in the singular fibers of g .

Lemma 6.9. *The cohomology group $H^2(X, T_X)$ is non-zero when $n \geq 4$.*

Proof. Using Serre duality, it suffices to check that $H^0(X, \Omega_X^1 \otimes \omega_X) \neq 0$. Twisting (10) by ω_X and taking the induced long exact sequence on cohomology, one sees that $H^0(X, \Omega_X^1 \otimes \omega_X)$ contains a subgroup $H^0(X, \omega_X \otimes g^*\mathcal{O}_{\mathbb{P}^1}(-2))$. Using that $\omega_X \simeq g^*\mathcal{O}_{\mathbb{P}^1}(n-2)$ and $g_*\mathcal{O}_X \simeq \mathcal{O}_{\mathbb{P}^1}$, we conclude that:

$$H^0(X, \omega_X \otimes g^*\mathcal{O}_{\mathbb{P}^1}(-2)) \simeq H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(n-4)).$$

and hence this group is nonzero for $n \geq 4$. \square

Lemma 6.10. *The cohomology group $\text{Ext}^2(\Omega_X^1(\log S), \mathcal{O}_X)$ is non-zero when $n \geq 4$.*

Recall that the obstruction to a deformation of (X, S) lies in $\text{Ext}^2(\Omega_X^1(\log S), \mathcal{O}_X)$.

Proof. We utilize the long exact sequence obtained by applying $R\text{Hom}(-, \mathcal{O}_X)$ to the short exact sequence

$$0 \longrightarrow \Omega_X^1 \longrightarrow \Omega_X^1(\log S) \longrightarrow \mathcal{O}_S \longrightarrow 0,$$

the last terms of which are

$$\cdots \longrightarrow \text{Ext}^2(\mathcal{O}_S, \mathcal{O}_X) \longrightarrow \text{Ext}^2(\Omega_X^1(\log S), \mathcal{O}_X) \longrightarrow \text{Ext}^2(\Omega_X^1, \mathcal{O}_X) \longrightarrow 0.$$

The final term $\text{Ext}^2(\Omega_X^1, \mathcal{O}_X) \cong H^2(X, T_X)$ is non-zero when $n \geq 4$ by Lemma 6.9 and the result follows. \square

7. COMPACTIFICATIONS

In this section we discuss the natural compactification of \mathcal{E}_n (resp. \mathcal{P}_n) given by taking the closure in the proper moduli stack $\mathcal{M}_{c(\epsilon), v(\epsilon)}^{\text{KSBA}}$ (resp. $\mathcal{M}_v^{\text{KSB}}$). By the main theorem, these compactifications are irreducible components of the KSBA (resp. KSB) moduli spaces for $n \neq 4$.

We use two tools to understand these compactifications: twisted stable maps and wall-crossing as developed in [AV97, AV02, AB19, AB23] and [AB21, Inc20, ABIP23, MZ23] respectively.

The starting point is the observation that an elliptic surface $(X \rightarrow C, S)$ with at worst I_k fibers is equivalent to a morphism

$$g: C \rightarrow \overline{\mathcal{M}}_{1,1}$$

to the moduli stack of pointed elliptic curves. The I_k fibers lie over the preimages of $\infty \in \overline{\mathcal{M}}_{1,1}$ and k is the ramification of g at a given preimage. The composition to the coarse moduli space $j: C \rightarrow \overline{\mathcal{M}}_{1,1}$ is the j -invariant of the elliptic surface. The degree of the j -map satisfies $\deg(j) = 12n$, where n is the height. The space of maps admits a compactification by a proper Deligne-Mumford stack

$$\mathcal{K}_n := \overline{\mathcal{K}}_{0,0}(\overline{\mathcal{M}}_{1,1}, n)$$

parametrizing 0-pointed, genus 0 and degree n twisted stable maps (cf. [AV02]).

Definition 7.1. A 0-pointed twisted stable map of genus g and degree n is a commutative diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{j'} & \overline{\mathcal{M}}_{1,1} \\ \pi \downarrow & & \downarrow \\ C & \xrightarrow{j} & \overline{\mathcal{M}}_{1,1} \end{array}$$

such that

- (1) \mathcal{C} is a stacky curve with at worst nodal singularities, and $\pi: \mathcal{C} \rightarrow C$ is the coarse moduli space;
- (2) π is an isomorphism over the non-singular locus of C ;
- (3) \mathcal{C} is formally locally isomorphic around each node to

$$[(\mathrm{Spec} k[[x, y]]/(xy))/\mu_r] \quad \text{by} \quad (x, y) \mapsto (\xi \cdot x, \xi^{-1} \cdot y);$$

- (4) j' is a representable morphism which induces the map j ; and
- (5) j is a stable map of genus g and degree $12n$.

Since the bottom half of the diagram is determined by the top half, we often just write $(j': \mathcal{C} \rightarrow \overline{\mathcal{M}}_{1,1})$.

Theorem 7.2. *The stack \mathcal{K}_n is irreducible.*

Proof. By [AB23, Thm. 5.6], any genus 0 twisted stable map $g_0: \mathcal{C}_0 \rightarrow \overline{\mathcal{M}}_{1,1}$ can be deformed to a family of maps

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{g} & \overline{\mathcal{M}}_{1,1} \\ \downarrow & & \\ \mathrm{Spec} R & & \end{array}$$

over the spectrum of a DVR R with closed point 0 and generic point $\eta \in \mathrm{Spec} R$ such that \mathcal{C}_η is a smooth genus 0 curve. Thus, the locus of maps with smooth source is dense in \mathcal{K}_n . On the other hand, by definition, if \mathcal{C} is smooth, then $\mathcal{C} \rightarrow C$ is an isomorphism and so $\mathcal{C} \cong \mathbb{P}^1$. We conclude that the space of twisted stable maps to $\overline{\mathcal{M}}_{1,1}$ of degree n and genus 0 with *smooth* source curve is simply the space $\mathcal{W}_n^{\mathrm{nod}}$ of Weierstrass elliptic fibrations of height n with at worst nodal fibers, which is irreducible by Theorem 4.2(2). \square

Proposition 7.3 ([AB21]). *There exists a commutative diagram*

$$\begin{array}{ccc} & \mathcal{K}_n^\nu & \\ & \swarrow & \searrow \bar{\Phi}_n \\ \mathcal{K}_n & & \bar{\mathcal{E}}_{n,\epsilon} \xrightarrow{\bar{\Psi}_n} \bar{\mathcal{P}}_n \end{array}$$

where \mathcal{K}_n^ν is the normalization of \mathcal{K}_n , $0 < \epsilon \ll 1$ and $\bar{\Phi}_n$ and $\bar{\Psi}_n$ extend $\Phi_n|_{\mathcal{W}_n^{\mathrm{nod}}}$ and Ψ_n respectively.

7.1. From twisted stable maps limits to KSBA limits. There are two stability conditions that we will consider for elliptic surfaces. The first one comes from *twisted stable maps*, as in Definition 7.1, and the second one from KSB, as in Section 2.3.

Definition 7.4. An elliptic surface $(X, S) \rightarrow C$ is *twisted stable maps-stable* (abbrv. tsm-stable) if there is 0-pointed twisted stable map $(j': \mathcal{C} \rightarrow \overline{\mathcal{M}}_{1,1})$ of a certain genus such that:

- (1) C is the coarse moduli space of \mathcal{C} ;

- (2) if we denote by $(\mathcal{X}, \mathcal{S}) \rightarrow \mathcal{C}$ the pull-back of the universal curve and the universal section to \mathcal{C} , then (X, S) is the coarse moduli space of $(\mathcal{X}, \mathcal{S})$; and
- (3) the projection morphism $X \rightarrow C$ coincides with the induced morphism between coarse moduli spaces of $\mathcal{X} \rightarrow \mathcal{C}$.

Remark 7.5. One can check that the locus where \mathcal{C} is a stack (but not a scheme) can be determined from the data of $X \rightarrow C$: it is along the nodes $\mathfrak{n} \in C$ such that the fiber of $X \rightarrow C$ is non-reduced. Indeed, the fibers of $\mathcal{X} \rightarrow \mathcal{C}$ are parametrized by $\overline{\mathcal{M}}_{1,1}$, so if we take the reduced structure on the geometric fiber of $X \rightarrow C$ over \mathfrak{n} , the resulting curve is a quotient of a Deligne-Mumford stable 1-pointed genus 1 curve (E, p) , by a subgroup Γ of $\text{Aut}(E, p)$, and $\Gamma = \{1\}$ if and only if \mathcal{C} is a scheme at the node \mathfrak{n} ; see [AB19].

The morphism Φ_n is described as follows. For each point $p \in \mathcal{K}'_n$, consider a one parameter family $\text{Spec}(R) \rightarrow \mathcal{K}'_n$ over the spectrum of a DVR R , which maps the generic point η to the locus parametrizing minimal Weierstrass fibrations in \mathcal{K}_n , and the special point, which we denote by 0 , to p . One has the resulting family of tsm-stable elliptic surfaces:

$$(\mathcal{X}, \mathcal{S}) \longrightarrow \mathcal{C} \longrightarrow \text{Spec}(R).$$

Choose $0 < \epsilon \ll 1$, and consider the canonical model

$$(\mathcal{X}^c, \epsilon \mathcal{S}^c) \longrightarrow \text{Spec}(R) \quad \text{of} \quad (\mathcal{X}, \epsilon \mathcal{S}) \longrightarrow \text{Spec}(R).$$

From [Inc20, Thm. 1.2], for ϵ small enough, the special fiber of such a canonical model will be an elliptic surface $\mathcal{X}^c \rightarrow \mathcal{C}$ with a section \mathcal{S} and irreducible fibers. The image of p via $\overline{\Phi}_n$ will be obtained by contracting $\mathcal{S}^c \subseteq \mathcal{X}^c$, and its special fiber will be obtained by contracting the section \mathcal{S}_0^c of each irreducible component of \mathcal{X}_0^c . It turns out that it is slightly easier to understand \mathcal{X}^c rather than the threefold obtained from it by contracting \mathcal{S}^c , so we will focus on \mathcal{X}^c . To obtain the canonical model $(\mathcal{X}^c, \epsilon \mathcal{S}^c)$, one has to run an MMP and use the abundance theorem; this is worked out in [AB21, Inc20], we report the salient steps.

In *loc. cit.* it is proven that there is a *specific* MMP with scaling that one can run, such that only a specific type of flip is needed, the so-called *flip of La Nave* (see [AB21, Appendix A]; see also [LN02, Thm. 7.1.2], [AB21, Sec. 6.2.2] or [Inc20, 3.2] for a description of the flip of La Nave). More specifically, it is proven in [Inc20, Thm. 6.5] that the specific MMP mentioned above can be factored as

$$(\mathcal{X}, \mathcal{S}) = (\mathcal{X}^{(1)}, \epsilon \mathcal{S}^{(1)}) \dashrightarrow (\mathcal{X}^{(2)}, \epsilon \mathcal{S}^{(2)}) \dashrightarrow \dots \dashrightarrow (\mathcal{X}^{(m)}, \epsilon \mathcal{S}^{(m)}) = (\mathcal{X}^c, \epsilon \mathcal{S}^c)$$

where $f_i : \mathcal{X}^{(i)} \dashrightarrow \mathcal{X}^{(i+1)}$ is either a flip of La Nave, or a divisorial contraction of some irreducible component of $\mathcal{X}_p^{(i)}$. Via these steps, it is proven in [AB21, Section 6] and [Inc20, Cor. 6.7] that the central fiber of $\mathcal{X}^{(i)}$ is a nodal union of irreducible components which are either:

- (1) pseudo-elliptic surfaces, or
- (2) elliptic surfaces.

The flip of La Nave contracts an irreducible component of the special fiber of the section $\mathcal{S}^{(i)}$; this will result in a pseudo-elliptic component attached to a so-called *intermediate fiber* (cf. [AB17, Def. 4.9]).

Now, to control how the special fiber of $(\mathcal{X}^{(i)}, \mathcal{S}^{(i)})$ change after each step of the MMP, we have to control the intersection pairing on some elliptic surfaces; this is the goal of the remaining part of this section.

Lemma 7.6. *Assume that $\mathcal{X}^{(i)} \dashrightarrow \mathcal{X}^{(i+1)}$ is a flip of La Nave, which contracts a section $S_P \subseteq \mathcal{X}^{(i)}$ and flips out the curve $A \subseteq \mathcal{X}^{(i+1)}$. Let S^+ be the section of the elliptic surface given by the irreducible component $E^+ \subseteq \mathcal{X}_0^{(i+1)}$ containing A , and let S^- be the section in the elliptic surface E^- , given by the proper transform of E^+ in $\mathcal{X}_0^{(i)}$. Then one has*

$$(S^-)^2 = (S^+)^2 - \frac{1}{A^2} \quad \text{and} \quad S_P^2 = -\frac{1}{A^2}.$$




FIGURE 1. Special fibers $\mathcal{X}_0^{(i)} \dashrightarrow \mathcal{X}_0^{(i+1)}$ of a flip of La Nave

Proof. Let $\mathcal{S}^{(j)}$ be the proper transform of the section on $\mathcal{X}^{(j)}$ for every j . Then, if S_1, \dots, S_k are the irreducible components of $\mathcal{S}_0^{(j)}$, we have that

$$(\mathcal{S}_\eta^{(j)})^2 = \sum_{\ell=1}^k (\mathcal{S}^{(j)} \cdot S_\ell) = \sum_{\ell=1}^k (\mathcal{S}_\ell^2)_{\mathcal{X}_\ell^{(j)}},$$

where $\mathcal{S}_\eta^{(j)}$ is the restriction of $\mathcal{S}^{(j)}$ to the generic fiber of $\mathcal{X}^{(i)} \rightarrow \text{Spec}(R)$, and the last intersection pairing is computed on each irreducible component $\mathcal{X}_\ell^{(j)}$ of $\mathcal{X}^{(j)}$. The first equality follows from the flatness of $\mathcal{S}^{(j)} \rightarrow \text{Spec}(R)$, and so its self-intersection is constant along the fibers of $\mathcal{X}^{(i)} \rightarrow \text{Spec}(R)$.

Now, from how the flip of La Nave is constructed, there are exactly two irreducible components of $\mathcal{X}^{(i)}$ where f_i is not an isomorphism. One is E^- , the proper transform of E^+ , and the other is P^- , the irreducible component whose section S_P will be contracted after the flip. Denote by S^+ (resp. S^-) the section on E^+ (resp. E^-). Then we have that

$$(S^-)^2 + (S_P)^2 = (S^+)^2.$$

Moreover, there is a morphism $p : E^+ \rightarrow E^-$ which contracts A , and $p_* S^+ = S^-$. So

$$(S^-)^2 = (S^- \cdot p_* S^+) = (p^* S^- \cdot S^+) = (S^+ + \alpha A \cdot S^+).$$

Here, we can compute

$$(S^+ + \alpha A \cdot A) = 0 \quad \text{so} \quad \alpha = \frac{-1}{A^2}.$$

Therefore one has $(S^-)^2 = (S^+)^2 - \frac{1}{A^2}$ as desired. \square

It follows from [Inc20, Thm. 9.9] that when $0 < \epsilon \ll 1$, there are no pseudo-elliptic surfaces among the irreducible components of \mathcal{X}^c . Therefore, all the pseudo-elliptic surfaces which appear after a flip of La Nave will eventually get contracted, and the corresponding intermediate fiber along which the pseudo-elliptic component is attached will become a cuspidal fiber (cf. [AB17]). Moreover, from [Inc20, Prop. 4.14], the type of intermediate fiber that appears after a flip of La Nave is uniquely determined by A^2 . Combining this with [AB17, Thm. 1.1] and [AB21, Table 2], where the authors compute the log-canonical threshold of a cusp in a minimal Weierstrass fibration, we obtain the following result.

Lemma 7.7. *Suppose that $f_i : \mathcal{X}^{(i)} \rightarrow \mathcal{X}^{(i+1)}$ contracts a pseudo-elliptic surface. Let $A \subseteq \mathcal{X}^{(i)}$ be the intermediate component of the intermediate fiber along which is attached the pseudo-elliptic surface contracted*

by f_i . Let A^c be the cuspidal fiber given by the proper transform of A in $\mathcal{X}^{(i+1)}$, and X be the irreducible component of $\mathcal{X}_0^{(i+1)}$ containing A^c . Then we have that

$$-\frac{1}{A^2} = 1 - \text{lct}(X; A^c).$$

Proposition 7.8. Let $(X', S') \rightarrow C'$ be a tsm-stable elliptic surface, and let (X, S) be an irreducible component of X' . Let

- (1) C be the irreducible component of C' such that X maps to C ;
- (2) $f : X \rightarrow C$ be the corresponding map;
- (3) p_1, \dots, p_k be the nodes of C' along C ;
- (4) $F_i := (f^* p_i)^{\text{red}}$, and m_i be the multiplicity of F_i in $f^* p_i$.

If $j : C \rightarrow \overline{M}_{1,1}$ is the corresponding j -map, then one has

$$(S^2) = -\frac{\deg(j)}{12} \quad \text{and} \quad \left(K_X + \sum_{i=1}^k F_i\right) \cdot S = 2g(C) - 2 + k + \frac{\deg(j)}{12}.$$

Moreover, if we denote by Y the surface obtained by replacing F_1, \dots, F_n by cuspidal fibers F_1^c, \dots, F_n^c , and denote by S_Y the proper transform of S on Y , then one has

$$\left(K_Y + \sum_{i=n+1}^k F_i\right) \cdot S_Y = \left(K_X + \sum_{i=1}^k F_i\right) \cdot S - \sum_{i=1}^n \text{lct}(Y; F_i^c)$$

and

$$(S_Y^2) = -\frac{\deg(j)}{12} - \sum_{i=1}^n (1 - \text{lct}(Y; F_i^c)).$$

In particular, when $g(C) = 0$, we have that

$$\begin{aligned} \left(K_X + \sum_{i=1}^k F_i\right) \cdot S &= k - 2 + \frac{\deg(j)}{12}, \quad \text{and} \\ \left(K_Y + \sum_{i=n+1}^k F_i\right) \cdot S &= k - n - 2 + \frac{\deg(j)}{12} + \sum_{i=1}^n (1 - \text{lct}(Y; F_i^c)). \end{aligned}$$

When we say that Y is obtained by replacing F_1, \dots, F_n with cusps, we mean the following. One can take a resolution of X around the intersection point of S and F_i . The resulting surface will admit a morphism to C and if one can contract all the irreducible fiber components not meeting S , the fiber F gets replaced by a singular fiber. For example, it is shown in [AB17] that if one takes the minimal resolution, the new singular fibers are cusps.

Remark 7.9. Proposition 7.8 is the reason it is simpler to work with \mathcal{X}^c rather than with the threefold obtained by contracting $\mathcal{S}^c \subseteq \mathcal{X}^c$: we can use the canonical bundle formula on $\mathcal{X}^{(i)}$, as $\mathcal{X}^{(i)}$ admits a fibration with irreducible fibers which generically are elliptic curves.

Remark 7.10. Observe that in the formula computing (S_Y^2) , if all the fibers of Y are reduced (e.g. when $n = k$), then $(S_Y^2) \in \mathbb{Z}$, since in this case the section of Y is contained in the smooth locus of Y .

Proof of Proposition 7.8. Recall that the fibers of $f : X \rightarrow C$ are either one-pointed genus one stable curves, or the twisted fibers of Section 3.2. In particular, they are irreducible, so from the canonical bundle formula [FL20, Ex. 2.7] one has

$$K_X = f^* \left(K_C + \frac{\deg(j)}{12} \mathcal{O}(1) + \sum_{i=1}^k (1 - m'_i) p_i \right)$$

where $\{p_1, \dots, p_k\}$ are points in C where the fibers of $X \rightarrow C$ could be worse than nodal singularities (which by assumption are supported on the nodes of C'). From canonical bundle formula $m'_i = \text{lct}(X; m_i F_i)$. In [AB17], the authors compute it explicitly, and it is shown that $m'_i = \frac{1}{m_i}$, or in other terms, (X, F_i) is log-canonical.

Similarly, it is proven in [AB19, Prop. 5.3] that

$$\left(K_X + S + \sum_{i=1}^k F_i\right) \cdot S = 2g(S) - 2 + k.$$

Since

$$F_i \cdot S = \frac{1}{m_i} f^{-1}(p_i) \cdot S = \frac{1}{m_i},$$

putting this together leads to

$$2g(S) - 2 + \frac{\deg(j)}{12} + \sum_{i=1}^k (1 - m_i) + (S^2) + \sum_{i=1}^k m_i = 2g(S) - 2 + k$$

and thus

$$(S^2) = -\frac{\deg(j)}{12}.$$

The other equality now follows similarly, and the “moreover” part follows again from the canonical bundle formula. \square

Algorithm 7.11 (explicit construction of $(\mathcal{X}^c, \mathcal{S}^c)$). We are finally ready to explain how to compute the canonical model $(\mathcal{X}^c, \epsilon \mathcal{S}^c)$, starting from $(\mathcal{X}, \mathcal{S})$. From the explicit descriptions of the steps of the MMP, one can proceed as follows. Let S be an irreducible component of $\mathcal{S}^{(i)}$ which is a leaf on the dual graph of $\mathcal{S}^{(i)}$, and which belongs to the irreducible component $X \subseteq \mathcal{X}_0^{(i)}$. We can use Proposition 7.8 to compute $(K_X + \sum F_i) \cdot S$. There are two cases:

- (1) If $(K_X + \sum F_i) \cdot S \leq 0$, we flip the irreducible component X via a flip of La Nave. Since S is a leaf, there is a unique irreducible component $X' \subseteq \mathcal{X}_0^{(i)}$ to which X is attached along a fiber F . The flip of La Nave will replace F with an intermediate fiber. We know that the resulting pseudo-elliptic component P (namely, the proper transform of X) has to be contracted by taking the canonical model $(\mathcal{X}^c, \mathcal{S}^c)$, either by a step of our special MMP (if there is a $(K_{\mathcal{X}^{(i)}} + \epsilon \mathcal{S}^{(i)})$ -extremal ray, whose contraction will contract P) or by taking the canonical model of our minimal model (i.e. by $f^{(m-1)}$). We choose to contract it right away, so that the resulting contraction will replace F with a cusp, whose log-canonical threshold can be computed using Lemma 7.6, and so that each irreducible component of the resulting special fiber will be an elliptic surface with a section and with irreducible fibers (which will allow us to use Proposition 7.8 once again).
- (2) If $(K_X + \sum F_i) \cdot S > 0$ we do nothing and move to the next leaf.

The previous propositions explain how the intersection pairing change after point (1) above, so we can iterate the step above until for each S we have $(K_X + \sum F_i) \cdot S > 0$. At this point, the resulting elliptic surface will be the special fiber of $(\mathcal{X}^c, \epsilon \mathcal{S}^c)$.

Remark 7.12. One might wonder how the algorithm would change if we were to consider other coefficients for the section \mathcal{S} . It turns out that, for that case, the control on the steps of our special MMP will provide other (slightly more complicated) combinatorial invariants that one has to consider; see [Inc20, Thm. 1.4 and Def. 7.3].

7.2. Combinatorial description. The goal of this subsection is to introduce a combinatorial data which will package all the information in Algorithm 7.11. We will first introduce objects that correspond to the locally closed strata of \mathcal{K}_n – the compactification of the Weierstrass locus $\mathcal{W}_n^{\text{nod}}$ by twisted stable maps.

Definition 7.13. A *sliced tree* $\Gamma = (V, E, E_0, \text{jdeg})$ is a graph with vertices V and edges E , which is a tree, together with the following structure. We choose a *jdeg* function

$$\text{jdeg}: V \longrightarrow \frac{1}{12} \mathbb{Z}_{\geq 0},$$

a subset $E_0 \subset E$ of *sliced edges*, and for each $e \in E_0$ connecting v and w , we assign a pair (e_v, e_w) of fractions, called *slicings*, from the following list:

$$\left(\frac{1}{2}, \frac{1}{2}\right), \quad \left(\frac{1}{3}, \frac{2}{3}\right), \quad \left(\frac{1}{4}, \frac{3}{4}\right), \quad \left(\frac{1}{6}, \frac{5}{6}\right).$$

As usual, a vertex $v \in V$ is called a *leaf* if it is adjacent to only one other vertex. We define

$$\text{jdeg}(\Gamma) := \sum_{v \in V} \text{jdeg}(v).$$

Moreover, we define $E(v)$ (resp. $E_0(v)$) to be the set of edges (resp. sliced edges) adjacent to $v \in V$, and we require that for each vertex v we have that

$$\text{jdeg}(v) + \sum_{e \in E_0(v)} e_v \in \mathbb{Z}.$$

Definition 7.14. We say that a sliced tree $\Gamma = (V, E, E_0, \text{jdeg})$ is *tsm-stable* if

- (1) $\text{jdeg}(v) \geq 0$ for all $v \in V$; and
- (2) $|E(v)| \geq 3$ if $\text{jdeg}(v) = 0$.

Definition 7.15. Let $(X, S) \rightarrow C$ be a tsm-stable elliptic surface. The *sliced tree* Γ associated to $(X, S) \rightarrow C$ is given as follows.

- (1) Each irreducible component of X corresponds to a vertex of Γ .
- (2) The jdeg of each vertex is the degree of the corresponding j -map divided by 12.
- (3) There is an edge between vertices v and w if the corresponding irreducible components intersect.
- (4) The edge connecting v and w is sliced if the corresponding irreducible components are glued along a non-reduced fiber.
- (5) The slicing is defined as follows. If two irreducible components X_v and X_w of X intersect along a non-reduced fiber F , let $Y_v \rightarrow X_v$ (resp. $Y_w \rightarrow X_w$) be the minimal resolution of X_v (resp. X_w) along $S \cap F$, and let $Z_v \rightarrow Y_v$ (resp. $Z_w \rightarrow Y_w$) be the surface obtained by contracting all the fiber components not meeting S . Then the fiber F is replaced with a different fiber $F_v \subseteq Z_v$ (resp. $F_w \subseteq Z_w$). The slicing adjacent to v (resp. w) is $1 - \text{lct}(Z_v; F_v)$ (resp. $1 - \text{lct}(Z_w; F_w)$).

Remark 7.16.

- (a) The surfaces Z_v and Z_w of Definition 7.15 (5) can also be constructed by taking the minimal Weierstrass fibration birational to X_v and X_w respectively, if X_v and X_w are normal. The fibers F_v and F_w would be the cuspidal fibers which replace F .
- (b) The condition $\text{jdeg}(v) + \sum_{e \in E_0(v)} e_v \in \mathbb{Z}$ follows from Remark 7.10.
- (c) One might wonder why the possible markings are only those listed in Definition 7.13, for example, why $(\frac{1}{2}, \frac{1}{6})$ is not allowed. This follows from Definition 7.15 (4). Indeed, it turns out that if the action is of the type $(x, y) \mapsto (\xi \cdot x, \xi^{-1} \cdot y)$, then the only slicing allowed are those of Definition 7.13.

The following proposition is now straightforward.

Proposition 7.17. *The stack \mathcal{K}_n admits a locally closed stratification with strata labeled by the stable sliced trees of $\text{jdeg}(\Gamma) = n$:*

$$\mathcal{K}_n = \bigsqcup_{\Gamma, \text{jdeg}(\Gamma)=n} \mathcal{K}_\Gamma.$$

Remark 7.18. Table 1 explains what is the slicing (and so $1 -$ the minimal log-canonical threshold) of each type of cusp which appears in a minimal Weierstrass fibration. Moreover, if one replaces a cuspidal fiber $f^{-1}(p)$ with a twisted fiber as in [AB17], the corresponding elliptic surface in a neighborhood of p comes from a twisted stable map $\phi: \mathcal{C} \rightarrow \overline{\mathcal{M}}_{1,1}$. From the analysis in [AB17] and [BPS22, Thm. 1.6 & Sec. 7], the denominator d of each slicing corresponds to the order of the stabilizers of $p \in \mathcal{C}$, and the numerator comes from the character of the irreducible representation of μ_d on $\phi^*\mathcal{O}(1)$. For example, type III* cusps get replaced with twisted fibers such that \mathcal{C} has automorphism group μ_4 , with the action on the fibers of $\phi^*\mathcal{O}(1)$ being $\zeta \cdot v = \zeta^3 v$.

Remark 7.19. Some of these strata may be empty. For example, one can consider a tree with a vertex whose jdeg equal to $\frac{1}{6}$, and with 5 edges exiting from it, each sliced with coefficient $\frac{1}{6}$. This should correspond

TABLE 1. Correspondence between Kodaira fiber types and slicings

Kodaira type	Slicing, or $1 - \text{lt}$	Kodaira type	Slicing, or $1 - \text{lt}$
I	0	I*	$\frac{1}{2}$
II	$\frac{1}{6}$	II*	$\frac{5}{6}$
III	$\frac{1}{4}$	III*	$\frac{3}{4}$
IV	$\frac{1}{3}$	IV*	$\frac{2}{3}$


to an elliptic surface, with j -map of degree 2, and with five non-reduced fibers. One can replace this elliptic surface with its minimal Weierstrass fibration, and one would have a Weierstrass fibration with five type II cusps and j -map of degree 2. But such a Weierstrass fibration would be given by

$$A \in H^0(\mathcal{O}_{\mathbb{P}^1}(4)) \quad \text{and} \quad B \in H^0(\mathcal{O}_{\mathbb{P}^1}(6)),$$

where the polynomial A should vanish along the five points corresponding to cuspidal fibers. This forces A to be identically 0. By [BPS22, Thm. 1.6], if the edges are sliced with coefficient $\frac{1}{6}$, the corresponding cusp will be such that B cannot have a double root. So B should vanish at six distinct points, and therefore there should be six cusps rather than five, which is the desired contradiction.

Example 7.20. Consider a tsm-stable elliptic surface $(\pi: X \rightarrow C, S)$, whose associated sliced tree Γ is as on the right. Then it should be interpreted as providing the following information.

- C has 6 irreducible components C_v , $v \in \{a, b, c, d, f, h\}$, each isomorphic to \mathbb{P}^1 .
- $\text{jdeg}(\Gamma) = 6$, hence $g \in \mathcal{K}_6$, i.e. it is a degeneration of a Weierstrass fibration whose j -map has degree 72.
- The degree of the j -map restricted to C_v and divided by 12 is the label $\text{jdeg}(v)$ of the vertex v .



In this example, the sliced edges are (bc) , (bd) and (dh) , where the slicings are $(\frac{1}{6}, \frac{5}{6})$, $(\frac{1}{2}, \frac{1}{2})$ and $(\frac{1}{2}, \frac{1}{2})$ respectively.

In order to describe boundary strata in the closure of the KSBA moduli space, we need to contract certain irreducible components of surfaces via Algorithm 7.11. This will reduce the number of vertices, but create singularities on the surfaces, which can be tracked combinatorially by attaching half-edges. Since the contraction process resembles the process of pruning trees, we name the corresponding object accordingly.

Definition 7.21. A *pruned tree* $\Pi = (V, E, E_0, \text{jdeg}, F, T)$ is a sliced tree (V, E, E_0, jdeg) together with a set K , called *klt-markings*, a set L called *lc-markings*, and the following additional structure. Each klt-marking and lc-marking is attached to one vertex, and similar to the notation in Definition 7.13, we denote the set of klt-markings (resp. lc-markings) attached to a vertex $v \in V$ by $T(v)$ (resp. $F(v)$). Each klt-marking $t \in T(v)$ attached to $v \in V$ is assigned a number

$$t_v \in \left\{ \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{3}{4}, \frac{1}{6}, \frac{5}{6} \right\}.$$

Remark 7.22. The numbers in Definition 7.21 are $(1 - \text{lt of cuspidal fibers})$ in minimal Weierstrass fibrations, see [AB17].

Definition 7.23. The *weight* of a vertex v of a pruned tree is defined as follows:

$$\text{wt}(v) := \#\{\text{edges adjacent to } v\} - 2 + \text{jdeg}(v) + \sum_{t_v \in T(v)} t_v + |F(v)|.$$

Definition 7.24. A pruned tree $\Pi = (V, E, E_0, \text{jdeg}, F, T)$ is *stable* if each vertex $v \in V$ has $\text{wt}(v) > 0$.

Remark 7.25. We now explain the reason behind the previous definition. We will associate to each KSBA-limit, as in Section 7.1, a pruned tree Π . More explicitly, if $(\mathcal{X}, S) \rightarrow \text{Spec}(R)$ is a one-parameter family of

t-sm-stable elliptic surfaces, whose generic fiber is a Weierstrass fibration, we explained in Section 7.1 how to take the relative canonical model of $(\mathcal{X}, \epsilon\mathcal{S}) \rightarrow \text{Spec}(R)$. If Γ is the sliced tree associated to $(\mathcal{X}_0, \mathcal{S}_0)$, each step of the algorithm flips and contracts a component of $(\mathcal{X}_0, \mathcal{S}_0)$ and that corresponds to removing a leaf of Γ and replacing it with an lc cusp or a klt cusp. From Algorithm 7.11, the algorithm terminates when each irreducible component of such a canonical model will satisfy that $(K_X + \sum F_i).S > 0$, and from Proposition 7.8 this translates into a condition that $\text{wt}(v) > 0$ for every vertex in the resulting graph. Moreover, the lc markings (resp. klt markings) should correspond to strictly lc cusps (resp. klt cusps).

The discussion in Algorithm 7.11 leads to the following.


Proposition 7.26. *Let $\bar{\Phi}_n: \mathcal{K}_n^\nu \rightarrow \bar{\mathcal{E}}_n$ be the same as in Proposition 7.3. Let $\Gamma = (V, E, E_0, \text{jdeg})$ be a stable sliced tree parametrizing a stratum \mathcal{S}_Γ in \mathcal{K}_n^ν . Let Π be obtained from Γ by the following process of pruning:*

- (1) Set $\Pi = \Gamma$ with $F = T = \emptyset$.
- (2) If there is a leaf $v \in V$ of weight ≤ 0 , denote by w the vertex adjacent to it, and do the following:
 - (a) If $t = 1$ then replace $F(w)$ by $F(w) \cup \{v\}$;
 - (b) if $t = 0$, do nothing; otherwise
 - (c) replace $T(w)$ by $T(w) \cup \{\text{wt}(v) + 1\}$.
 If there are no leaves of weight ≤ 0 , stop.
- (3) Repeat (2).


The resulting pruned tree, denoted $\bar{\Phi}_n(\Gamma) := \Pi$, is stable as a pruned tree. Moreover, the image of the stratum $\bar{\Phi}_n(\Gamma)$ parametrizes surface pairs (X, S) with sliced tree given by Π and whose components have lc cusps (resp. klt cusps) of given type indexed by $F(w)$ (resp. $T(w)$).

Since $\bar{\Phi}_n$ is surjective, every point of $\bar{\mathcal{E}}_n$ is in the image of some stratum. Thus, the combinatorial data of any surface pair (X, S) parametrized by $\bar{\mathcal{E}}_n$ is encoded by a stable pruned graph $\bar{\Phi}_n(\Gamma) = \Pi$ for some stable sliced graph Γ .

Example 7.27. Let Γ be the sliced tree exhibited in Example 7.20. Then the pruned tree $\bar{\Phi}_6(\Gamma)$ is the one on the right. Every leaf of the pruned tree on the right has positive weight, hence it is stable and does not need to be pruned any further.



Example 7.28. We now give another example of pruning, where each squiggly arrow corresponds to pruning all outermost leaves once. The sliced edges have coefficients $(\frac{1}{2}, \frac{1}{2})$, $(\frac{1}{2}, \frac{1}{2})$, $(\frac{1}{3}, \frac{2}{3})$, $(\frac{1}{6}, \frac{5}{6})$ respectively, the edges labelled with (y) represent a klt marking with marking y , whereas the edges labelled with an x represent lc markings.



Proof of Proposition 7.26. As mentioned above, the klt-markings (resp. lc-markings) of a given vertex, correspond to the klt (resp. lc) cuspidal fibers in the corresponding irreducible components. From Proposition 7.8, the weight of a vertex v on a pruned tree corresponds to the intersection number $(K_{\mathcal{X}^{(i)}}. S_v)$, where S_v is the irreducible component of $\mathcal{S}_0^{(i)}$ contained in the irreducible component of \mathcal{X}_0 corresponding to the vertex v . Therefore from Algorithm 7.11, the only thing we need to check is the following. Let v be a leaf with non-positive weight, attached to a vertex w , and let X_w^- be the irreducible component corresponding to v with X_w^+ the irreducible component attached to it. Then after a flip of La Nave and a contraction of the pseudo-elliptic component, if we denote by X_v^+ the proper transform of X_j^- , and by F the cuspidal fiber to which the pseudo-elliptic surface given by the proper transform of X_v^+ is contracted, then we claim that

$$1 - \text{lct}(X_w^{\text{cusp}}; F) = \text{wt}(i) + 1$$

where X_w^{cusp} is the proper transform of X_w^- containing F .




FIGURE 2. Notations as in the proof of Proposition 7.26.


This is because from Lemma 7.7, the log-canonical threshold is given by the reciprocal of the self-intersection of a certain intermediate component, which from Lemma 7.6 is given (up to sign) by (S_v^2) , where $S_v \subseteq X_v$ is the section component. Since X_v is attached only along a single fiber, the desired statement now follows from Proposition 7.8. \square

Corollary 7.29. *The value $\sum_v \text{wt}(v)$ remains constant during the pruning process.*

Proof. This follows from how the algorithm of Proposition 7.26 works. \square

Corollary 7.30. *Let X be a KSB-stable surface parametrized by $\overline{\mathcal{P}}_3$, and Π be the pruned tree associated to X . Then Π has two leaves (i.e. Π is a chain), and has at most six vertices.*

Observe that one can construct an element with six irreducible components. It suffices to construct a twisted stable map with the following tsm-stable graph Γ , and with slicings $(\frac{5}{6}, \frac{1}{6}), (\frac{2}{3}, \frac{1}{3}), (\frac{1}{2}, \frac{1}{2}), (\frac{1}{3}, \frac{2}{3}), (\frac{1}{6}, \frac{5}{6})$ in order from left to right.



Such a twisted stable map exists, or in other terms, the stratum in \mathcal{K}_3 corresponding to the graph above is not empty; we briefly sketch how to construct it. First, for each vertex v of the diagram above, we consider a map from a root-stack of $\mathbb{P}^1 \rightarrow \overline{\mathcal{M}}_{1,1}$ such that the corresponding map on coarse moduli spaces has degree $12 \text{jdeg}(v)$. Then we glue the corresponding maps along the stacky points, in a way so that the resulting morphism is balanced.

For example, for the first two vertices on the left, one can proceed as follows. For the leftmost vertex, we start by considering an elliptic K3 which we denote by $Y \rightarrow \mathbb{P}^1$, with a single klt cusp, and from the slicing and Table 1, this should be of type II*. From [Mir89, Table (IV).3.1], one can construct it explicitly via its Weierstrass equation, taking two polynomials

$$A \in H^0(\mathcal{O}_{\mathbb{P}^1}(8)), B \in H^0(\mathcal{O}_{\mathbb{P}^1}(12))$$

which have all distinct and single roots, except for a single point $p \in \mathbb{P}^1$ where A has a zero of multiplicity 4, and B has multiplicity 5. This is clearly possible, and the two polynomials A and B will induce a morphism $\mathbb{P}^1 \rightarrow [\mathbb{A}^2/\mathbb{G}_m]$, with the action with weights 4 and 6, inducing $f: \mathbb{P}^1 \setminus \{p\} \rightarrow \overline{\mathcal{M}}_{1,1}$. By [BPS22, Thm. 1.6 & Thm. 3.3] (see also [BV24]), one can construct a root stack $\mathcal{P}^1 \rightarrow \mathbb{P}^1$ so that f extends to $\mathcal{P}^1 \rightarrow \overline{\mathcal{M}}_{1,1}$.

We can check that \mathcal{P}^1 has μ_6 as automorphism group on the stacky point. Indeed, the two sections A and B give a map $\mathbb{P}^1 \rightarrow [\mathbb{A}^2/\mathbb{G}_m]$ which locally is of the form $z \mapsto (z^4, z^5)$, and its \mathbb{G}_m -orbit in \mathbb{A}^2 is of the form

$$\mathbb{G}_m \times \mathbb{P}^1 \setminus \{p\} \rightarrow \mathbb{A}^2, (\lambda, z) \mapsto (\lambda^4 z^4, \lambda^6 z^5).$$

Extending this map to \mathbb{P}^1 boils down to replacing λ with $z^{\frac{m}{d}}$, so that the two sections $(z^{4+\frac{4m}{d}}, z^{5+\frac{6m}{d}})$ do not have a pole at p and do not vanish simultaneously. The smallest positive d that one can take will lead to a representable morphism $\mathbb{P}^1 \rightarrow \overline{\mathcal{M}}_{1,1}$, and it is $d=6$ with $m=-5$; so \mathcal{P}^1 will have a μ_6 as stabilizer group on the stacky point. Since the morphism $\mathbb{P}^1 \rightarrow \overline{\mathcal{M}}_{1,1}$ is representable, such a stacky point will go to the only point of $\overline{\mathcal{M}}_{1,1}$ with μ_6 as automorphisms. Similarly, for the second leftmost vertex, one can consider two homogeneous polynomials A, B of degree 4 and 6 respectively, which have distinct and single roots on \mathbb{P}^1 , except at two points p_1 and p_2 . From the slicings, Table 1 and [Mir89, Table (IV).3.1], we require that at p_1 the polynomials A and B must vanish of order 1 (this will correspond to a type II fiber), and at p_2 instead A should vanish with multiplicity 3 and B with multiplicity 4. This is again possible, and proceeding as before this will lead to a stacky \mathbb{P}^1 , which we denote by \mathcal{P}_2^1 , with two stacky points. As before, one can check that \mathcal{P}_2^1 has, at p_1 , automorphism group which is μ_6 , and at p_2 which is μ_3 . We can glue \mathcal{P}^1 and \mathcal{P}_2^1 along the point with automorphism μ_6 , and the corresponding maps to $\overline{\mathcal{M}}_{1,1}$ will glue. On the glued point, one can check that we will have a twisted vertex. We can proceed with this recipe to get the desired twisted map $\mathcal{C} \rightarrow \overline{\mathcal{M}}_{1,1}$.

Proof of Corollary 7.30. We begin by observing the following:

$$\begin{aligned} \text{wt}(\Pi) &= \text{wt}(\Gamma) \\ &= \sum_{i \in V(\Gamma)} (\#\{\text{edges containing } i\} - 2 + \text{jdeg}(i)) \\ &= \sum_{i \in V(\Gamma)} (\#\{\text{edges containing } i\} - 2) + \text{jdeg}(\Gamma) \\ &= \text{jdeg}(\Gamma) - 2 = 1, \end{aligned}$$

where we used that the Euler characteristic of Γ , defined by $\#\{\text{vertices}\} - \#\{\text{edges}\}$, is equal to 1 since Γ is a tree. As a consequence, one has

$$(11) \quad \sum_{i \in V(\Pi)} \left(\text{jdeg}(i) + \sum_{t_j \in T(i)} t_j + |F(i)| \right) = 3.$$

Let us prove that Π can have at most two leaves. If not, let ℓ, j, k three leaves. The stability condition guarantees that

$$\begin{aligned} \text{jdeg}(\ell) + \sum_i |F(\ell)| + \sum_{i \in T(\ell)} t_i &> 1, \\ \text{jdeg}(j) + \sum_i |F(j)| + \sum_{i \in T(j)} t_i &> 1 \\ \text{jdeg}(k) + \sum_i |F(k)| + \sum_{i \in T(k)} t_i &> 1. \end{aligned}$$

Since the graph is connected, there is no edge between i, j and ℓ . Then if we add up the previous inequalities, we get a contradiction of Equation (11). We conclude that Π has at most two edges, and thus Π is a chain. We now prove that there cannot be more than four internal vertices in the chain.

Let's label the edges on the chain as follows: let l_1, l_2 be the leaves and n_1, \dots, n_k the internal vertices. We know that

$$\text{jdeg}(l_i) + \sum_i |F(l_i)| + \sum_{i \in T(l_i)} t_i > 1$$

for each leaf $i = 1, 2$, and from how the numbers t_i are defined, we have that

$$\text{jdeg}(l_i) + \sum_i |F(l_i)| + \sum_{i \in T(l_i)} t_i \geq \frac{7}{6}.$$

Similarly for $1 \leq i \leq k$,

$$\text{jdeg}(n_i) + \sum_i |F(n_i)| + \sum_{i \in T(n_i)} t_{n_i} \geq \frac{1}{6}.$$

Combining this with Equation (11), we get that $k \leq 4$. \square

More generally, the previous proof shows that the quantity

$$\sum_{i \in V(\Pi)} \left(\text{jdeg}(i) + \sum_{t_j \in T(i)} t_j + |F(i)| \right) = \text{wt}(\Pi) + 2 = \text{jdeg}(\Gamma)$$

whenever $\overline{\Phi}_n(\Gamma) = \Pi$. The latter quantity is constant in families of twisted stable maps and thus agrees with the height of a surface parametrized by the generic point of \mathcal{K}_n . This motivates the following.

Definition 7.31. The *height* of a stable pruned tree is defined by

$$\text{ht}(\Pi) = \sum_{i \in V(\Pi)} \left(\text{jdeg}(i) + \sum_{t_j \in T(i)} t_j + |F(i)| \right) \in \mathbb{N}.$$

Proposition 7.32. *The pruned tree associated to any KSB-stable surface parametrized by $\overline{\mathcal{P}}_n$ has height n .*

REFERENCES

- [AB17] Kenneth Ascher and Dori Bejleri, *Log canonical models of elliptic surfaces*, *Advances in Mathematics* **320** (2017), 210–243.
- [AB19] ———, *Moduli of fibered surface pairs from twisted stable maps*, *Math. Ann.* **374** (2019), no. 1-2, 1007–1032.
- [AB21] ———, *Moduli of weighted stable elliptic surfaces and invariance of log plurigenera*, *Proc. Lond. Math. Soc.* (3) **122** (2021), no. 5, 617–677. With an appendix by Giovanni Inchiostro.
- [AB23] ———, *Smoothability of relative stable maps to stacky curves*, *Épjournal Géom. Algébrique* **7** (2023), Art. 2, 22.
- [ABIP23] Kenneth Ascher, Dori Bejleri, Giovanni Inchiostro, and Zsolt Patakfalvi, *Wall crossing for moduli of stable log varieties*, *Annals of Mathematics*, to appear (2023).
- [AH11] Dan Abramovich and Brendan Hassett, *Stable varieties with a twist*, *Classification of algebraic varieties*, 2011, pp. 1–38.
- [AP23] Valery Alexeev and Rita Pardini, *Explicit compactifications of moduli spaces of Campedelli and Burniat surfaces*, arXiv preprint arXiv:0901.4431 (2023).
- [AV02] Dan Abramovich and Angelo Vistoli, *Compactifying the space of stable maps*, *Journal of the American Mathematical Society* **15** (2002), no. 1, 27–75.
- [AV97] ———, *Complete moduli for fibered surfaces* **Recent progress in intersection theory (Bologna, 1997)** (1997).
- [BHPS13] Bhargav Bhatt, Wei Ho, Zsolt Patakfalvi, and Christian Schnell, *Moduli of products of stable varieties*, *Compositio Mathematica* **149** (2013), no. 12, 2036–2070.
- [BPS22] Dori Bejleri, Jun-Yong Park, and Matthew Satriano, *Height moduli on cyclotomic stacks and counting elliptic curves over function fields*, 2022. Available at <https://arxiv.org/abs/2210.04450>.
- [BPVdV84] Wolf Barth, Chris Peters, and Antonius Van de Ven, *Compact complex surfaces*, Springer, 1984.
- [BV23] Giulio Bresciani and Angelo Vistoli, *An arithmetic valuative criterion for proper maps of tame algebraic stacks*, *Manuscripta Mathematica* (2023), 1–11.
- [BV24] ———, *An arithmetic valuative criterion for proper maps of tame algebraic stacks*, *manuscripta mathematica* **173** (2024), no. 3, 1061–1071.
- [CDLI24] Samir Canning, Andrea Di Lorenzo, and Giovanni Inchiostro, *The integral chow rings of moduli of Weierstrass fibrations*, *Transactions of the American Mathematical Society* (2024).
- [DM69] P. Deligne and D. Mumford, *The irreducibility of the space of curves of given genus*, *Inst. Hautes Études Sci. Publ. Math.* **36** (1969), 75–109.
- [FL20] Enrica Floris and Vladimir Lazić, *A travel guide to the canonical bundle formula*, *Birational Geometry and Moduli Spaces* (2020), 37–55.
- [Fuj14] Osamu Fujino, *Fundamental theorems for semi log canonical pairs*, *Algebr. Geom.* **1** (2014), no. 2, 194–228.
- [GKKP11] Daniel Greb, Stefan Kebekus, Sándor J. Kovács, and Thomas Peternell, *Differential forms on log canonical spaces*, *Publ. Math. Inst. Hautes Études Sci.* **114** (2011), 87–169.
- [Gro66] Alexander Grothendieck, *Éléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné): IV. étude locale des schémas et des morphismes de schémas, troisième partie*, *Inst. Hautes Études Sci. Publ. Math.* **28** (1966).
- [Hac04] Paul Hacking, *Compact moduli of plane curves*, *Duke Math. J.* **124** (2004), no. 2, 213–257.

- [Har13] Robin Hartshorne, *Algebraic geometry*, Vol. 52, Springer Science & Business Media, 2013.
- [Inc20] Giovanni Inchiostro, *Moduli of Weierstrass fibrations with marked section*, *Advances in Mathematics* **375** (2020), 107374.
- [KM98] János Kollár and Shigefumi Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [Kol09a] János Kollár, *Hulls and husks*, 2009. Available at <https://arxiv.org/abs/0805.0576>.
- [Kol09b] ———, *Lectures on resolution of singularities (am-166)*, Vol. 166, Princeton University Press, 2009.
- [Kol23] ———, *Families of varieties of general type*, Vol. 231, Cambridge University Press, 2023.
- [KSB88] J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, *Invent. Math.* **91** (1988), no. 2, 299–338.
- [Liu12] Wenfei Liu, *Stable degenerations of surfaces isogenous to a product II*, *Trans. Amer. Math. Soc.* **364** (2012), no. 5, 2411–2427.
- [LN02] Gabriele La Nave, *Explicit stable models of elliptic surfaces with sections*, 2002.
- [Mir81] Rick Miranda, *The moduli of Weierstrass fibrations over \mathbf{P}^1* , *Math. Ann.* **255** (1981), no. 3, 379–394.
- [Mir89] ———, *The basic theory of elliptic surfaces*, *Dottorato di Ricerca in Matematica*. [Doctorate in Mathematical Research], ETS Editrice, Pisa, 1989.
- [MZ23] Fanjun Meng and Ziquan Zhuang, *MMP for locally stable families and wall crossing for moduli of stable pairs*, 2023. Available at <https://arxiv.org/abs/2311.01319>.
- [PS21] Jun-Yong Park and Johannes Schmitt, *Arithmetic geometry of the moduli stack of Weierstrass fibrations over \mathbb{P}^1* , 2021. Available at <https://arxiv.org/abs/2107.12231>.
- [Rol10] Sönke Rollenske, *Compact moduli for certain Kodaira fibrations*, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **9** (2010), no. 4, 851–874.
- [Sei87] Wolfgang K. Seiler, *Global moduli for elliptic surfaces with a section*, *Compositio Math.* **62** (1987), no. 2, 169–185.
- [SS10] Matthias Schütt and Tetsuji Shioda, *Elliptic surfaces*, *Algebraic geometry in East Asia—Seoul 2008, 2010*, pp. 51–160.
- [Sta24] The Stacks project authors, *The stacks project*, 2024.
- [vO06] Michael van Opstall, *Stable degenerations of surfaces isogenous to a product of curves*, *Proc. Amer. Math. Soc.* **134** (2006), no. 10, 2801–2806.