

ROOT STACK VALUATIVE CRITERION FOR GOOD MODULI SPACES

DORI BEJLERI, GIOVANNI INCHIOSTRO, AND MATTHEW SATRIANO

ABSTRACT. We prove a root stack valuative criterion for good moduli space maps and for gerbes for reductive groups under some mild assumptions on the residue characteristic. We give several applications to parahoric extension for torsors, rational points on stacks, gerbes and homogeneous spaces, and the geometry of fibrations.

1. INTRODUCTION

The goal of this paper is to prove the following root stack valuative criterion for good moduli spaces.

Theorem 1.1. *Let $\mathcal{X} \rightarrow X$ be a good moduli space where \mathcal{X} is an Artin stack with affine diagonal and of finite type over a locally Noetherian scheme S . Let R be a DVR with fraction field K and residue field k . Given a commutative diagram of solid arrows*

$$\begin{array}{ccc} \mathrm{Spec} K & \xrightarrow{\quad} & \mathcal{X} \\ \downarrow & \nearrow \text{dashed} & \downarrow \\ \sqrt[n]{\mathrm{Spec} R} & \dashrightarrow & \mathrm{Spec} R \xrightarrow{\quad} X \end{array}$$

there exists a root stack $\sqrt[n]{\mathrm{Spec} R} \rightarrow \mathrm{Spec} R$ and dotted arrows filling in the diagram. Moreover, if $\mathrm{Spec} K \rightarrow \mathcal{X}$ maps to the closed point of $\mathcal{X} \times_X K$, then one can choose an extension such that also the closed point of $\sqrt[n]{\mathrm{Spec} R}$ maps to the closed point of $\mathcal{X} \times_X k$.

Good moduli spaces for algebraic stacks were introduced by Alper in his thesis [Alp13], and are a generalization both of coarse moduli spaces for Deligne-Mumford stacks in characteristic 0, and of morphisms, appearing in the context of GIT, of the form $[W(\mathcal{L})^{ss}/G] \rightarrow W//_{\mathcal{L}}G$, where W is a projective variety with an action of a linearly reductive group G , \mathcal{L} is a G -linearized ample line bundle on W and $W(\mathcal{L})^{ss}$ is its semistable locus. On a first approximation, these are morphisms $\mathcal{X} \rightarrow X$ which are of the form $[\mathrm{Spec}(A)/\mathrm{GL}_N] \rightarrow \mathrm{Spec}(A^{\mathrm{GL}_N})$ étale locally on X rather than Zariski locally, as in classical GIT [AHR25, Theorem 6.1]. Good moduli space maps if they exist are universal for maps to algebraic spaces and have very strong topological and cohomological properties. They are a fundamental tool in the study of algebraic stacks with positive dimensional stabilizers.

The key difference between the root stack valuative criterion Theorem 1.1 and the usual existence part of the valuative criterion for a good moduli space [AHLH23, Theorem A.8] is that both the fraction field and the residue field of R are preserved by the extension $\sqrt[n]{\mathrm{Spec} R} \rightarrow \mathrm{Spec} R$. This allows for many arithmetic applications (Section 1.1) and for globalizing the valuative criterion from a DVR to a global curve C (Proposition 1.5 below). For Deligne-Mumford stacks over \mathbb{C} , the analogous valuative criterion has been used extensively in the theory of twisted stable maps [AV02, ACV03] and fibered surfaces [AB19]. Recently it was generalized by Bresciani and Vistoli [BV24a] to tame stacks in any characteristic and appears in numerous applications including to heights of rational points on tame stacks [ESZB23, BPS24, DY22], resolution of indeterminacy for rational maps to stacks [Jeo25], and classification of singular fibers in fibrations [Rey25].

Date: July 11, 2025.

DB was partially supported by NSF grant DMS-2401483.

MS was partially supported by a Discovery Grant from the National Science and Engineering Research Council of Canada and a Mathematics Faculty Research Chair from the University of Waterloo.

Beyond the realm of algebraic stacks with finite inertia, partial progress was made in [DLI22, Theorem 3.9] and [DLI24, Theorem 1.2]. For the stack of K -semistable Fano varieties and of boundary polarized Calabi-Yau pairs, a proof of properness that shows one only needs to adjoin a root of the uniformizer of R was given in [BLXZ25] and [ABB⁺23, Remark 7.2] respectively, though not necessarily that this can be done μ_n -equivariantly as in Theorem 1.1. See also [SU24] for related results in the context of motivic integration.

The fact that \mathcal{X} has a good moduli space is used to reduce to the case that \mathcal{X} is a gerbe for a smooth connected linearly reductive group scheme but in that case we prove the following result which in positive and mixed characteristic is more general than the setting of a good moduli space. Recall that a group scheme G is special in the sense of Serre if all G -torsors can be trivialized Zariski locally.

Remark 1.2. We can factor the resulting morphism $\sqrt[n]{\mathrm{Spec} \bar{R}} \rightarrow \mathcal{X}$ through its relative coarse moduli space [AOV11, Theorem 3.1] which is necessarily a root stack of possibly smaller degree [BV24a, Proposition 3.12] so without loss of generality we can assume that this filling is representable.

Theorem 1.3. *Let R and K be as above and let $\mathcal{X} \rightarrow \sqrt[n]{\mathrm{Spec} \bar{R}}$ be a gerbe for a reductive group scheme $G \rightarrow \sqrt[n]{\mathrm{Spec} \bar{R}}$ over a root stack of $\mathrm{Spec} R$. Suppose either that G is special, or that the order of the Weyl groups of the fibers of $G \rightarrow \sqrt[n]{\mathrm{Spec} \bar{R}}$ is coprime to the residue characteristic, or G is an extension of the form $1 \rightarrow G_1 \rightarrow G \rightarrow G_2 \rightarrow 1$ such that the Weyl groups of the fibers of $G_1 \rightarrow \sqrt[n]{\mathrm{Spec} \bar{R}}$ have order coprime to the residue characteristic, and such that G_2 is special. Then any K -point $x_K : \mathrm{Spec} K \rightarrow \mathcal{X}$ extends to a section $x : \mathcal{R} \rightarrow \mathcal{X}$ where $\mathcal{R} \rightarrow \sqrt[n]{\mathrm{Spec} \bar{R}}$ is a further root stack.*

Remark 1.4. The above theorem is the first step in generalizing Theorem 1.1 to the case of *adequate moduli spaces* [Alp14], at least under some assumptions. In characteristic 0, adequate and good moduli spaces are equivalent but in positive and mixed characteristic, adequate moduli spaces are a larger class. For example, if G is reductive and the residue characteristic is positive, then $BG \rightarrow S$ is an adequate moduli space but usually not a good moduli space. From Example 4.3 we can see that the following conditions on the stabilizer G at a closed points seem to be necessary: the reduced connected $(G_0)_{red} \subset G$ is as in Theorem 1.3 and the component group $G/(G_0)_{red}$ is a tame finite group scheme. Even if we impose these assumptions, there are two key places where the proof of Theorem 1.1 uses the good moduli space rather than adequate moduli space assumption: 1) the application of canonical reduction of stabilizers [ER21] in Theorem 3.3 and in particular the fact that the reduced connected component of the identity behaves well in families, and 2) the replacement for Kempf's Theorem when K is not perfect in Lemma 3.1.

Next we globalize the valuative criterion to show that rational maps from a global curve to an Artin stack with a proper good moduli space can be extended after a root stack. Proposition 1.5 yields the existence of *tuning stacks* for K -points of \mathcal{X} in the sense of [ESZB23] and is the first step in developing the theory of heights on stacks for Artin stacks with a good moduli space.

Proposition 1.5. *Suppose C is a locally excellent, regular, 1-dimensional scheme (e.g. a smooth curve over a field, or a Dedekind domain flat over \mathbb{Z}). Let \mathcal{X} be a stack as in Theorem 1.1 with proper good moduli space or a gerbe over a root stack of C as in Theorem 1.3. Then any rational map $\varphi : C \dashrightarrow \mathcal{X}$ extends to a morphism $f : \mathcal{C} \rightarrow \mathcal{X}$ from some root stack $\mathcal{C} \rightarrow C$ along the complement of a domain of definition for φ .*

Corollary 1.6. *Let C be as in Proposition 1.5 and let $K = k(C)$ be its fraction field. Let \mathcal{X} be a stack as in Theorem 1.1 with a proper good moduli space or a gerbe over a root stack of C as in Theorem 1.3. Then any K -point $x_K : \mathrm{Spec} K \rightarrow \mathcal{X}$ extends to a morphism $\mathcal{C} \rightarrow \mathcal{X}$ from some root stack of C .*

Finally, Theorem 1.1 combined with the existence criterion of [AHLH23] immediately implies the following relative version. We refer the reader to *loc.cit.* for the definitions of Θ -reductive and S -complete morphisms.

Corollary 1.7. *Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be a finite type morphism of Artin stacks. Suppose that*

- (i) π has affine diagonal,
- (ii) the relative inertia groups at all closed points of the fibers of π are linearly reductive,
- (iii) π is Θ -reductive,
- (iv) π is S -complete, and
- (v) π satisfies the existence part of the valuative criterion of properness.

Then for any commutative diagram of solid arrows below where R is a DVR with fraction field K

$$\begin{array}{ccccc}
 & & \text{Spec } K & \longrightarrow & \mathcal{X} \\
 & \swarrow \text{dashed} & \downarrow & \searrow \text{dashed} & \downarrow \\
 \sqrt[n]{\text{Spec } R} & \dashrightarrow & \text{Spec } R & \longrightarrow & \mathcal{Y}
 \end{array}$$

there exists a root stack $\sqrt[n]{\text{Spec } R}$ and dotted arrows filling in the diagram.

Remark 1.8. Note that assumption (ii) follows from the other assumptions in characteristic 0.

1.1. Applications. Throughout this section, R is a DVR with fraction field K and residue field k .

1.1.1. Torsors and parahoric extension. Consider the special case of Theorem 1.3 where $\mathcal{X} = \mathcal{B}G$ is the classifying stack of a reductive group over a root stack of R . We say that a reductive group has tame Weyl group if its Weyl group has order coprime to the residue characteristics (Definition 4.1). Note that some condition on the group is necessary as illustrated by Example 4.3. Then Theorem 1.3 implies that every G -torsor over K is pulled back from some G -torsor over some root stack of $\text{Spec } R$. In fact the order of the root stack necessary is also uniformly bounded in terms of G .

Corollary 1.9. *Let $G \rightarrow \sqrt[n]{\text{Spec } R}$ be a group over a root stack, with G reductive and with tame Weyl group. Then there exists an integer n depending only on G such that the natural map*

$$\underline{\text{Hom}}(\sqrt[n]{\text{Spec } R}, \mathcal{B}G) \rightarrow \underline{\mathbf{H}}^1(K, G)$$

is essentially surjective where $\underline{\text{Hom}}$ denotes the groupoid of maps and $\underline{\mathbf{H}}^1$ denotes the groupoid of torsors.

Note that if R is strictly henselian, the groupoid of G -torsors on $\sqrt[n]{\text{Spec } R}$ is equivalent to that of G -torsors over $\mathcal{B}\mu_{n,k}$. Thus we get an essentially surjective functor

$$\underline{\text{Hom}}(\mathcal{B}\mu_{n,k}, \mathcal{B}G) = [\text{Hom}(\mu_{n,k}, G)/G] \rightarrow \underline{\mathbf{H}}^1(K, G)$$

where the action of G is by conjugation. By [MT15, Lemma 7.1], all such homomorphisms come from co-characters of G over the residue field k .

Remark 1.10. Compare Corollary 1.9 with [RS22, Lemma 3.5(d)].

Remark 1.11. The kernel of the natural map in Corollary 1.9 should be related to the $\frac{1}{n}\mathbb{Z}$ -points of the Bruhat-Tits building of G over K (see [GWZ25, Theorem 3.17]).

One of the main interests in Proposition 4.10 is that it can be interpreted as a parahoric extension theorem for G -bundles on K . Indeed by work of many authors [BS15, PR24, Dam24, BČE⁺25] [SSW24, Section 5], a reductive group over a root stack $\mathcal{R} \rightarrow \text{Spec } R$ is equivalent data to a parahoric model $\mathcal{G} \rightarrow \text{Spec } R$ of G_K and the category of parahoric \mathcal{G} -torsors over R is equivalent to the category of G -torsors on \mathcal{R} . Thus combining Proposition 2.10 with Proposition 1.5 and Corollary 1.6, we get the following parahoric extension result.

Proposition 1.12. *Suppose G is as in Theorem 1.3. Then any G -bundle over K extends to a parahoric bundle over R . The same holds if R and K are replaced by a global curve as in Proposition 1.5 and its function field respectively.*

Example 1.13. Suppose R has residue characteristic $\neq 2$ and let $C \subset \mathbb{P}_K^2$ be a smooth conic which corresponds to a map $\text{Spec } K \rightarrow \mathcal{B}PGL_2$. Let $\mathcal{C} \rightarrow \text{Spec } R$ be the closure of C in \mathbb{P}_R^2 . Then the central fiber is either a smooth conic, a doubled line, or a union of two lines. In the first case, the PGL_2 -torsor extends to R . In the second case, the normalization of the pullback of \mathcal{C} to $\sqrt[n]{\text{Spec } R}$ has good reduction and thus the torsor extends to the second root stack. In the third case, the total space has an A_n singularity at the node of the central fiber for some $n \geq 0$ (where by convention we an A_0 singularity is a regular point). If $n = 0$, blowing up the node and contracting the strict transform of the central fiber yields a model with good reduction so we are in the first case. If $n = 1$, blowing up the node and contracting the strict transform of the central fiber yields a model with a double line so we are in the second case. For $n \geq 2$, blowing up the

node and contracting the strict transform of the central fiber yields a new central fiber which is still a union of two lines but now has an A_{n-2} singularity. Repeating this operation puts us back in either the first or the second case depending on the parity of n . Thus for $\mathcal{B}\mathrm{PGL}_2$ in odd characteristic, any torsor over K extends up to a degree 2 root stack. \diamond

1.1.2. The Lang-Nishimura Theorem for Artin stacks. Following [BV24a], we give a generalization of Lang-Nishimura to Artin stacks with a good moduli space. We expect that this will have applications to questions on essential dimension, fields of definition and fields of moduli for Artin stacks as in e.g. [BV24b, BRV11].

Proposition 1.14 (Lang–Nishimura theorem). *Let S be a scheme and $\mathcal{X} \dashrightarrow \mathcal{Y}$ a rational map of algebraic stacks over S , with \mathcal{X} locally noetherian and integral and \mathcal{Y} admitting a good moduli space $\mathcal{Y} \rightarrow Y$ which is proper over S . Let k be a field, $s : \mathrm{Spec} k \rightarrow S$ a morphism. Assume that s lifts to a regular point $\mathrm{Spec} k \rightarrow \mathcal{X}$; then it also lifts to a morphism $\mathrm{Spec} k \rightarrow \mathcal{Y}$.*

The following is a straightforward corollary.

Corollary 1.15. *Suppose that a linearly reductive and special group G acts on an affine scheme $\mathrm{Spec} A$ and that the ring of invariants A^G is an R -algebra. Assume also that $\mathrm{Spec} A \otimes_R K$ has a K -point, and the composition $\mathrm{Spec} K \rightarrow \mathrm{Spec} A \rightarrow \mathrm{Spec} A^G$ extends to $\mathrm{Spec} R \rightarrow \mathrm{Spec} A^G$. Then $\mathrm{Spec} A \otimes_R k$ has a k -point.*

1.1.3. Gerbes and homogeneous spaces. Next we give some applications of Theorem 1.3 to gerbes and homogeneous spaces over DVRs which may be viewed as a version of Grothendieck-Serre for gerbes and non-principal homogeneous spaces.

Proposition 1.16. *Let $G \rightarrow \mathrm{Spec} R$ be a reductive group which is either special or has tame Weyl group. Then any class $\mathcal{G} \in H^2(R, G)$ which is trivial over $\mathrm{Spec} K$ becomes trivial after passing to a root stack. In particular, if $\mathcal{G}|_K = 0$ then $\mathcal{G}|_k = 0$.*

As a special case, we can consider a homogeneous space V for a group H which yields a gerbe $[V/H]$ for the stabilizer group.

Corollary 1.17. *Let $H \rightarrow \mathrm{Spec} R$ be a special affine group scheme. Let $V \rightarrow \mathrm{Spec} R$ be an affine homogeneous space for H and suppose that the geometric stabilizer groups are reductive groups with tame Weyl group. If $V(K) \neq \emptyset$ then $V(k) \neq \emptyset$.*

As unirationality for homogeneous spaces is equivalent to admitting a rational point, we obtain the following.

Corollary 1.18. *Unirationality specializes along DVRs for homogeneous spaces for special reductive groups with reductive geometric stabilizer with tame Weyl group.*

1.1.4. Applications to fibrations. Suppose \mathcal{X} is a stack with a proper good moduli space which parametrizes a class of projective varieties, for example $\overline{\mathcal{M}}_g$ or the KSBA moduli space of stable varieties, the K -moduli stack parametrizing K -semistable Fano varieties, or the moduli space of boundary polarized Calabi-Yau surface pairs [ABB⁺23, BL24]. Then the root stack valuative criterion yields certain nice integral models for fibrations and a method for classifying the singular fibers of fibrations whose general fiber is parametrized by the stack \mathcal{X} as follows.

Let $Y_\eta \rightarrow \mathrm{Spec} K$ be an object over of \mathcal{X} and let $\mathcal{Y} \rightarrow \mathcal{R}_n$ be the extension to a family over the n^{th} root stack given by Theorem 1.1. Then taking the coarse moduli space yields an integral model $Y \rightarrow \mathrm{Spec} R$ whose central fiber is a quotient of an object parametrized by \mathcal{X} by μ_n . This turns the question of classifying singular fibers to one of classifying μ_n actions on the objects of \mathcal{X} . See [AB19], [BPS24, Section 7] and [Rey25] for examples involving stable curves.

Corollary 1.19. *Let R be a DVR containing \mathbb{C} . Suppose $Y_\eta \rightarrow \mathrm{Spec} K$ is a K -semistable Fano variety. Then there exists a proper model $Y \rightarrow \mathrm{Spec} R$ with klt singularities whose central fiber is the quotient of a K -polystable Fano variety by μ_n . In particular, the central fiber is irreducible and of multiplicity n .*

Example 1.13 can be seen as a special case of Corollary 1.19 for the K -semistable Fano variety \mathbb{P}^1 .

Corollary 1.20. *Let R be a DVR containing \mathbb{C} . Suppose $(Y_\eta, D_\eta) \rightarrow \mathrm{Spec} K$ is a boundary polarized Calabi-Yau surface pair in the sense of [ABB⁺23, BL24]. Then there exists a proper model $(Y, D) \rightarrow \mathrm{Spec} R$ with semi-log canonical singularities whose central fiber is the quotient of a boundary polarized Calabi-Yau by a μ_n action. In particular, $(Y, D + (Y_0)_{\mathrm{red}})$ is semi-log canonical and Y_0 has multiplicity n .*

Moreover, by Corollary 1.6, Corollaries 1.19 and 1.20 have analogues for fibrations over global curves.

1.2. Conventions. By reductive group scheme $G \rightarrow S$ we mean, as in [Con14, Definition 3.1.1], a smooth S -affine group scheme, such that the geometric fibers are connected reductive groups. When we don't specify, the cohomology will be the fppf cohomology. We will pass between an algebraic stack over a base scheme B to the corresponding category fibered in groupoids over $Sch(B)$ without mentioning it specifically; we expect that this will not cause any confusion. Whenever we mention the site of an algebraic stack \mathcal{X} , it will be the flat-fppf site: its objects are flat morphisms $B \rightarrow \mathcal{X}$ from a scheme, and a covering of an object is an fppf-covering. All algebraic stacks will be noetherian, unless mentioned otherwise. All algebraic stacks in the body of the paper will have affine diagonal. The n^{th} root stack of a DVR R , denoted $\sqrt[n]{\text{Spec } R}$ or \mathcal{R}_n , is the algebraic stack $[\text{Spec } R[t^{1/n}]/\mu_n]$ where t is a uniformizer of R .

1.3. Acknowledgements. We thank Jarod Alper, Harold Blum, Patrick Brosnan, Elden Elmanto, Michael Groechenig, Tom Haines, Sam Molcho, Swarnava Mukhopadhyay, Danny Ofek, Zinovy Reichstein, Jason Starr for helpful conversations. We thank Andres Fernandez Herrero for helpful conversations and for giving us detailed feedback on a previous version of this manuscript. We especially thank Sid Mathur for many helpful conversations about gerbes.

2. GROUPS, TORSORS AND GERBES ON ALGEBRAIC STACKS

In this section we develop some preliminary results on reductive groups and their torsors and gerbes over an algebraic stack. We also refer the reader to [BIMS25] where this material will be developed in more depth.

2.1. Reductive groups over a stack. We begin with the following

Definition 2.1. Let \mathcal{X} be an algebraic stack. A group G over \mathcal{X} is a sheaf on $\mathcal{X}_{\text{fppf}}$ which is a sheaf in groups, and such that there is a fppf-cover $U \rightarrow \mathcal{X}$ such that the sheaf $G|_U$ is represented by a scheme over U . We say that G is a smooth (resp. separated, of finite type, fppf) group over \mathcal{X} , if there is an fppf-cover as above such that $G|_U$ is represented by a smooth (resp. separated, of finite type, fppf) group scheme over U .

Definition 2.2. Let $G \rightarrow \mathcal{X}$ be a smooth separated group over an algebraic stack \mathcal{X} . We say that G is reductive (resp. linearly reductive, a torus) if for every fppf-cover $B \rightarrow \mathcal{X}$ from a scheme B , the pull-back $G \times_{\mathcal{X}} B \rightarrow B$ is a reductive (resp. linearly reductive, a torus) B -group.

Lemma 2.3. *Let $G \rightarrow \mathcal{X}$ a reductive group over a tame algebraic stack as above, with connected fibers. Then there is an algebraic stack \mathcal{T} , with a smooth affine and representable morphism $\mathcal{T} \rightarrow \mathcal{X}$ such that for every morphism $f : B \rightarrow \mathcal{X}$, there is an equivalence*

$$\text{Hom}_{\mathcal{X}}(B, \mathcal{T}) \longleftrightarrow \{\text{maximal tori of } G \times_{\mathcal{X}} B\}.$$

Proof. Note that $B \mapsto \{\text{maximal tori of } G \times_{\mathcal{X}} B\}$ is an fppf sheaf. Indeed, the restriction of this functor to B_{fppf} is represented by the scheme $\text{Tor}_{G_B/B} \rightarrow B$, as proven in [Con14, Theorem 3.2.6]. Thus, \mathcal{T} is a sheaf over \mathcal{X} . Then to prove representability, it is enough to show $\mathcal{T} \rightarrow \mathcal{X}$ is affine. For this, we may look fppf locally, where we can assume \mathcal{X} is a scheme; we then apply [Mil13, XII, 5.4]. \square

Lemma 2.4. *Let \mathcal{X} be an algebraic stack, and $T \rightarrow \mathcal{X}$ a torus over \mathcal{X} which fppf-locally is isomorphic to \mathbb{G}_m^n . Then the sheaf $\underline{\text{Isom}}_{\text{grp}}(\mathbb{G}_m^n \times \mathcal{X}, T)$ over $\mathcal{X}_{\text{fppf}}$, that sends $U \rightarrow \text{Isom}_{\text{grp}}(\mathbb{G}_m^n(U), T(U))$, is represented by an algebraic stack, which is étale over \mathcal{X} but not noetherian for $n > 1$.*

Proof. We see that $\underline{\text{Isom}}_{\text{grp}}(\mathbb{G}_m^n \times \mathcal{X}, T)$ is a $\text{GL}_{n, \mathcal{X}}(\mathbb{Z}) = \text{Aut}(\mathbb{G}_{m, \mathcal{X}}^n)$ -torsor. Since $\text{GL}_{n, \mathcal{X}}(\mathbb{Z})$ is étale over \mathcal{X} , it follows that $\underline{\text{Isom}}_{\text{grp}}(\mathbb{G}_m^n \times \mathcal{X}, T)$ is étale over \mathcal{X} . Note it is not noetherian as $\text{GL}_{n, \mathcal{X}}(\mathbb{Z})$ is not noetherian. \square

The following lemma is a generalization of [Ser10, Theorem 3'']; see also [GCP24, Lemma 2.7].

Lemma 2.5. *Let G be a reductive group over a field k and suppose μ_r acts on G . Then there exists a μ_r -invariant maximal torus of G defined over k .*

Proof. If G is a torus there is nothing to prove, so we will assume G is not a torus. Suppose k has characteristic exponent $p \geq 1$ and write $r = p^a m$ where m is coprime to p . Following the proof of [Ser10, Theorem 3''], we induct on m and the dimension of G . When $m = 0$, $\mu_r = \mu_{p^a}$ is a connected group scheme. Let $\varphi : \mu_r \rightarrow \text{Aut}(G)$ be the morphism inducing the action. Since $\text{Out}(G)$ is étale over k [Con14, Theorem 7.1.9(1)], the composition $\mu_r \rightarrow \text{Out}(G)$ is trivial so φ must factor through the adjoint group $G^{\text{ad}} = G/Z(G)$. By [MT15, Lemma 7.1], the morphism $\varphi : \mu_r \rightarrow G^{\text{ad}}$ factors through a torus which we may take to be maximal by Grothendieck's theorem on maximal tori (see [Con14, Remark A.1.2]). Lifting this maximal torus of G^{ad} to G , we obtain a maximal torus of G which is μ_r -invariant as required. If $m > 0$, then $\mu_r \cong \mu_{p^a} \times \mu_m$ where μ_m is étale of order coprime to the characteristic. Then the invariant subgroup of the μ_m action G^{μ_m} satisfies the conclusions of [Ser10, Proposition 3]. Indeed as in *loc. cit.*, the conclusions can be checked after passing to \bar{k} in which case $\mu_m \cong \mathbb{Z}/m\mathbb{Z}$ where *loc. cit.* applies directly. In particular, $G_1 = (G^{\mu_m})^0$ is reductive and it is positive dimensional if G is not a torus. Now we proceed exactly as in the end of the proof of [Ser10, Theorem 3''] with A replaced by μ_r and $A' = \langle s \rangle$ by μ_m , we report the salient steps below. We consider the $\mu_r/\mu_m = \mu_{p^a}$ action on G_1 and take a μ_{p^a} -invariant torus in G_1 , which exists by the base case $m = 0$ above, and we take G_2 its centralizer. Observe that μ_r acts on G_2 . As $\dim(G_2) < \dim(G)$, by induction G_2 admits a μ_r -maximal torus. This will have the same dimension as a maximal torus for G ; thus we have found a maximal torus of G (contained in G_2) which is fixed by μ_r . \square

Remark 2.6. The proof shows more generally that the conclusion of [Ser10, Theorem 3''] holds whenever A is a linearly reductive finite group scheme with a composition series $1 = A_0 \subset A_1 \dots \subset A_l = A$ such that A_i is normal in A and A_{i+1}/A_i is a k -form of either μ_r or $\mathbb{Z}/r\mathbb{Z}$ for some r .

Proposition 2.7. *Let R be a henselian DVR, and let \mathcal{R} be the r^{th} root stack of $\text{Spec } R$ at the closed point. Let $G \rightarrow \mathcal{R}$ be a reductive group with connected fibers. Then there is a torus $T \rightarrow \mathcal{R}$ with a morphism $i : T \subseteq G$ such that, for every geometric point $\text{Spec } l \rightarrow \mathcal{R}$, the fiber $T_l \rightarrow G_l$ is a maximal torus.*

Proof. For any map of algebraic stacks $f : \mathcal{X}' \rightarrow \mathcal{X}$ let $\Gamma(\mathcal{X}'/\mathcal{X})$ denote the groupoid of sections of f . Let $\mathcal{T} \rightarrow \mathcal{R}$ be the stack of maximal tori of G which is smooth and affine over \mathcal{R} by Lemma 2.3. We wish to show that $\Gamma(\mathcal{T}/\mathcal{R})$ is nonempty. Let $\mathcal{R}_0 = B\mu_r$ be the residual gerbe of the closed point of $\text{Spec } R$ and let $\mathcal{T}_0 = \mathcal{T} \times_{\mathcal{R}} \mathcal{R}_0$, $G_0 = G \times_{\mathcal{R}} \mathcal{R}_0$. The data of G_0 is equivalent to the data of a reductive group G_k over k , the residue field of R , together with a μ_r -action on it. By Lemma 2.5, there exists a μ_r -invariant maximal torus of G_k , i.e. the groupoid of sections $\Gamma(\mathcal{T}_0/\mathcal{R}_0)$ is nonempty. Since R is henselian, the pair $(\mathcal{R}, \mathcal{R}_0)$ is henselian by [AHR25, Theorem 3.6], and \mathcal{R} has the resolution property as it is a global quotient of an affine scheme by \mathbb{G}_m . Thus by [AHR25, Proposition 7.9], $\Gamma(\mathcal{T}/\mathcal{R}) \rightarrow \Gamma(\mathcal{T}_0/\mathcal{R}_0)$ is essentially surjective, and in particular nonempty. \square

2.2. Gerbes and torsors over stacks. We recall some background on torsors, gerbes and bands. Let \mathcal{X} be an algebraic stack and G/\mathcal{X} be a group stack as in the previous section.

Definition 2.8. A G -torsor over \mathcal{X} is an fppf sheaf $P \rightarrow \mathcal{X}$ with an action $G \times_{\mathcal{X}} P \rightarrow P$ for which there exists an fppf cover $U \rightarrow \mathcal{X}$ such that $P_U \cong G_U$ with the left action of G_U . A gerbe over \mathcal{X} is an fppf stack $\mathcal{Y} \rightarrow \mathcal{X}$ for which there exists an fppf cover $U \rightarrow \mathcal{X}$ and a group $G_U \rightarrow U$ such that $\mathcal{Y}_U \cong BG_U$ where BG_U is the classifying stack of $G_U \rightarrow U$.

We refer to [Gir65, Chapter IV] for background on bands which we recall here; see also [EHKV01, Section 3.1]. Let $\mathcal{G} \rightarrow \mathcal{X}$ be a gerbe which is trivialized after an fppf cover $U \rightarrow \mathcal{X}$ and suppose that the group G_U is pulled back from a group stack $G \rightarrow \mathcal{X}$. Over $U \times_{\mathcal{X}} U$ we have descent data for BG_U which is a section of the stack $\text{Isom}_U(BG_U, BG_U) = [\text{Aut}(G_U)/G_U]$ where G_U acts by conjugation. Passing to π_0 yields a section of $\text{Out}(G_U)$ over $U \times_{\mathcal{X}} U$ satisfying the cocycle condition, i.e. an element of $H^1(\mathcal{X}, \text{Out}(G))$. Note that there is a natural map $H^1(\mathcal{X}, \text{Aut}(G)) \rightarrow H^1(\mathcal{X}, \text{Out}(G))$ and that elements of $H^1(\mathcal{X}, \text{Aut}(G))$ classify \mathcal{X} -forms of G .

Definition 2.9. Let $\mathcal{G} \rightarrow \mathcal{X}$ be a gerbe as above. The band of \mathcal{G} is the element $\text{Band}(\mathcal{G}) \in H^1(\mathcal{X}, \text{Out}(G))$ described above. Similarly, if $G' \rightarrow \mathcal{X}$ is a form of G , we define the band of G' as the image $\text{Band}(G') \in H^1(\mathcal{X}, \text{Out}(G))$ of the class of G' under the natural map $H^1(\mathcal{X}, \text{Aut}(G)) \rightarrow H^1(\mathcal{X}, \text{Out}(G))$. We say \mathcal{G} is banded by G' if $\text{Band}(\mathcal{G}) = \text{Band}(G')$ and call \mathcal{G} a G' -gerbe.

When G is abelian, G -torsors and G -gerbes are classified by fppf cohomology $H^1(\mathcal{X}, G)$ and $H^2(\mathcal{X}, G)$ respectively (e.g. [Ols16, Theorems 12.1.5 & 12.2.8]). When G is non-abelian, we follow [Gir65] and define $H^1(\mathcal{X}, G)$ and $H^2(\mathcal{X}, G)$ respectively as the set of isomorphism classes of G -torsors and G -gerbes on X . The following is a generalization of [Dou75, Ch. V, sect. 3.2, p. 75] to stacks.

Proposition 2.10. *Let $\mathcal{G} \rightarrow \mathcal{X}$ be a gerbe over an algebraic stack \mathcal{X} . Suppose that there exists an fppf cover $U \rightarrow \mathcal{X}$ such that $\mathcal{G}|_U \cong \mathcal{B}G_U$ where $G_U \rightarrow U$ is reductive and U is connected. Then there exists a reductive group $G \rightarrow \mathcal{X}$ and a form $G' \rightarrow \mathcal{X}$ of G such that $\text{Band}(G') = \text{Band}(\mathcal{G})$.*

Proof. Since U is connected, the type of $G_U \rightarrow U$ is constant [DGA⁺11, XXII Section 2]. Let $G_0 \rightarrow \text{Spec } \mathbb{Z}$ be the Chevalley group of this type, that is, the unique split reductive group with the given root data [Con14, Theorem 6.1.17]. After possibly refining U , we may assume that $G_U \cong G_0 \times_{\mathbb{Z}} U$. Let $G := G_0 \times_{\mathbb{Z}} \mathcal{X}$. Then $\text{Band}(\mathcal{G}) \in H^1(\mathcal{X}, \text{Out}(G))$. On the other hand, for the Chevalley model, $\text{Aut}(G_0) \rightarrow \text{Out}(G_0)$ splits [Con14, Theorem 7.1.9 (3)]. Fix such a splitting. This induces a splitting $\text{Aut}(G) \rightarrow \text{Out}(G)$ over \mathcal{X} and thus a lift of $\text{Band}(\mathcal{G})$ to an element of $H^1(\mathcal{X}, \text{Aut}(G))$. Let G' be the corresponding \mathcal{X} -form of G . This is our desired group scheme. \square

3. PRELIMINARY REDUCTIONS

In this section we make some preliminary reductions that show it suffices to prove Theorem 1.1 in the case where the stack is a gerbe whose geometric automorphism groups are reductive. Throughout this section, we let R be a DVR with fraction field K and residue field k and we assume that \mathcal{X} is an algebraic stack with good moduli space $\mathcal{X} \rightarrow \text{Spec } R$.

3.1. Passing to the polystable locus. We first show that it suffices to prove Theorem 1.1 in the case where the K -point of \mathcal{X} is polystable (see e.g. [BES24, Definition 3.2 and Lemma 3.3]). We begin with the following.

Lemma 3.1. *Let $V = \text{Spec } A \rightarrow \text{Spec } K$ with a GL_n action such that $[V/\text{GL}_n] \rightarrow \text{Spec } K$ is a good moduli space. Let $x \in V(K)$ be any K -point. Then there is a one parameter subgroup $\lambda : \mathbb{G}_m \rightarrow \text{GL}_n$ defined over K such that $\lim_{t \rightarrow 0} \lambda(t)x$ exists and belongs to the unique closed orbit of V .*

Proof. For perfect fields this follows from [Kem78], especially [Kem78, §4]. In characteristic p instead we proceed as follows. By [BHMR17, Theorem 1.3], the cocharacter closure of $\text{GL}_n(K)x$ contains a unique cocharacter closed $\text{GL}_n(K)$ orbit and there is a $\lambda : \mathbb{G}_m \rightarrow \text{GL}_n$ defined over K such that $\lim_{t \rightarrow 0} \lambda(t)x$ exists and belongs to this cocharacter closed orbit. We wish to show that this limit in fact belongs to the closed orbit. Thus it suffices to check that every cocharacter closed orbit of the $\text{GL}_n(K)$ action on V is contained in the unique closed orbit of V . We refer the reader to [BHMR17] for the definition of cocharacter closed and cocharacter closure. Toward this end, let $z \in V$ be a point in a cocharacter-closed orbit. After passing to the separable closure K^s of K , we can write $[V/\text{GL}_n] \times_K K^s \cong [W/G]$ where W is affine and $G \subset \text{GL}_n$ is the stabilizer of the closed orbit. By assumption, G is linearly reductive so its connected component is a torus $T \cong \mathbb{G}_{m,K^s}^n$. By [BHMR17, Theorem 1.5(ii)], z remains cocharacter closed over K^s and under the isomorphism with $[W/G]$ corresponds to a cocharacter closed orbit of $G(K)$. Again by [Kem78], after passing further to the algebraic closure \overline{K}/K^{sep} , there is a 1-parameter subgroup $\lambda_{\overline{K}} : (\mathbb{G}_m)_{\overline{K}} \rightarrow (\mathbb{G}_m^n)_{\overline{K}}$ such that $\lim_{t \rightarrow 0} \lambda_{\overline{K}}(t)z$ exists and is contained in the closed orbit. But λ is of the form $t \mapsto (t^{a_1}, \dots, t^{a_n})$ with $a_i \in \mathbb{Z}$, so it is the pullback of a character $\lambda_{K^s} : (\mathbb{G}_m)_{K^s} \rightarrow (\mathbb{G}_m^n)_{K^s}$. Thus $\lim_{t \rightarrow 0} \lambda_{K^s}(t)z$ exists and is contained in the closed orbit. Since z is cocharacter closed, this limit must also be contained in $G(K)z$ so z is contained in the closed orbit of $V \times_K K^s$. As belonging to the closed orbit can be checked after a separable field extensions, then z is also in the closed orbit of V , as desired. \square

Proposition 3.2. *Let R be a DVR with generic point $\eta = \text{Spec } K$. Consider an algebraic stack \mathcal{X} with a good moduli space $\xi : \mathcal{X} \rightarrow \text{Spec } R$. Assume that*

- (1) \mathcal{X} admits a K -point $\phi : \eta \rightarrow \mathcal{X}$, and
- (2) every $\psi : \eta \rightarrow \mathcal{X}$ sending η to the polystable point of \mathcal{X}_{η} extends to a section over some root stack $\mathcal{R} = \sqrt[n]{\text{Spec } R}$ of $\text{Spec } R$.

Then ϕ extends to a section over some root stack \mathcal{R} of $\text{Spec } R$.

Proof. We first prove that there is a map $\Theta_K := [\mathbb{A}_K^1/\mathbb{G}_m] \rightarrow \mathcal{X}$ sending 1 to $\phi(\eta)$ and 0 to the unique polystable point of \mathcal{X}_K over η . Indeed, from [AHR25, Theorem 6.1], we can write $\mathcal{X}_K = [V/\mathrm{GL}_n]$ with V affine and $\mathcal{X}_K \rightarrow \eta$ a good moduli space. By Lemma 3.1, there is a 1-parameter subgroup defined over K such that $\lambda : \mathbb{G}_m \rightarrow \mathrm{GL}_n$ such that $\lim_{t \rightarrow 0} \lambda(t)\phi(\eta)$ is in the unique closed orbit of the GL_n action on V . Taking quotients of the induced map $\mathbb{A}_K^1 \rightarrow \mathcal{X}_K$ yields the required map $\Theta_K \rightarrow \mathcal{X}$.

By assumption, up to replacing $\mathrm{Spec} R$ with a root stack \mathcal{R} , we can extend $\mathrm{Spec} K \xrightarrow{\iota} \mathbb{A}_K^1 \rightarrow \mathcal{X}$ to a morphism $\mathcal{R} \rightarrow \mathcal{X}$, where ι is the inclusion of the origin in \mathbb{A}_K^1 . Consider then the following diagram

$$\begin{array}{ccc} \mathrm{Spec} K & \longrightarrow & \mathrm{Spec} R \\ \downarrow \iota & & \downarrow \beta \\ \mathbb{A}_K^1 & \xrightarrow{\alpha} & \mathrm{Spec} A. \end{array}$$

where A is the fiber product of $K[t] \times_K R$. More explicitly,

$$A = \{p(t) \in K[t] : p(0) \in R\} = R \left[t, \frac{t}{\pi}, \frac{t}{\pi^2}, \dots \right]$$

where π is a uniformizer in R . As $\mathrm{Spec} K \rightarrow \mathbb{A}_K^1$ is a closed embedding and $\mathrm{Spec} K \rightarrow \mathrm{Spec} R$ is affine, $\mathrm{Spec} A$ is a pushout of the previous diagram in algebraic stacks by [AHHLR24, Theorem 4.2]. Consider the n -th root stack of $\mathrm{Spec} A$ at the Cartier divisor $\pi = 0$. This leads to the following cartesian diagram:

$$\begin{array}{ccc} \mathrm{Spec} K & \longrightarrow & \mathcal{R} \\ \downarrow \iota & & \downarrow b \\ \mathbb{A}_K^1 & \xrightarrow{a} & \sqrt[n]{\mathrm{Spec} A}. \end{array}$$

We argue now that this diagram is also a pushout in algebraic stacks. By [AHHLR24, Theorem 4.2] it suffices to check that $\mathcal{O}_{\sqrt[n]{\mathrm{Spec} A}} \rightarrow a_* \mathcal{O}_{\mathbb{A}_K^1} \times_{(a \circ \iota)_* \mathcal{O}_{\mathrm{Spec} K}} b_* \mathcal{O}_{\mathcal{R}}$ is an isomorphism. This can be checked flat-locally on $\sqrt[n]{\mathrm{Spec} A}$, so consider the cover $\mathrm{Spec} A[r]/(r^n - \pi) \rightarrow \sqrt[n]{\mathrm{Spec} A}$. This cover pulls back to the diagram of schemes on the left, leading to the diagram of rings on the right

$$\begin{array}{ccccc} \mathrm{Spec} K[r]/(r^n - \pi) & \longrightarrow & \mathrm{Spec} R[r]/(r^n - \pi) & & K[r]/(r^n - \pi) \longleftarrow R[r]/(r^n - \pi) \\ \downarrow & & \downarrow & & \uparrow \scriptstyle t=0 \\ \mathrm{Spec} K[t, r]/(r^n - \pi) & \longrightarrow & \mathrm{Spec} A[r]/(r^n - \pi) & & K[t, r]/(r^n - \pi) \longleftarrow A[r]/(r^n - \pi) \end{array}$$

It suffices to note that the right-most diagram is a pull-back in rings.

By the universal property of pushouts, we have a map $\sqrt[n]{\mathrm{Spec} A} \rightarrow \mathcal{X}$. Since

$$A = \mathrm{colim}_l R \left[\frac{t}{\pi^m} \mid m \leq l \right] = \mathrm{colim}_l R \left[\frac{t}{\pi^l} \right]$$

and \mathcal{X} is of finite type, by the functorial characterization of locally finitely presented morphisms, the map $\sqrt[n]{\mathrm{Spec} A} \rightarrow \mathcal{X}$ factors via $\sqrt[n]{\mathrm{Spec} R[\frac{t}{\pi^k}]} = \sqrt[n]{\mathrm{Spec} R[s, t]/(s\pi^k - t)} \rightarrow \mathcal{X}$ for some $k \gg 0$. Consider the morphism $i : \mathrm{Spec} R \rightarrow \mathrm{Spec} R[s, t]/(s\pi^k - t)$ induced by the R -algebra homomorphism $j : R[s, t]/(s\pi^k - t) \rightarrow R$ that sends $s \mapsto \pi$ and $t \mapsto \pi^{k+1}$. We have that:

- (1) the image $i(\eta)$ of the generic point of $\mathrm{Spec} R$ is contained in $D(t) \cap D(\pi) = \mathbb{A}_K^1 \setminus \{0\}$ as $j(t\pi) \neq 0$
- (2) the image $i(x)$ of the closed point of $\mathrm{Spec} R$ is the prime ideal (π, s, t) , which agrees with the image of the closed point of $\mathrm{Spec} R$ via β .

Then it induces a morphism $\sqrt[n]{\mathrm{Spec} R} \rightarrow \sqrt[n]{\mathrm{Spec} R[s, t]/(s\pi^k - t)}$, which if postcomposed with the morphism $\sqrt[n]{\mathrm{Spec} R[s, t]/(s\pi^k - t)} \rightarrow \mathcal{X}$ yields a morphism $\sqrt[n]{\mathrm{Spec} R} \rightarrow \mathcal{X}$ whose restriction to the generic point $\eta \rightarrow \mathcal{X}$ is isomorphic to $\phi : \eta \rightarrow \mathcal{X}$ by (1), and which sends the special point to a point over the closed point of $\mathrm{Spec} R$ by (2). \square

3.2. Reduction to the case of a gerbe. The next result shows that it suffices to prove Theorem 1.1 for gerbes over root stacks.

Theorem 3.3. *Let R be a DVR, with generic point $\eta = \text{Spec } K$. Consider a stack \mathcal{X} with good moduli space $\pi : \mathcal{X} \rightarrow \text{Spec } R$, and suppose $\phi : \eta \rightarrow \mathcal{X}^{ps} \subset \mathcal{X}$ is a polystable K -point. Then there exists a root stack $\mathcal{R}_n = \sqrt[n]{\text{Spec } R}$ and a diagram as follows*

$$\begin{array}{ccccc} & & \mathcal{G} & \xrightarrow{p} & \mathcal{X} \\ & \nearrow \phi' & \downarrow \pi' & & \downarrow \pi \\ \eta & \longrightarrow & \mathcal{R}_n & \xrightarrow{q} & \text{Spec } R \end{array}$$

where q is the coarse moduli space map, ϕ and $p \circ \phi'$ are isomorphic and π' is a gerbe banded by a smooth, connected, linearly reductive group scheme $G \rightarrow \mathcal{R}_n$. Moreover, the closed point of \mathcal{G} maps to a polystable point in \mathcal{X} .

The argument follows closely many of the ideas in [BES24].

Proof. Let $x \in \mathcal{X}_K$ be the polystable point of \mathcal{X}_K and let \mathcal{X}'' be the closure of $\{x\} \subset \mathcal{X}$. Since $\mathcal{X} \rightarrow \text{Spec } R$ is universally closed, and since closed substacks of a stack with a separated good moduli space still admit a good moduli space [Alp13, Lemma 4.14], the stack \mathcal{X}'' admits a good moduli space which is closed in $\text{Spec } R$ and which contains η . In other terms, the composition $\mathcal{X}'' \rightarrow \mathcal{X} \rightarrow \text{Spec } R$ is a good moduli space. Moreover, the point η is a stable point of $\mathcal{X}'' \rightarrow \text{Spec } R$ so $\mathcal{X}'' \rightarrow \text{Spec } R$ is a stable good moduli space, i.e. the stable locus is dense. Thus we may apply canonical reduction of stabilizers [ER21] which yields a modification $\mathcal{X}' \rightarrow \mathcal{X}''$ such that

- $\mathcal{X}' \rightarrow \mathcal{X}''$ is an isomorphism over η ,
- the dimension of the stabilizers of the geometric points of \mathcal{X}' is constant, and
- \mathcal{X}' admits a good moduli space which is obtained by a sequence of blow-ups of $\text{Spec } R$.

Since R is a DVR, the sequence of blowups of $\text{Spec } R$ must be the identity and the composition $(\mathcal{X}')_{red} \rightarrow \mathcal{X}' \rightarrow \text{Spec } R$ is a good moduli space by [Alp13, Theorem 4.16(viii)] so without loss of generality we may assume \mathcal{X}' is reduced. By [ER21, Proposition B.2] and its proof (especially [ER21, Proposition B.6], see also [BES24, Proposition 2.3]) we can factor $\mathcal{X}' \rightarrow \text{Spec } R$ as $\mathcal{X}' \rightarrow \mathcal{Y} \rightarrow \text{Spec } R$ where the first map is a gerbe whose geometric automorphism groups are the smooth connected components of the identity of the geometric automorphism groups of \mathcal{X}' and $\mathcal{Y} \rightarrow \text{Spec } R$ is a tame coarse moduli space. By [Alp13, Proposition 3.13], $\mathcal{X}' \rightarrow \mathcal{Y}$ is also a good moduli space so the geometric automorphism groups are also linearly reductive.

Note that by construction the K -point $\phi : \eta \rightarrow \mathcal{X}$ factors through $\mathcal{X}' \rightarrow \mathcal{X}$. Now [BV24a, Theorem 3.1] applies to the map $\mathcal{Y} \rightarrow \text{Spec } R$ so $\eta \rightarrow \mathcal{X}' \rightarrow \mathcal{Y}$ extends to a section of $\mathcal{Y} \rightarrow \text{Spec } R$ up to replacing $\text{Spec } R$ with a root stack \mathcal{R}_n . Then $\mathcal{G} := \mathcal{X}' \times_{\mathcal{Y}} \mathcal{R}_n \rightarrow \mathcal{R}_n$ is a good moduli gerbe with smooth, connected, linearly reductive geometric automorphism groups. It follows from Proposition 2.10 that any such gerbe is banded by a smooth, connected, linearly reductive group scheme over \mathcal{R}_n (i.e. a form of G over \mathcal{R}_n). Moreover, by construction ϕ factors through $\phi' : \eta \rightarrow \mathcal{G}$ as required.

As for the moreover part, observe that at each saturated blow-up in the algorithm of Edidin and Rydh [ER21], we are taking an open substack (namely, using their notation, a *saturated blow-up*) of the blow-up of the polystable point over the closed point of \mathcal{R} . Hence it suffices to show that the saturated blow-up intersects the exceptional divisor of the blow-up, and each blow-up in the Edidin-Rydh algorithm is centered at a point on the exceptional divisor of the previous blow-up. Both these statements follow from [ER21, Proposition 3.7 (3)]: the first one follows from [ER21, Proposition 3.7 (3)] directly, the second one follows as at each step we are blowing up the polystable point, which is the unique closed point (over the closed point of $\text{Spec } R$). As the exceptional divisor is closed, it will contain the unique closed polystable point. Then any point on the last exceptional divisor we extract will map to the polystable point of \mathcal{X} . \square

4. GERBE CASE

The goal of this section is to prove the root stack valuative criterion for gerbes banded by a reductive group. When the residue characteristic is finite, we need a tameness assumption on the reductive group.

Definition 4.1. Let $G \rightarrow S$ be a reductive group over a stack S . We say G has tame Weyl group if for all geometric points $x : \operatorname{Spec} k \rightarrow S$, the order of the Weyl group of G_x is coprime to the characteristic of k .

Theorem 4.2. Let R be a henselian DVR with fraction field K and let \mathcal{R} be a root stack of $\operatorname{Spec} R$. Let $\mathcal{G} \rightarrow \mathcal{R}$ be a gerbe banded by a reductive group $G \rightarrow \mathcal{R}$ which is either special or has tame Weyl group. Suppose that \mathcal{G} admits a K -point $\phi : \operatorname{Spec} K \rightarrow \mathcal{G}$. Then ϕ extends to a section $\mathcal{R} \rightarrow \mathcal{G}$ up to replacing \mathcal{R} with a further root stack.

The following example illustrates that some assumption on G is necessary.

Example 4.3. Let R be a mixed characteristic $(0, 2)$ henselian DVR with imperfect residue field k , for example the henselization of $\mathbb{Z}[t]_{(2)}$, and let $t \in R$ be an element which is not a square in k . Consider the conic $C = \{x^2 + y^2 = tz^2\} \subset \mathbb{P}_K^2$. This can be interpreted as a map $\operatorname{Spec} K \rightarrow \mathcal{B}\operatorname{PGL}_2$. We claim that this map does not extend to any root stack of $\operatorname{Spec} R$. Suppose toward contradiction that it did extend over a root stack \mathcal{R}_n . Then the pullback C' of C to $R' = R[2^{1/n}]$ has good reduction, where R' has the same residue field k . Let $C' \rightarrow \operatorname{Spec} R'$ be a smooth proper model. Then by smoothness C' admits a k'' -point where k''/k is a separable extension. By the henselian assumption, this extends to an R'' -point where R''/R' the unique finite étale extension corresponding to k''/k . In particular, $C'' = C \times_R R''$ has a K'' -rational point. This implies its discriminant must be a square. On the other hand, the discriminant of $x^2 + y^2 = tz^2$ is $8t$, so after possibly adjoining a further root of 2, this implies that t is a square in K'' . Thus its reduction is a square in k'' which contradicts the fact that k''/k is separable so C cannot have good reduction over any root stack. \diamond

The proof will proceed as follows. First we will prove that, up to replacing \mathcal{R} with a further root stack, the gerbe \mathcal{G} has to be trivial. This will be achieved by replacing \mathcal{G} with a gerbe over \mathcal{R} banded by the center of G , which is an extension of a torus by a tame finite group. Using [BV24a] we will further reduce the problem to the case in which G is a torus, which we handle explicitly. Once we know that \mathcal{G} is trivial, the problem will become to lift a class $c \in H^1(\eta, G|_\eta)$. If G is special, this is automatic. In the case where G has tame Weyl group, we will reduce again to the case of a torus by showing that G admits a maximal torus T , lifting c to a class in $H^1(\eta, N|_\eta)$ where N is the normalizer of T in G , and reducing again to the case of a torus using that N is an extension of a torus by a finite tame group (so using [BV24a]).

4.1. Torus case. In this section we include a few preparatory results that will be needed later. Whenever we take a cohomology group on an algebraic stack, we will use the flat-fppf cohomology.

Lemma 4.4. Let \mathcal{X} be a tame regular algebraic stack with a dense open substack that is a scheme. Then $H^2(\mathcal{X}, \mathbb{G}_m)$ is torsion and if \mathcal{X} admits a coarse moduli space $\pi : \mathcal{X} \rightarrow X$, the cokernel of $\pi^* : H^1(X, \mathbb{G}_m) \rightarrow H^1(\mathcal{X}, \mathbb{G}_m)$ is torsion.

Proof. We first prove that $H^2(\mathcal{X}, \mathbb{G}_m)$ is torsion. It follows from the same argument as in [Lie08, Proposition 3.1.3.3] that the morphism $H^2(\mathcal{X}, \mathbb{G}_m) \rightarrow H^2(U, \mathbb{G}_m)$ is injective, where U is a dense open substack which is a scheme (see also [Ach24, Prop. 2.23]). More specifically, consider a class $c \in H^2(\mathcal{X}, \mathbb{G}_m)$ which restrict to the trivial class generically. This corresponds to a gerbe $\mathcal{G} \rightarrow \mathcal{X}$ which generically is trivial. So $\mathcal{G}|_U$ has a 1-twisted line bundle \mathcal{L}_η , and its push-forward via $\mathcal{G}|_U \rightarrow \mathcal{G}$ will be a 1-twisted line bundle. As $H^2(U, \mathbb{G}_m)$ is torsion from [Lie08, Corollary 3.1.3.4], also $H^2(\mathcal{X}, \mathbb{G}_m)$ is torsion. To check that $H^1(\mathcal{X}, \mathbb{G}_m)$ is torsion, it suffices to use [AH11, Lemma 2.3.7] to argue that for every line bundle \mathcal{L} on \mathcal{X} , a power $\mathcal{L}^{\otimes N}$ descends to X . \square

Proposition 4.5. Let R be a henselian DVR, let \mathcal{R} be a root stack of $\operatorname{Spec} R$ and let $T \rightarrow \mathcal{R}$ a smooth group stack with geometric fibers isomorphic to \mathbb{G}_m^n for a fixed n . Then there is a finite étale morphism $\pi : \mathcal{R}' \rightarrow \mathcal{R}$ where \mathcal{R}' is also a root stack of a henselian DVR such that $\pi^{-1}T \cong \mathbb{G}_m^n \times \mathcal{R}'$.

In other terms, we can split the torus after a finite étale morphism of the root stack \mathcal{R} . This is well-known if \mathcal{R} is a scheme.

Proof. Let p be the residue characteristic of the closed point of $\operatorname{Spec} R$, let μ_d the stabilizer of the closed point of \mathcal{R} and π the uniformizer of R . If $p > 0$ let e and d' be such that $d = p^e d'$ and p does not divide d' , so $\mu_d \cong \mu_{p^e} \times \mu_{d'}$; if $p = 0$ let $d = d'$. Observe now that $\operatorname{Spec} R[t]/(t^d - \pi) \cong \operatorname{Spec} R[t_1, t_2]/(t_1^{p^e} - \pi, t_2^{d'} - t_1)$

and $\mathcal{R} \cong [\mathrm{Spec} R[t_1, t_2]/(t_1^{p^e} - \pi, t_2^{d'} - t_1)/\mu_{d'} \times \mu_{p^e}]$ where $\mu_{d'}$ acts on t_2 and μ_{p^e} on t_1 . So we have a space X (namely, $\mathrm{Spec} R[t_1, t_2]/(t_1^{p^e} - \pi, t_2^{d'} - t_1)$) with an action of a group of the form $G = G_1 \times G_2$ (namely, $G_1 = \mu_{p^e}$ and $G_2 = \mu_{d'}$). We can factor $X \rightarrow [X/G_1 \times G_2]$ as $X \rightarrow [X/G_1] \rightarrow [X/G_1 \times G_2]$. So we can then factor the fppf $\mu_{d'}$ -cover $\mathrm{Spec} R[t]/(t^d - \pi) \rightarrow \mathcal{R}$ as

$$\mathrm{Spec} R[t]/(t^d - \pi) \rightarrow {}^{p^e}\sqrt{\mathrm{Spec} R[\sqrt[d']{\pi}]} \xrightarrow{\xi} \mathcal{R}.$$

Observe that ξ is finite and étale as it is a $\mu_{d'}$ -torsor, so it suffices to prove the desired statement for $R[\sqrt[d']{\pi}]$, which is still a henselian DVR, and for $\mathcal{R} = {}^{p^e}\sqrt{\mathrm{Spec} R[\sqrt[d']{\pi}]}$: we can assume $d = p^e$ and $p > 0$.

Observe now that if k is a separably closed field of characteristic p , and T is a torus on $\mathcal{B}\mu_{p^e}$ over $\mathrm{Spec} k$, then $T \cong \mathcal{B}\mu_{p^e} \times \mathbb{G}_m^n$ for some n . Indeed, as T splits after the fppf-cover $\mathrm{Spec} k \rightarrow \mathcal{B}\mu_{p^e}$, T is given by a homomorphism $\mu_{p^e} \rightarrow \mathrm{Aut}(T) = \mathrm{GL}_n(\mathbb{Z})$. As $\mathrm{GL}_n(\mathbb{Z})$ is discrete over $\mathrm{Spec} k$, the only homomorphism is the trivial one.

Consider then R^{sh} the strict henselization of R , consider the morphisms $\mathrm{Spec} R^{sh} \rightarrow \mathrm{Spec} R$ and $\mathcal{R}^{sh} := \mathcal{R} \times_{\mathrm{Spec} R} \mathrm{Spec} R^{sh}$. Over \mathcal{R}^{sh} one can consider the isom-sheaf $\mathcal{I}^{sh} := \underline{\mathrm{Isom}}_{\mathrm{grp}}(\mathbb{G}_m^n \times \mathcal{R}^{sh}, T^{sh})$ of Lemma 2.4 where $T^{sh} = T|_{\mathcal{R}^{sh}}$. Then there is a section $(\mathcal{B}\mu_{p^e})_{k^s} \rightarrow \mathcal{I}$, where k^s is the separable closure of the residue field of R . As $\mathcal{I}^{sh} \rightarrow \mathcal{R}^{sh}$ is étale, this section lifts uniquely to any infinitesimal thickening of $(\mathcal{B}\mu_{p^e})_{k^s}$ in \mathcal{R}^{sh} , so from [AHR25, Theorem 3.4] it lifts to a section over \mathcal{R}^{sh} . As \mathcal{I}^{sh} is the pull-back of $\underline{\mathrm{Isom}}_{\mathrm{grp}}(\mathbb{G}_m^n \times \mathcal{R}, T)$ which is locally of finite type over \mathcal{R} , and as R^{sh} is the limit of finite and étale morphisms over R , there is a finite and étale morphism $\mathrm{Spec} R' \rightarrow \mathrm{Spec} R$ such that T trivializes when pulled back to $\mathcal{R} \times_{\mathrm{Spec} R} \mathrm{Spec} R'$; note this fiber product is a root stack over $\mathrm{Spec} R'$. \square

Lemma 4.6. *Let $p : \mathcal{Y} \rightarrow \mathcal{X}$ be a finite and étale morphism of degree d of algebraic stacks and $T \rightarrow \mathcal{Y}$ a smooth commutative scheme. Then p induces a morphism from the flat-fppf topos of \mathcal{Y} to the one of \mathcal{X} , and there are morphisms $T \rightarrow p_*p^{-1}T$ and $p_*p^{-1}T \rightarrow T$ whose composition is the multiplication by d .*

Observe that this is again well-known for schemes, on the small étale site [Sta21, Tag 03SH]. We will need the corresponding statement for the fppf-site, in the case when T is smooth; for doing so the key input is [Gro68, Theorem 11.7].

Proof. As p is finite and étale, from [Sta21, Tag 0GR1] it induces a morphism of topoi. In particular, there are functors $p_* : \mathcal{Sch}(\mathcal{Y}_{\mathrm{fppf}}) \rightarrow \mathcal{Sch}(\mathcal{X}_{\mathrm{fppf}})$ and $p^{-1} : \mathcal{Sch}(\mathcal{X}_{\mathrm{fppf}}) \rightarrow \mathcal{Sch}(\mathcal{Y}_{\mathrm{fppf}})$. The sheaf represented by $T \times_{\mathcal{X}} \mathcal{Y}$ agrees with $p^{-1}T$. Moreover, there is an adjunction morphism $\mathrm{Id} \rightarrow p_*p^{-1}$ that gives a map $T \rightarrow p_*p^{-1}T$. We claim that, as T is smooth, there is also a map $p_*p^{-1}T \rightarrow T$ such that the composition $T \rightarrow p_*p^{-1}T \rightarrow T$ is the multiplication by d .

To check this, it suffices to prove that for every flat morphism $U \rightarrow \mathcal{X}$ from a separated scheme

- (1) there is a map $p_*p^{-1}T(U) \rightarrow T(U)$, such that if precomposed with $T(U) \rightarrow p_*p^{-1}T(U)$ gives the multiplication by d , and
- (2) this map is functorial in U (i.e. it is a map of sheaves).

For (1), consider then the restriction morphisms $r_U : \mathcal{Sch}(U_{\mathrm{fppf}}) \rightarrow \mathcal{Sch}(U_{\mathrm{ét}})$ and $r_V : \mathcal{Sch}(V_{\mathrm{fppf}}) \rightarrow \mathcal{Sch}(V_{\mathrm{ét}})$ where $V = U \times_{\mathcal{X}} \mathcal{Y}$, and the subscript ét stands for the small étale site. Then:

- $T(U) = r_U(T|_U)(U)$. Indeed, we can pick an étale cover $\mathcal{U} \rightarrow U$ where \mathcal{U} is a disjoint union of affines (so also $\mathcal{U} \times_U \mathcal{U}$ will be affine), and from Grothendieck's theorem [Gro68, Theorem 11.7] $T(\mathcal{U}) = r_U T(\mathcal{U})$ and $T(\mathcal{U} \times_U \mathcal{U}) = r_U T(\mathcal{U} \times_U \mathcal{U})$ as T is smooth. Then from the sheaf properties, the global sections $T(U)$ and $r_U(T|_U)(U)$ are the colimit of the same diagram, so $T(U) = r_U(T|_U)(U)$.
- As we can compute p_* , p^{-1} and global sections in terms of limits of global sections of T on affine schemes which are étale over either V or U , proceeding as above $r_V(p^{-1}T|_U)$ and $p_{\mathrm{ét}}^{-1}r_U(T|_U)$ are canonically isomorphic and similarly with p_* , where we denoted by $p_{\mathrm{ét}}$ the morphism between the small étale sites $V_{\mathrm{ét}} \rightarrow U_{\mathrm{ét}}$.
- For the small étale sites we have morphisms $r_U(T|_U)(U) \rightarrow ((p_{\mathrm{ét}})_*(p_{\mathrm{ét}})^{-1})r_U(T|_U)(U) \rightarrow r_U(T|_U)(U)$ from [Sta21, Tag 03SH], so we also have morphisms $T(U) \rightarrow (p_*p^{-1}T)(U) \rightarrow T(U)$ whose composition is multiplication by d .

For part (2) instead, it suffices to check that if one has a cartesian diagram as follows

$$\begin{array}{ccc} V' & \longrightarrow & V \\ \downarrow & & \downarrow f \\ U' & \longrightarrow & U \end{array}$$

with $V \rightarrow U$ finite and étale, then the trace map of [Sta21, Tag 03SH] commutes with base change. Recall that if \mathcal{A} is a sheaf of abelian groups on the small étale site of U , the trace map is defined in [Sta21, Tag 03SH] as the composition of

$$\mathcal{A} \rightarrow (f_{\text{ét}})_*(f_{\text{ét}})^{-1}\mathcal{A} \cong (f_{\text{ét}})!(f_{\text{ét}})^{-1}\mathcal{A} \rightarrow \mathcal{A}$$

where $(f_{\text{ét}})_*$, $(f_{\text{ét}})^{-1}$ and $(f_{\text{ét}})!$ are maps between the categories abelian sheaves on the small étale site of U and V . The two arrows come from natural adjunctions, whereas the isomorphism comes from the map defined in [Sta21, Tag 0F4L], which is an isomorphism for f finite and étale [Sta21, Tag 03S7]. Then to check the desired statement, it suffices to observe that both the adjunction maps $r_U(T_U) \rightarrow (f_{\text{ét}})_*(f_{\text{ét}})^{-1}r_U(T_U)$ and $(f_{\text{ét}})!(f_{\text{ét}})^{-1}r_U(T_U) \rightarrow r_U(T_U)$, and the isomorphism $(f_{\text{ét}})_*(f_{\text{ét}})^{-1}r_U(T_U) \cong (f_{\text{ét}})!(f_{\text{ét}})^{-1}r_U(T_U)$ commute with base change.

As f is étale, the first map $\mathcal{A} \rightarrow (f_{\text{ét}})_*(f_{\text{ét}})^{-1}\mathcal{A}$ evaluated at an étale morphism $W \rightarrow U$ agrees with the restriction

$$\mathcal{A}(W) \rightarrow \mathcal{A}(W \times_U V).$$

The second map instead, up to shrinking W , agrees with the addition map

$$\bigoplus_{W \rightarrow V \text{ over } U} (f_{\text{ét}})^{-1}\mathcal{A}(W) = \bigoplus_{W \rightarrow V \text{ over } U} \mathcal{A}(W) \rightarrow \mathcal{A}(W)$$

as $(f_{\text{ét}})!$ is the sheafification of $(f_{\text{ét}})_p!$ [Sta21, Tag 03S2]. These commute with base change. Similarly the isomorphism $\mathcal{A}(W \times_U V) \rightarrow \bigoplus_{W \rightarrow V \text{ over } U} \mathcal{A}(W)$ defined in [Sta21, Tag 0F4L], where we shrank W so that there are $\deg(f)$ sections of $W \rightarrow V$ over U , again commutes with base change. \square

Corollary 4.7. *Let R be a henselian DVR, let \mathcal{R} be a root stack of $\text{Spec } R$ at its closed point and $T \rightarrow \mathcal{R}$ a torus. Then $H^i(\mathcal{R}, T)$ is torsion for $i = 1, 2$.*

Proof. From Proposition 4.5 there is a finite étale morphism $p : \mathcal{R}' \rightarrow \mathcal{R}$ of degree d such that $T|_{\mathcal{R}'} \cong \mathbb{G}_m^n$ for some n , with \mathcal{R}' the root stack of a DVR. From Lemma 4.6 the map

$$T \rightarrow p_*p^{-1}T \rightarrow T$$

of abelian fppf-sheaves of \mathcal{X} is the multiplication by d , for $i = 1, 2$. So there are maps whose composition is multiplication by d

$$H^i(\mathcal{R}, T) \rightarrow H^i(\mathcal{R}, p_*p^{-1}T) \rightarrow H^i(\mathcal{R}, T).$$

It suffices to check that $H^i(\mathcal{R}, p_*p^{-1}T) \cong H^i(\mathcal{R}', p^{-1}T)$. Indeed, \mathcal{R}' is such that $p^{-1}T = \mathbb{G}_m^n$ and $H^i(\mathcal{R}', \mathbb{G}_m^n)$ is torsion by Lemma 4.4. The equality $H^i(\mathcal{R}, p_*p^{-1}T) \cong H^i(\mathcal{R}', p^{-1}T)$ follows from the Leray spectral sequence, once we prove that $R^ip_*p^{-1}T = 0$ for $i > 0$; we will now prove this.

From [Sta21, Tag 0GR2] the sheaf $R^ip_*p^{-1}T$ is the sheafification of the presheaf that sends $U \rightarrow \mathcal{R}$ to $H^i(U \times_{\mathcal{R}} \mathcal{R}', p^{-1}T)$. In particular to check that $R^ip_*p^{-1}T = 0$ it suffices to check that every separated scheme $U \rightarrow \mathcal{R}$ admits an étale cover $\mathcal{U} \rightarrow U$ in affine schemes such that $H^i(\mathcal{U} \times_{\mathcal{R}} \mathcal{R}', p^{-1}T) = 0$. Let us denote by $V := \mathcal{R}' \times_{\mathcal{R}} U$ with the second projection f . As f is finite, $R^i(f_{\text{ét}})_*(p^{-1}T)|_{V_{\text{ét}}} = 0$ from [Sta21, Tag 0A4K], where the subscript ét stands for the restriction of f and $p^{-1}T$ to the small étale sites of V and U . Then we can find an étale cover $\mathcal{U} \rightarrow U$ of U in affine schemes such that $H_{\text{ét}}^i(\mathcal{U} \times_U V, p^{-1}T|_{V_{\text{ét}}}) = 0$. The vanishing of $R^ip_*p^{-1}T$ follows from Grothendieck's theorem [Gro68, Theorem 11.7]: étale cohomology of affine schemes with value in $p^{-1}T$ (which is smooth) agrees with fppf, so $H_{\text{ét}}^i(\mathcal{U} \times_U V, p^{-1}T|_{V_{\text{ét}}}) = H^i(\mathcal{U} \times_U V, p^{-1}T)$. Then $R^ip_*p^{-1}T = 0$ as every scheme $U \rightarrow \mathcal{R}$ admits a cover $\mathcal{U} \rightarrow U$ where $R^ip_*p^{-1}T(\mathcal{U}) = 0$, and $R^ip_*p^{-1}T$ is a sheaf. \square

Lemma 4.8. *Let \mathcal{R} be the root stack of the spectrum of a DVR R , let η be its generic point and let $T \rightarrow \mathcal{R}$ be a torus. Assume we are given $x_\eta \in H^1(\eta, T|_\eta)$. Then, up to replacing \mathcal{R} with a further root stack, we can extend x_η to $x \in H^1(\mathcal{R}, T)$.*

Proof. First observe that $H^1(\eta, T|_\eta)$ is torsion by Lemma 4.4. In the fppf topology of \mathcal{R} , we have the following exact sequence:

$$1 \rightarrow F \rightarrow T \xrightarrow{\cdot d} T \rightarrow 1$$

where d is the order of x_η . As T is a torus, F is tame. Then there is a finite tame group F over \mathcal{R} and an element $c_\eta \in H^1(\eta, F|_\eta)$ that maps to x_η . It suffices to extend c_η to $c \in H^1(\mathcal{R}, F)$. Observe that the element c_η corresponds to a morphism $\eta \rightarrow \mathcal{B}F$. As F is tame, from [BV24a] we can extend this morphism to a map $\mathcal{R} \rightarrow \mathcal{B}F$, up to replacing \mathcal{R} with a further root stack. This corresponds to an element $c_F \in H^1(\mathcal{R}, F)$ and its image in $H^1(\mathcal{R}, T)$ is the desired extension x . \square

Proposition 4.9. *Let \mathcal{R} be the root stack of a henselian DVR R . Let $T \rightarrow \mathcal{R}$ be a torus, and $\mathcal{T} \rightarrow \mathcal{R}$ a gerbe banded by T which is generically trivial. Then, up to replacing \mathcal{R} with a further root stack, \mathcal{T} is trivial.*

Proof. Let $c \in H^2(\mathcal{R}, T)$ be the class representing \mathcal{T} . First observe that by Corollary 4.7, the class c is torsion. Let d be the order of c , and consider the exact sequence in the fppf-topology

$$1 \rightarrow F \rightarrow T \xrightarrow{\cdot d} T \rightarrow 1.$$

As d is the order of c , there is $c_F \in H^2(\mathcal{R}, F)$ which maps to c via $\beta : H^2(\mathcal{R}, F) \rightarrow H^2(\mathcal{R}, T)$. Consider then the following commutative square, where η is the generic point of \mathcal{R} :

$$\begin{array}{ccccc} H^1(\mathcal{R}, T) & \xrightarrow{\alpha} & H^2(\mathcal{R}, F) & \xrightarrow{\beta} & H^2(\mathcal{R}, T) \\ \downarrow & & \downarrow & & \downarrow \\ H^1(\eta, T) & \xrightarrow{\alpha|_\eta} & H^2(\eta, F) & \xrightarrow{\beta|_\eta} & H^2(\eta, T). \end{array}$$

Since $\beta|_\eta((c_F)|_\eta) = 0$, there is an element $x_\eta \in H^1(\eta, T)$ that maps to $(c_F)|_\eta$. From Lemma 4.8, up to replacing \mathcal{R} with a further root stack, we can extend x_η to $x \in H^1(\mathcal{R}, T)$. In particular, up to replacing c_F with $c_F - \alpha(x)$, we can assume that $(c_F)|_\eta$ is trivial. So \mathcal{F} , the F -gerbe over \mathcal{R} corresponding to c_F , has a section over η . Thus by [BV24a] it has a section after replacing \mathcal{R} with a further root stack. In other terms, $c_F = 0$ after passing to a further root stack. Thus also $c = 0$. \square

4.2. Proof of Theorem 4.2. We are finally ready to prove Theorem 4.2; we begin with the following auxiliary statement.

Proposition 4.10. *Let R be a henselian DVR, let $G \rightarrow \mathcal{R}$ be a reductive group which is either special or has tame Weyl group (as in Definition 4.1) where \mathcal{R} is a root stack of $\text{Spec } R$ at its closed point. Let η be the generic point of \mathcal{R} . Then any section of $(\mathcal{B}G)_\eta \rightarrow \eta$ can be extended to a section of $\mathcal{R} \rightarrow \mathcal{B}G$ up to replacing \mathcal{R} with a further root stack.*

Proof. In the special case, there is nothing to prove as $\eta \rightarrow \mathcal{B}G$ classifies the trivial torsor which extends to the trivial torsor over \mathcal{R} . Thus suppose G is reductive with tame Weyl group. Let T be a maximal torus of G , which exists by Proposition 2.7 and let N be the normalizer of T in G . The inclusion $N \rightarrow G$ induces a map $H^1(\cdot, N) \rightarrow H^1(\cdot, G)$. By [Ser97, Lemma III.2.2.1 (b)] the class c_η lifts to an element $d_\eta \in H^1(\eta, N)$, as all maximal tori of a reductive group over an algebraically closed field are conjugate. In particular, it suffices to check that the map $d_\eta : \eta \rightarrow \mathcal{B}N_\eta$ lifts to a $d : \mathcal{R} \rightarrow \mathcal{B}N$, up to replacing \mathcal{R} with a further root stack. Recall that the normalizer of a maximal torus T is an extension of T by the Weyl group which by assumption is tame over \mathcal{R} :

$$1 \rightarrow T \rightarrow N \rightarrow W \rightarrow 1.$$

In particular, we can factor $\mathcal{B}N \rightarrow \mathcal{R}$ as $\mathcal{B}N \xrightarrow{\alpha} \mathcal{B}W \rightarrow \mathcal{R}$. By [BV24a], there is a lift of $\alpha \circ d_\eta$ up to replacing \mathcal{R} with a further root stack, so we have a morphism $\mathcal{R} \rightarrow \mathcal{B}W$ extending $\alpha \circ d_\eta$. We need now to lift it to $\mathcal{B}N \times_{\mathcal{B}W} \mathcal{R}$, but $\mathcal{B}N \times_{\mathcal{B}W} \mathcal{R} \rightarrow \mathcal{R}$ is a gerbe for a form of the torus T by [Sta21, Tag 0CJY]. Then Proposition 4.9 applies so that after replacing \mathcal{R} by a further root stack, we can extend $\eta \rightarrow \mathcal{B}N \times_{\mathcal{B}W} \mathcal{R}$ to a section $\mathcal{R} \rightarrow \mathcal{B}N \times_{\mathcal{B}W} \mathcal{R}$ as required. \square

Remark 4.11. In the special case of a gerbe over R itself rather than a root stack of $\mathrm{Spec} R$, the order of the root stack needed to extend the K -point of $\mathcal{B}G$ can be bounded uniformly in terms of the Weyl group of G . Indeed by work of Chernousov, Gille and Reichstein [CGR06, CGR08] there exists a finite subgroup scheme $S \subset G$ which is an extension of the Weyl group of G by a diagonalizable group such that $H^1(K, S) \rightarrow H^1(K, G)$ is surjective. Thus we can lift the K -point of $\mathcal{B}G$ to $\mathcal{B}S$. By the tameness assumption on the Weyl group of G , the stack $\mathcal{B}S$ is tame so we may apply [BV24a] to this stack and then the degree of the root stack necessary is bounded by the order of S .

We are now ready to prove Theorem 4.2.

Proof of Theorem 4.2. From Proposition 4.10, it suffices to show that, up to replacing \mathcal{R} with a further root stack, we have that $\mathcal{G} \cong \mathcal{B}G$. Since \mathcal{G} is banded by G , it corresponds to a class $c \in H^2(\mathcal{R}, G)$. Let Z be the center of G (which exists by [Con14, Lemma 2.2.4]). Observe that Z is linearly reductive as $Z \subseteq C(T)$ where $C(T)$ is the centralizer of a maximal torus T of G (which again exists by [Con14, Lemma 2.2.4]), and $C(T) = T$ as it is true on geometric fibers and both sides commute with base change by [Con14, Theorem 3.3.4]. It follows from [Gir65, Thm. IV.3.3.3] (see also the proof of [BES24, Proposition 2.1]) that the map $H^2(\mathcal{R}, Z) \rightarrow H^2(\mathcal{R}, G)$ and the map $H^2(\eta, Z) \rightarrow H^2(\eta, G)$ defined by sending a gerbe \mathcal{Z} to $(\mathcal{Z} \times \mathcal{B}G) // Z$ are bijections, where $(\mathcal{Z} \times \mathcal{B}G) // Z$ is the rigidification of $\mathcal{Z} \times \mathcal{B}G$ by the diagonal subgroup Z of the inertia stack $\mathcal{I}(\mathcal{Z} \times \mathcal{B}G) = \mathcal{I}\mathcal{Z} \times \mathcal{I}\mathcal{B}G$. In particular, it suffices to show that if Z is a commutative linearly reductive group over \mathcal{R} , and $\mathcal{Z} \rightarrow \mathcal{R}$ is a gerbe banded by Z which is generically trivial, it can be trivialized up to replacing \mathcal{R} with a further root stack. We can further rigidify by the smooth connected component of the identity $(Z^0)_{sm}$ ([AHR25, Theorem 9.9] and [BES24, Remark 2.2]) and factor $\mathcal{Z} \rightarrow \mathcal{R}$ as $\mathcal{Z} \rightarrow \mathcal{Z}' \rightarrow \mathcal{R}$ where $\mathcal{Z} \rightarrow \mathcal{Z}'$ is a gerbe for $(Z^0)_{sm}$, and $\mathcal{Z}' \rightarrow \mathcal{R}$ is finite and tame [BES24, Proposition 2.3]. Applying [BV24a] to the generically trivial tame gerbe $\mathcal{Z}' \rightarrow \mathcal{R}$, after replacing \mathcal{R} by a further root stack, we can assume that $\mathcal{Z}' \rightarrow \mathcal{R}$ has a section. Pulling back $\mathcal{Z} \rightarrow \mathcal{Z}'$ along this section, we obtain a gerbe for $(Z^0)_{sm}$ so it suffices to treat the case in which $(Z^0)_{sm} = Z$ is a smooth commutative connected reductive group, i.e. a torus. Then Proposition 4.9 applies so that \mathcal{Z} and thus \mathcal{G} is trivialized after replacing \mathcal{R} by a further root stack. \square

5. PROOFS OF THE MAIN THEOREMS AND APPLICATIONS

We are finally ready to prove Theorem 1.1 and Theorem 1.3.

Proof of Theorem 1.1. We first assume that R is henselian. From Proposition 3.2 it suffices to treat the case when the generic point of $\mathrm{Spec} R$ maps to a polystable point of \mathcal{X} . From Theorem 3.3 it suffices to assume that $\mathcal{X} \rightarrow X$ is a gerbe banded by a connected, smooth, linearly reductive group. This case is treated in Theorem 4.2. To reduce to the henselian case we use [Ryd11, Theorem B]. Take $\mathrm{Spec} A \rightarrow \mathrm{Spec} R$ the Henselization morphism. We know that the n -th root stack of $\mathrm{Spec} A$ at its closed point will admit a lifting to \mathcal{X} . But the following diagram is a pushout, which concludes the argument

$$\begin{array}{ccc} \mathrm{Spec} \mathrm{Frac}(A) & \longrightarrow & \sqrt[n]{\mathrm{Spec} A} \\ \downarrow & & \downarrow \\ \mathrm{Spec} \mathrm{Frac}(R) & \longrightarrow & \sqrt[n]{\mathrm{Spec} R}. \end{array}$$

\square

Proof of Theorem 1.3. When R is henselian, this is exactly Theorem 4.2. To reduce to the henselian case, we can use the same pushout diagram as in the above proof of Theorem 1.1. Finally, to handle extensions of groups, suppose we have an exact sequence $1 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 1$ of reductive groups over a root stack \mathcal{R} where G_1 has tame Weyl group and G_3 is special. Let $\mathcal{X}_2 \rightarrow \mathcal{R}$ be a gerbe banded by G_2 and $\mathrm{Spec} K \rightarrow \mathcal{X}_2$ be a K -point. As in the proof of Theorem 4.2, after passing to a further root stack of \mathcal{R} , we may assume that $\mathcal{X}_2 \cong \mathcal{B}G_2$. Composing with $\mathcal{B}G_2 \rightarrow \mathcal{B}G_3$ yields a K -point of $\mathcal{B}G_3$ which extends to a \mathcal{R} -point after taking a further root stack by assumption. Now consider $\mathcal{X}_1 := \mathcal{R} \times_{\mathcal{B}G_3} \mathcal{B}G_2$ with its induced K -point $\mathrm{Spec} K \rightarrow \mathcal{X}_1$. This is a gerbe banded by a form of G_1 so by assumption, the K -point extends after taking a further root stack. Composing with $\mathcal{X}_1 \rightarrow \mathcal{X}_2 = \mathcal{B}G_2$ yields the required extension. \square

Proof of Proposition 1.5 and Corollary 1.6. For Proposition 1.5, let $x_1, \dots, x_m \in C$ be points of indeterminacy for φ and let U be the open complement so that φ restricts to a morphism $U \rightarrow \mathcal{X}$. Let $R_i = \mathcal{O}_{C, x_i}$ which is a DVR and let $K = k(C) = \text{Frac } R_i$ be the function field. The K -point $\text{Spec } K \rightarrow \mathcal{X}$ extends to root stack points $\sqrt[n]{\text{Spec } R_i} \rightarrow \mathcal{X}$ by properness of X and Theorem 1.1 or by Theorem 1.3 for some integers n_i . Let $\mathcal{C} \rightarrow C$ be the global root stack $\sqrt[n_1, \dots, n_m]{(C, x_1, \dots, x_m)}$. We wish to extend $U \rightarrow \mathcal{X}$ to a morphism $\mathcal{C} \rightarrow \mathcal{X}$. By induction on m , we can extend the morphism one point at a time so suppose without loss of generality that $m = 1$. Consider the diagram

$$\begin{array}{ccc} \text{Spec } K & \longrightarrow & \sqrt[n]{\text{Spec } R} \\ \downarrow & & \downarrow \\ U & \longrightarrow & \mathcal{C} \end{array}$$

This is a flat Mayer-Vietoris square in the sense of [HR23, Definition 1.2] and \mathcal{C} is locally excellent by assumption so this square is a pushout square in algebraic stacks by [HR23, Theorem A]. Thus there is a morphism $\mathcal{C} \rightarrow \mathcal{X}$ by the universal property of pushouts.

For Corollary 1.6, note that \mathcal{X} is locally of finite presentation by assumption so the given K -point $\text{Spec } K \rightarrow \mathcal{X}$ spreads out to a morphism $U \rightarrow \mathcal{X}$ for some open set $U \subset C$ [Sta21, Tag 0CMX]. The result then follows from Proposition 1.5. \square

Proof of Corollary 1.9. By Theorem 1.3 and Remark 4.11, there exists an n depending only on G such that any K -point extends to a $\sqrt[n]{\text{Spec } R}$ point of $\mathcal{B}G$. \square

Proof of Proposition 1.14 and Corollary 1.15. Exactly the same argument as in [BV24a, Theorem 4.1] goes through, except we replace [BV24a, Corollary 3.2] with Theorem 1.1 which gives a morphism $\mathcal{R} \rightarrow \mathcal{Y}$ that, if pre-composed with the covering $\text{Spec}(R) \rightarrow \mathcal{R}$ and the closed embedding of the closed point $\text{Spec}(k_R) \rightarrow \text{Spec}(R)$, gives the desired point on \mathcal{Y} .

For Corollary 1.15, we can apply Proposition 1.14 to the rational map $\text{Spec } R \dashrightarrow [\text{Spec } A/G]$ to get a k -point of $[\text{Spec } A \otimes_R k/G_k]$ which lifts to a k -point of $A \otimes_R k$ by the special assumption. \square

Proof of Proposition 1.16. Let $\mathcal{G} \in H^2(R, G)$ such that $\mathcal{G}|_K = 0$. Then $\mathcal{G}_K \cong \mathcal{B}G_K$ so there exists a point $\text{Spec } K \rightarrow \mathcal{G}$ which by Theorem 1.3 extends to a root stack $\mathcal{R}_n \rightarrow \mathcal{G}$. Composing with $\text{Spec } k \rightarrow \mathcal{R}_n$ yields a k -point of $\mathcal{G}|_k$ which exhibits the triviality of $\mathcal{G}|_k$. \square

Proof of Corollary 1.17 and Corollary 1.18. Let $x_K : \text{Spec } K \rightarrow [V/H]$ be the composition of a K -point of V with the quotient map. This exhibits the triviality of the gerbe $[V/H]_K$. By Proposition 1.16 the gerbe $[V/H]_k$ is also trivial and since H is special, any k -point of $[V/H]_k$ lifts to a k -point of V .

If V_K is unirational then it admits a K -point. Thus V_k also admits a k -point. By [Bor91, Theorem 18.2(ii)], H_k is unirational and given a k -point $x \in V(k)$, we get a surjection $H_k \rightarrow V_k$ by $h \mapsto hx$ which implies V_k is unirational. \square

Remark 5.1. In fact Theorem 1.3 applied to $[V/H]$ shows something slightly stronger. Given any K -point $x_K \in V(K)$, there exists a root extension $R' = R[t^{1/n}]$ where t is a uniformizer and an element $h \in H(R')$ such that hx_K extends to an R' -point of V .

Proof of Corollary 1.19 and Corollary 1.20. The K -moduli stack $\mathcal{M}^{K\text{-ss}}$ of K -semistable Fano varieties has a proper good moduli space $M^{K\text{-ps}}$ by [LWX19, BLX22, ABHLX20, LXZ22]. Thus by Theorem 1.1, there exists a root stack \mathcal{R}_n and a family $\mathcal{Y} \rightarrow \mathcal{R}_n$ of K -semistable Fano varieties with polystable central fiber extending $Y_\eta \rightarrow \text{Spec } K$. The total space has klt singularities by [Oda13, Theorem 1.3] and inversion of adjunction for klt singularities [fai92, Sections 16 & 17] and quotients of klt singularities are klt so the coarse space $Y \rightarrow \text{Spec } R$ has klt singularities and central which is a quotient of a K -polystable Fano variety by μ_n . The multiplicity n can be deduced by noting that $\mathcal{R}_n \rightarrow \text{Spec } R$ is totally ramified of order n (see e.g. [BPS24, Lemma 7.11]).

For the moduli of boundary polarized Calabi-Yau surface pairs \mathcal{M}^{CY} , the argument is similar, except here we only have an *asymptotically good moduli space* [ABB⁺23, Definition 13.1]. More precisely, we bound the Cartier index of K_Y to obtain an open substack \mathcal{M}_m^{CY} which has a good moduli space M_m^{CY} . By the proofs of [ABB⁺23, Theorem 1.3] and [BL24, Theorem 1.1], the good moduli space is proper and independent of

m for $m \gg 0$. Thus we apply Theorem 1.3 to \mathcal{M}_m^{CY} for m sufficiently large. Then the proof is exactly as above using that quotients of slc singularities are slc [AP12, Lemma 2.3] and inversion of adjunction for log canonical singularities [Kaw07, Main Theorem]. \square

REFERENCES

- [AB19] Kenneth Ascher and Dori Bejleri. Moduli of fibered surface pairs from twisted stable maps. *Math. Ann.*, 374(1-2):1007–1032, 2019.
- [ABB⁺23] Kenneth Ascher, Dori Bejleri, Harold Blum, Kristin DeVleming, Giovanni Inchiostro, Yuchen Liu, and Xiaowei Wang. Moduli of boundary polarized Calabi-Yau pairs. *arXiv e-prints*, page arXiv:2307.06522, July 2023.
- [ABHLX20] Jarod Alper, Harold Blum, Daniel Halpern-Leistner, and Chenyang Xu. Reductivity of the automorphism group of K -polystable Fano varieties. *Invent. Math.*, 222(3):995–1032, 2020.
- [Ach24] Niven Achenjang. On brauer groups of tame stacks, 2024.
- [ACV03] Dan Abramovich, Alessio Corti, and Angelo Vistoli. Twisted bundles and admissible covers. volume 31, pages 3547–3618. 2003. Special issue in honor of Steven L. Kleiman.
- [AH11] Dan Abramovich and Brendan Hassett. Stable varieties with a twist. In *Classification of algebraic varieties*, pages 1–38, 2011.
- [AHLR24] Jarod Alper, Jack Hall, Daniel Halpern-Leistner, and David Rydh. Artin algebraization for pairs with applications to the local structure of stacks and ferrand pushouts. In *Forum of Mathematics, Sigma*, volume 12, page e20. Cambridge University Press, 2024.
- [AHLH23] Jarod Alper, Daniel Halpern-Leistner, and Jochen Heinloth. Existence of moduli spaces for algebraic stacks. *Inventiones mathematicae*, 234(3):949–1038, 2023.
- [AHR25] Jarod Alper, Jack Hall, and David Rydh. The étale local structure of algebraic stacks. *arXiv e-prints*, page <https://arxiv.org/pdf/1912.06162.pdf>, April 2025.
- [Alp13] Jarod Alper. Good moduli spaces for Artin stacks. *Ann. Inst. Fourier (Grenoble)*, 63(6):2349–2402, 2013.
- [Alp14] Jarod Alper. Adequate moduli spaces and geometrically reductive group schemes. *Algebr. Geom.*, 1(4):489–531, 2014.
- [AOV11] Dan Abramovich, Martin Olsson, and Angelo Vistoli. Twisted stable maps to tame Artin stacks. *J. Algebraic Geom.*, 20(3):399–477, 2011.
- [AP12] Valery Alexeev and Rita Pardini. Non-normal abelian covers. *Compositio Mathematica*, 148(4):1051–1084, 2012.
- [AV02] Dan Abramovich and Angelo Vistoli. Compactifying the space of stable maps. *J. Amer. Math. Soc.*, 15(1):27–75, 2002.
- [BĈE⁺25] V. Balaji, Kestutis Česnavičius, Elden Elmanto, Arnab Kundu, and Alex Youcis. Grothendieck-Serre phenomena for parahoric groups. In preparation, 2025.
- [BES24] Dori Bejleri, Elden Elmanto, and Matthew Satriano. Proper splittings and projectivity for good moduli spaces. *arXiv preprint arXiv:2408.11057*, 2024.
- [BHMR17] Michael Bate, Sebastian Herpel, Benjamin Martin, and Gerhard Röhrle. Cocharacter-closure and the rational Hilbert–Mumford theorem. *Mathematische Zeitschrift*, 287:39–72, 2017.
- [BIMS25] Dori Bejleri, Giovanni Inchiostro, Siddharth Mathur, and Matthew Satriano. A primer on good moduli space gerbes. In preparation, 2025.
- [BL24] Harold Blum and Yuchen Liu. Good moduli spaces for boundary polarized calabi-yau surface pairs, 2024.
- [BLX22] Harold Blum, Yuchen Liu, and Chenyang Xu. Openness of K -semistability for Fano varieties. *Duke Math. J.*, 171(13):2753–2797, 2022.
- [BLXZ25] Harold Blum, Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. Relative stability theory of properness of k -moduli. In preparation, 2025.
- [Bor91] Armand Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1991.
- [BPS24] Dori Bejleri, Jun-Yong Park, and Matthew Satriano. Height moduli on cyclotomic stacks and counting elliptic curves over function fields, 2024.
- [BRV11] Patrick Brosnan, Zinovy Reichstein, and Angelo Vistoli. Essential dimension of moduli of curves and other algebraic stacks. *J. Eur. Math. Soc. (JEMS)*, 13(4):1079–1112, 2011. With an appendix by Najmuddin Fakhruddin.
- [BS15] V. Balaji and C. S. Seshadri. Moduli of parahoric \mathcal{G} -torsors on a compact Riemann surface. *J. Algebraic Geom.*, 24(1):1–49, 2015.
- [BV24a] Giulio Bresciani and Angelo Vistoli. An arithmetic valuative criterion for proper maps of tame algebraic stacks. *manuscripta mathematica*, 173(3):1061–1071, 2024.
- [BV24b] Giulio Bresciani and Angelo Vistoli. Fields of moduli and the arithmetic of tame quotient singularities. *Compos. Math.*, 160(5):982–1003, 2024.
- [CGR06] Vladimir Chernousov, Philippe Gille, and Zinovy Reichstein. Resolving g -torsors by abelian base extensions. *Journal of Algebra*, 296(2):561–581, 2006.
- [CGR08] Vladimir Chernousov, Philippe Gille, and Zinovy Reichstein. Reduction of structure for torsors over semilocal rings. *manuscripta mathematica*, 126:465–480, 2008.
- [Con14] Brian Conrad. Reductive group schemes. <http://math.stanford.edu/~conrad/papers/luminygsa3.pdf>, 2014. Société mathématique de France.

- [Dam24] Chiara Damiolini. On equivariant bundles and their moduli spaces. *C. R. Math. Acad. Sci. Paris*, 362:55–62, 2024.
- [DGA⁺11] M Demazure, A Grothendieck, M Artin, JE Bertin, P Gabriel, M Raynaud, and JP Serre. Séminaire de géométrie algébrique du bois marie 1962-64. schémas en groupes (SGA 3). Tome III: Structures des schémas en groupes reductif, 2011.
- [DLI22] Andrea Di Lorenzo and Giovanni Inchiostro. Degenerations of twisted maps to algebraic stacks. *arXiv preprint arXiv:2210.03806*, 2022.
- [DLI24] Andrea Di Lorenzo and Giovanni Inchiostro. Effective morphisms and quotient stacks. *Int. Math. Res. Not. IMRN*, (11):9091–9113, 2024.
- [Dou75] Jean-Claude Douai. 2-cohomologie galoisienne des groupes semi-simples définis sur les corps locaux. *C. R. Acad. Sci. Paris Sér. A-B*, 280(6):Aii, A321–A323, 1975.
- [DY22] Ratko Darda and Takehiko Yasuda. The Batyrev-Manin conjecture for DM stacks. *arXiv e-prints*, page arXiv:2207.03645, 2022.
- [EHKV01] Dan Edidin, Brendan Hassett, Andrew Kresch, and Angelo Vistoli. Brauer groups and quotient stacks. *American Journal of Mathematics*, 123(4):761–777, 2001.
- [ER21] Dan Edidin and David Rydh. Canonical reduction of stabilizers for Artin stacks with good moduli spaces. *Duke Math. J.*, 170(5):827–880, 2021.
- [ESZB23] Jordan S. Ellenberg, Matthew Satriano, and David Zureick-Brown. Heights on stacks and a generalized Batyrev-Manin-Malle conjecture. *Forum Math. Sigma*, 11:Paper No. e14, 54, 2023.
- [fli92] *Flips and abundance for algebraic threefolds*. Société Mathématique de France, Paris, 1992. Papers from the Second Summer Seminar on Algebraic Geometry held at the University of Utah, Salt Lake City, Utah, August 1991, Astérisque No. 211 (1992).
- [GCP24] Philippe Gille, Vladimir Chernousov, and Arturo Pianzola. Loop torsors. theory and applications. *arXiv preprint arXiv:2412.07294*, 2024.
- [Gir65] Jean Giraud. Cohomologie non abélienne; préliminaires. *C. R. Acad. Sci. Paris*, 260:2392–2394, 1965.
- [Gro68] Alexander Grothendieck. Le groupe de Brauer. II. Théorie cohomologique. In *Dix exposés sur la cohomologie des schémas*, volume 3 of *Adv. Stud. Pure Math.*, pages 67–87. North-Holland, Amsterdam, 1968.
- [GWZ25] Michael Groechenig, Dimitri Wyss, and Paul Ziegler. Twisted points of quotient stacks, integration and bps-invariants, 2025.
- [HR23] Jack Hall and David Rydh. Mayer-Vietoris squares in algebraic geometry. *J. Lond. Math. Soc. (2)*, 107(5):1583–1612, 2023.
- [Jeo25] Myeong Jae Jeon. Resolution of indeterminacy of rational maps to proper tame stacks, 2025.
- [Kaw07] Masayuki Kawakita. Inversion of adjunction on log canonicity. *Invent. Math.*, 167(1):129–133, 2007.
- [Kem78] George R Kempf. Instability in invariant theory. *Annals of Mathematics*, 108(2):299–316, 1978.
- [Lie08] Max Lieblich. Twisted sheaves and the period-index problem. *Compositio Mathematica*, 144(1):1–31, 2008.
- [LWX19] Chi Li, Xiaowei Wang, and Chenyang Xu. On the proper moduli spaces of smoothable Kähler-Einstein Fano varieties. *Duke Math. J.*, 168(8):1387–1459, 2019.
- [LXZ22] Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. Finite generation for valuations computing stability thresholds and applications to k-stability. *Annals of Mathematics*, 196(2):507–566, 2022.
- [Mil13] J. S. Milne. *Séminaire de géométrie algébrique du Bois Marie 1962–64. Schémas en groupes (SGA 3)*. Tom I, Tom III [book review of mr2867621; mr2867622]. *Eur. Math. Soc. Newsl.*, (87):50–52, 2013.
- [MT15] Johan Martens and Michael Thaddeus. Variations on a theme of Grothendieck, 2015.
- [Oda13] Yuji Odaka. The GIT stability of polarized varieties via discrepancy. *Ann. of Math. (2)*, 177(2):645–661, 2013.
- [Ols16] Martin Olsson. *Algebraic spaces and stacks*, volume 62. American Mathematical Soc., 2016.
- [PR24] Georgios Pappas and Michael Rapoport. On tamely ramified \mathcal{G} -bundles on curves. *Algebr. Geom.*, 11(6):796–829, 2024. With an appendix by Brian Conrad.
- [Rey25] Javier Reyes. Classification of singular fibers of hyperelliptic fibrations. In preparation, 2025.
- [RS22] Zinovy Reichstein and Federico Scavia. The behavior of essential dimension under specialization. *Épjournal Géom. Algébrique*, 6:Art. 21, 28, 2022.
- [Ryd11] David Rydh. Étale dévissage, descent and pushouts of stacks. *Journal of Algebra*, 331(1):194–223, 2011.
- [Ser97] Jean-Pierre Serre. *Galois cohomology*. Springer-Verlag, Berlin, 1997. Translated from the French by Patrick Ion and revised by the author.
- [Ser10] Jean-Pierre Serre. Bounds for the orders of the finite subgroups of $G(k)$. *arXiv preprint arXiv:1011.0346*, 2010.
- [SSW24] Mao Sheng, Hao Sun, and Jianping Wang. A nonabelian hodge correspondence for principal bundles in positive characteristic, 2024.
- [Sta21] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2021.
- [SU24] Matthew Satriano and Jeremy Usatine. Beyond twisted maps: applications to motivic integration, 2024.

(DB) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD, 20742
Email address: **dbejleri@umd.edu**

(GI) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON, USA
Email address: **ginchios@uw.edu**

(MS) DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO ON N2L3G1, CANADA
Email address: **msatrian@uwaterloo.ca**