

THE HIGHER ORDER FRACTIONAL CALDERÓN PROBLEM FOR LINEAR LOCAL OPERATORS: UNIQUENESS

GIOVANNI COVI, KELJO MÖNKKÖNEN, JESSE RAILO, AND GUNTHER UHLMANN

ABSTRACT. We study an inverse problem for the fractional Schrödinger equation (FSE) with a local perturbation by a linear partial differential operator (PDO) of the order smaller than the order of the fractional Laplacian. We show that one can uniquely recover the coefficients of the PDO from the Dirichlet-to-Neumann (DN) map associated to the perturbed FSE. This is proved for two classes of coefficients: coefficients which belong to certain spaces of Sobolev multipliers and coefficients which belong to fractional Sobolev spaces with bounded derivatives. Our study generalizes recent results for the zeroth and first order perturbations to higher order perturbations.

1. INTRODUCTION

Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $\Omega \subset \mathbb{R}^n$ a bounded open set where $n \geq 1$, $\Omega_e = \mathbb{R}^n \setminus \overline{\Omega}$ its exterior and $P(x, D)$ a linear partial differential operator (PDO) of order $m \in \mathbb{N}$

$$P(x, D) = \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha$$

where the coefficients $a_\alpha = a_\alpha(x)$ are functions defined in Ω . We study a nonlocal inverse problem for the perturbed fractional Schrödinger equation

$$(1) \quad \begin{cases} (-\Delta)^s u + P(x, D)u = 0 & \text{in } \Omega \\ u = f & \text{in } \Omega_e \end{cases}$$

where $(-\Delta)^s$ is a nonlocal pseudo-differential operator $(-\Delta)^s u = \mathcal{F}^{-1}(|\cdot|^{2s} \hat{u})$ in contrast to the local operator $P(x, D)$. In the inverse problem, one aims to recover the local operator P from the associated Dirichlet-to-Neumann map.

We always assume that 0 is not a Dirichlet eigenvalue of the operator $((-\Delta)^s + P(x, D))$, i.e.

If $u \in H^s(\mathbb{R}^n)$ solves $((-\Delta)^s + P(x, D))u = 0$ in Ω and $u|_{\Omega_e} = 0$, then $u = 0$.

Our data for the inverse problem is the Dirichlet-to-Neumann (DN) map $\Lambda_P: H^s(\Omega_e) \rightarrow (H^s(\Omega_e))^*$ which maps Dirichlet exterior values to a nonlocal version of the Neumann boundary value (see section 2 and 3.1). The main question that we study in this article is whether the DN map Λ_P determines uniquely the coefficients a_α in Ω . In other words, does $\Lambda_{P_1} = \Lambda_{P_2}$ imply that $a_{1,\alpha} = a_{2,\alpha}$ in Ω for all $|\alpha| \leq m$? We prove that the answer is positive under certain restrictions on the coefficients a_α and the order of the PDOs.

This gives positive answer to the uniqueness problem [10, Question 2.5] posed by the first three authors in a previous work. The precise statement in [10] asks to prove uniqueness for the higher order fractional Calderón problem in the case of a bounded domain with smooth boundary and PDOs with smooth coefficients (up to the boundary). The positive answer to this question follows from theorem 1.2. The study of the fractional Calderón problem was initiated by Ghosh, Salo and Uhlmann in the work [15] where the uniqueness for the associated inverse problem is proved when $m = 0$, $s \in (0, 1)$ and $a_0 \in L^\infty(\Omega)$.

Date: August 25, 2020.

Key words and phrases. Inverse problems, fractional Calderón problem, fractional Schrödinger equation, Sobolev multipliers.

We briefly note that by Peetre's theorem any linear operator $L: C_c^\infty(\Omega) \rightarrow C_c^\infty(\Omega)$ which does not increase supports, i.e. $\text{spt}(Lf) \subset \text{spt}(f)$ for all $f \in C_c^\infty(\Omega)$, is in fact a differential operator [30] (see also the original work [32]). Therefore our results apply to any local operator satisfying such properties and it is enough to study PDOs only. For a more general formulation of Peetre's theorem on the level of vector bundles, see [31].

1.1. Main results. We denote by $M(H^{s-|\alpha|} \rightarrow H^{-s})$ the space of all bounded Sobolev multipliers between the Sobolev spaces $H^{s-|\alpha|}(\mathbb{R}^n)$ and $H^{-s}(\mathbb{R}^n)$. We denote by $M_0(H^{s-|\alpha|} \rightarrow H^{-s}) \subset M(H^{s-|\alpha|} \rightarrow H^{-s})$ the space of bounded Sobolev multipliers that can be approximated with smooth compactly supported functions in the multiplier norm of $M(H^{s-|\alpha|} \rightarrow H^{-s})$. We also write $H^{r,\infty}(\Omega)$ for the local Bessel potential space with bounded derivatives. See section 2 for more detailed definitions.

Our first theorem is a generalization of [36, Theorem 1.1] which considered the case $m = 0$ with $s \in (0, 1)$. It also generalizes [10, Theorem 1.5] which considered the higher order cases $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ when $m = 0$.

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set where $n \geq 1$. Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $m \in \mathbb{N}$ be such that $2s > m$. Let*

$$P_j = \sum_{|\alpha| \leq m} a_{j,\alpha} D^\alpha, \quad j = 1, 2,$$

be linear PDOs of order m with coefficients $a_{j,\alpha} \in M_0(H^{s-|\alpha|} \rightarrow H^{-s})$. Given any two open sets $W_1, W_2 \subset \Omega_e$, suppose that the DN maps Λ_{P_i} for the equations $((-\Delta)^s + P_j)u = 0$ in Ω satisfy

$$\Lambda_{P_1} f|_{W_2} = \Lambda_{P_2} f|_{W_2}$$

for all $f \in C_c^\infty(W_1)$. Then $P_1|_\Omega = P_2|_\Omega$.

In theorem 1.1 one can pick the lower order coefficients ($|\alpha| < s$) from $L^p(\Omega)$ for high enough p (especially from $L^\infty(\Omega)$) and higher order coefficients ($s < |\alpha| < 2s$) from the closure of $C_c^\infty(\Omega)$ in $H^{r,\infty}(\Omega)$ for certain values of $r \in \mathbb{R}$. Lemmas 2.8 and 2.9 give more examples of Sobolev spaces which belong to the space of multipliers $M_0(H^{s-|\alpha|} \rightarrow H^{-s})$. We also note that when $|\alpha| = 0$, then the space of multipliers $M_0(H^s \rightarrow H^{-s})$ coincides with the one studied in [36].

It follows that the space of multipliers is trivial for higher order operators, i.e. $M(H^{s-|\alpha|} \rightarrow H^{-s}) = \{0\}$ when $s - |\alpha| < -s$. It would be possible to state theorem 1.1 for higher order PDOs, but that forces $a_\alpha = 0$ for all $|\alpha| > 2s$. For this reason we only consider PDOs whose order is $m < 2s$. See lemma 2.5 and the related remarks for more details.

Our second theorem generalizes [7, Theorem 1.1] and [15, Theorem 1.1] where similar results are proved when $m = 0, 1$ and $s \in (0, 1)$. It also generalizes [10, Theorem 1.5] where the case $m = 0$ and $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ was studied.

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain where $n \geq 1$. Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $m \in \mathbb{N}$ be such that $2s > m$. Let*

$$P_j(x, D) = \sum_{|\alpha| \leq m} a_{j,\alpha}(x) D^\alpha, \quad j = 1, 2,$$

be linear PDOs of order m with coefficients $a_{j,\alpha} \in H^{r_\alpha, \infty}(\Omega)$ where

$$(2) \quad r_\alpha := \begin{cases} 0 & \text{if } |\alpha| - s < 0, \\ |\alpha| - s + \delta & \text{if } |\alpha| - s \in \{1/2, 3/2, \dots\}, \\ |\alpha| - s & \text{if } \text{otherwise} \end{cases}$$

for any fixed $\delta > 0$. Given any two open sets $W_1, W_2 \subset \Omega_e$, suppose that the DN maps Λ_{P_i} for the equations $((-\Delta)^s + P_j(x, D))u = 0$ in Ω satisfy

$$\Lambda_{P_1} f|_{W_2} = \Lambda_{P_2} f|_{W_2}$$

for all $f \in C_c^\infty(W_1)$. Then $P_1(x, D) = P_2(x, D)$.

Our first theorem is formulated for general bounded open sets and the second theorem for Lipschitz domains. The difference arises in the proof of the well-posedness of the inverse problem. We note that theorem 1.2 holds for coefficients a_α which are smooth up to the boundary ($a_\alpha = g|_\Omega$ where $g \in C^\infty(\mathbb{R}^n)$). The conditions (1.2) imply that one can choose $a_\alpha \in L^\infty(\Omega)$ for every α such that $|\alpha| < s$. The case $|\alpha| = s$ never happens, as s is assumed not to be an integer. If $|\alpha| > s$, we have $a_\alpha \in H^{|\alpha|-s, \infty}(\Omega)$ when $|\alpha| - s \notin \{1/2, 3/2, \dots\}$. Thus the conditions (1.2) coincide with [7, 15] when $m = 0, 1$ and $s \in (0, 1)$.

Our article is roughly divided into two parts. The first part of the article (theorem 1.1 and section 3) generalizes the study of the uniqueness problem for singular potentials in [36] and the second part (theorem 1.2 and section 4) generalizes the uniqueness problem for bounded first order perturbations in [7].

The approach to prove theorems 1.1 and 1.2 is the following. First one shows that the inverse problem is well-posed and the corresponding bilinear forms are bounded. This leads to the boundedness of the DN maps and an Alessandrini identity. By a unique continuation property of the higher order fractional Laplacian one obtains a Runge approximation property for equation (1). Using the Runge approximation and the Alessandrini identity for suitable test functions one proves the uniqueness of the inverse problem.

1.2. On the earlier literature. Equation (1) and theorems 1.1 and 1.2 are related to the Calderón problem for the fractional Schrödinger equation first introduced in [15]. There one tries to uniquely recover the potential q in Ω by doing measurements in the exterior Ω_e . This is a nonlocal (fractional) counterpart of the classical Calderón problem arising in electrical impedance tomography where one obtains information about the electrical properties of some bounded domain by doing voltage and current measurements on the boundary [39, 40]. In [36] the study of the fractional Calderón problem is extended for “rough” potentials q , i.e. potentials which are in general bounded Sobolev multipliers. First order perturbations were studied in [7] assuming that the fractional part dominates the equation, i.e. $s \in (1/2, 1)$, and that the perturbations have bounded fractional derivatives. A higher order version ($s \in \mathbb{R}^+ \setminus \mathbb{Z}$) of the fractional Calderón problem was introduced and studied in [10]. These three articles [7, 10, 36] motivate the study of higher order (rough) perturbations to the fractional Laplacian $(-\Delta)^s$ in equation (1). The natural restriction for the order of $P(x, D)$ in theorems 1.1 and 1.2 is then $2s > m$ so that the fractional part governs the equation (1).

The fractional Calderón problem for $s \in (0, 1)$ has been studied in many settings. We refer to the survey [37] for a more detailed treatment. In the work [36] stability was proved for singular potentials, and in [34] the related exponential instability was shown. The fractional Calderón problem has also been solved under single measurement [14]. The perturbed equation is related to the fractional magnetic Schrödinger equation which is studied in [9, 24, 25, 26]. See also [4] for a fractional Schrödinger equation with a lower order nonlocal perturbation. Other variants of the fractional Calderón problem include semilinear fractional (magnetic) Schrödinger equation [19, 20, 24, 25], fractional heat equation [21, 35] and fractional conductivity equation [8] (see also [6, 13] for equations arising from a nonlocal Schrödinger-type elliptic operator). In the recent work [10], the first three authors of this article studied higher order versions ($s \in \mathbb{R}^+ \setminus \mathbb{Z}$) of the fractional Calderón problem and proved uniqueness for the Calderón problem for the fractional magnetic Schrödinger equation (up to a gauge). This article continues these studies by showing uniqueness for the fractional Schrödinger equation with higher order perturbations and gives positive answer to the question 2.5 posed in [10].

1.3. Examples of fractional models in the sciences. Equations involving fractional Laplacians like (1) have applications in mathematics and natural sciences. Fractional Laplacians appear in the study of anomalous and nonlocal diffusion, and these diffusion phenomena can be used in many areas such as continuum mechanics, graph theory and ecology just to mention a few [2, 5, 12, 27, 33]. Another place where the fractional counterpart of the classical Laplacian naturally shows up is the formulation of fractional quantum mechanics [22, 23]. For more applications of fractional mathematical models, see [5] and the references therein.

1.4. The organization of the article. In section 2 we introduce the notation and give preliminaries on Sobolev spaces and fractional Laplacians. We also define the spaces of rough coefficients (Sobolev multipliers) and discuss some of the basic properties. In section 3 we prove theorem 1.1 in detail. Finally, in section 4 we prove theorem 1.2 but as the proofs of both theorems are very similar we do not repeat all identical steps and we keep our focus in the differences of the proofs.

Acknowledgements. G.C. was partially supported by the European Research Council under Horizon 2020 (ERC CoG 770924). K.M. and J.R. were supported by Academy of Finland (Centre of Excellence in Inverse Modelling and Imaging, grant numbers 284715 and 309963). G.U. was partly supported by NSF, a Walker Family Endowed Professorship at UW and a Si Yuan Professorship at IAS, HKUST. G.C. would like to thank Angkana Rüland for helpful discussions and her hospitality during his visit to Max Planck Institute for Mathematics in the Sciences. J.R. and G.U. wish to thank Maarten V. de Hoop, Rice University and Simons Foundation for providing the support to participate 2020 MATH + X Symposium on Inverse Problems and Deep Learning, Mitigating Natural Hazards, where this project was initiated.

2. PRELIMINARIES

In this section we recall some basic theory of Sobolev spaces, Fourier analysis and fractional Laplacians on \mathbb{R}^n . We also introduce the spaces of Sobolev multipliers and prove a few properties for them. Some auxiliary lemmas which are needed in the proofs of our main theorems are given as well. We follow the references [1, 15, 29, 28, 38, 41] (see also section 3 in [10]).

2.1. Sobolev spaces. The (inhomogeneous) fractional L^2 -based Sobolev space of order $r \in \mathbb{R}$ is defined to be

$$H^r(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n) : \mathcal{F}^{-1}(\langle \cdot \rangle^r \hat{u}) \in L^2(\mathbb{R}^n)\}$$

equipped with the norm

$$\|u\|_{H^r(\mathbb{R}^n)} = \|\mathcal{F}^{-1}(\langle \cdot \rangle^r \hat{u})\|_{L^2(\mathbb{R}^n)}.$$

Here $\hat{u} = \mathcal{F}(u)$ is the Fourier transform of a tempered distribution $u \in \mathcal{S}'(\mathbb{R}^n)$, \mathcal{F}^{-1} is the inverse Fourier transform and $\langle x \rangle = (1 + |x|^2)^{1/2}$. We define the fractional Laplacian of order $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ as $(-\Delta)^s \varphi = \mathcal{F}^{-1}(|\cdot|^{2s} \hat{\varphi})$ where $\varphi \in \mathcal{S}(\mathbb{R}^n)$ is a Schwartz function. Then $(-\Delta)^s$ extends to a bounded operator $(-\Delta)^s : H^r(\mathbb{R}^n) \rightarrow H^{r-2s}(\mathbb{R}^n)$ for all $r \in \mathbb{R}$ by density of $\mathcal{S}(\mathbb{R}^n)$ in $H^r(\mathbb{R}^n)$.

Let $\Omega \subset \mathbb{R}^n$ be an open set and $F \subset \mathbb{R}^n$ a closed set. We define the following Sobolev spaces

$$\begin{aligned} H_F^r(\mathbb{R}^n) &= \{u \in H^r(\mathbb{R}^n) : \text{spt}(u) \subset F\} \\ \tilde{H}^r(\Omega) &= \text{closure of } C_c^\infty(\Omega) \text{ in } H^r(\mathbb{R}^n) \\ H^r(\Omega) &= \{u|_\Omega : u \in H^r(\mathbb{R}^n)\} \\ H_0^r(\Omega) &= \text{closure of } C_c^\infty(\Omega) \text{ in } H^r(\Omega). \end{aligned}$$

It follows that $\tilde{H}^r(\Omega) \subset H_0^r(\Omega)$, $\tilde{H}^r(\Omega) \subset H_{\bar{\Omega}}^r(\mathbb{R}^n)$, $(\tilde{H}^r(\Omega))^* = H^{-r}(\Omega)$ and $(H^r(\Omega))^* = \tilde{H}^{-r}(\Omega)$ for any open set Ω and $r \in \mathbb{R}$. If Ω is in addition a Lipschitz domain, then we have $\tilde{H}^r(\Omega) = H_{\bar{\Omega}}^r(\mathbb{R}^n)$ for all $r \in \mathbb{R}$ and $H_0^r(\Omega) = H_{\bar{\Omega}}^r(\mathbb{R}^n)$ when $r > -1/2$ such that $r \notin \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\}$.

More generally, let $1 \leq p \leq \infty$ and $r \in \mathbb{R}$. We define the Bessel potential space

$$H^{r,p}(\mathbb{R}^n) = \{u \in \mathcal{S}'(\mathbb{R}^n) : \mathcal{F}^{-1}(\langle \cdot \rangle^r \hat{u}) \in L^p(\mathbb{R}^n)\}$$

equipped with the norm

$$\|u\|_{H^{r,p}(\mathbb{R}^n)} = \|\mathcal{F}^{-1}(\langle \cdot \rangle^r \hat{u})\|_{L^p(\mathbb{R}^n)}.$$

We also write $\mathcal{F}^{-1}(\langle \cdot \rangle^r \hat{u}) =: J^r u$ where the Fourier multiplier $J = (\text{Id} - \Delta)^{1/2}$ is called the Bessel potential. We have the continuous inclusions $H^{r,p}(\mathbb{R}^n) \hookrightarrow H^{t,p}(\mathbb{R}^n)$ whenever $r \geq t$ [41]. By the

Mikhlin multiplier theorem one can show that $(-\Delta)^s: H^{r,p}(\mathbb{R}^n) \rightarrow H^{r-2s,p}(\mathbb{R}^n)$ is continuous whenever $s \geq 0$ and $1 < p < \infty$. The local version of the space $H^{r,p}(\mathbb{R}^n)$ is defined as earlier by the restrictions

$$H^{r,p}(\Omega) = \{u|_{\Omega} : u \in H^{r,p}(\mathbb{R}^n)\}$$

where $\Omega \subset \mathbb{R}^n$ is any open set. This space is equipped with the quotient norm

$$\|v\|_{H^{r,p}(\Omega)} = \inf\{\|w\|_{H^{r,p}(\mathbb{R}^n)} : w \in H^{r,p}(\mathbb{R}^n), w|_{\Omega} = v\}.$$

We have the continuous inclusions $H^{r,p}(\Omega) \hookrightarrow H^{t,p}(\Omega)$ whenever $r \geq t$ by the definition of the quotient norm.

We also define the spaces

$$\begin{aligned} H_F^{r,p}(\mathbb{R}^n) &= \{u \in H^{r,p}(\mathbb{R}^n) : \text{spt}(u) \subset F\} \\ \tilde{H}^{r,p}(\Omega) &= \text{closure of } C_c^\infty(\Omega) \text{ in } H^{r,p}(\mathbb{R}^n) \\ H_0^{r,p}(\Omega) &= \text{closure of } C_c^\infty(\Omega) \text{ in } H^{r,p}(\Omega) \end{aligned}$$

where $F \subset \mathbb{R}^n$ is a closed set. Note that $\tilde{H}^{r,p}(\Omega) \subset H_0^{r,p}(\Omega)$ since the restriction map $|_{\Omega}: H^{r,p}(\mathbb{R}^n) \rightarrow H^{r,p}(\Omega)$ is by definition continuous. One can also see that $\tilde{H}^{r,p}(\Omega) \subset H_{\Omega}^{r,p}(\mathbb{R}^n)$. If Ω is a bounded C^∞ -domain and $1 < p < \infty$, then we have [38, Theorem 1 in section 4.3.2]

$$\begin{aligned} \tilde{H}^{r,p}(\Omega) &= H_{\Omega}^{r,p}(\mathbb{R}^n), \quad s \in \mathbb{R} \\ H_0^{r,p}(\Omega) &= H^{r,p}(\Omega), \quad s \leq \frac{1}{p}. \end{aligned}$$

Some authors (especially in [7, 36]) use the notation $W^{r,p}(\Omega)$ for Bessel potential spaces. We have decided to use the notation $H^{r,p}(\Omega)$ so that these spaces are not confused with the Sobolev-Slobodeckij spaces which are in general different from the Bessel potential spaces [11].

The equation (1) we study is nonlocal. Instead of putting boundary conditions we impose exterior values for the equation. This can be done by saying that $u = f$ in Ω_e if $u - f \in \tilde{H}^s(\Omega)$. Motivated by this we define the (abstract) trace space $X = H^r(\mathbb{R}^n)/\tilde{H}^r(\Omega)$, i.e. functions in X are the same (have the same trace) if they agree in Ω_e . If Ω is a Lipschitz domain, then we have $X = H^r(\Omega_e)$ and $X^* = H_{\Omega_e}^{-r}(\mathbb{R}^n)$.

2.2. Properties of the fractional Laplacian. The fractional Laplacian admits two important properties which we need in our proofs. The first one is unique continuation property (UCP) which is used in proving the Runge approximation property.

Lemma 2.1 (UCP). *Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $r \in \mathbb{R}$ and $u \in H^r(\mathbb{R}^n)$. If $(-\Delta)^s u|_V = 0$ and $u|_V = 0$ for some nonempty open set $V \subset \mathbb{R}^n$, then $u = 0$.*

Lemma 2.1 is proved in [10] for $s > 1$ by reducing the problem to the UCP result for $s \in (0, 1)$ in [15]. Note that such property is not true for local operators like the classical Laplacian $(-\Delta)$. The second property we need is the Poincaré inequality, which is used in showing that the fractional Calderón problem is well-posed.

Lemma 2.2 (Poincaré inequality). *Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $K \subset \mathbb{R}^n$ compact set and $u \in H_K^s(\mathbb{R}^n)$. There exists a constant $c = c(n, K, s) > 0$ such that*

$$\|u\|_{L^2(\mathbb{R}^n)} \leq c \left\| (-\Delta)^{s/2} u \right\|_{L^2(\mathbb{R}^n)}.$$

Many different proofs for lemma 2.2 are given in [10]. We note that in the literature, the fractional Poincaré inequality is typically considered only when $s \in (0, 1)$.

Finally, we recall the fractional Leibniz rule, also known as the Kato-Ponce inequality. It is used to show the boundedness of the bilinear forms associated to the perturbed fractional Schrödinger equation in the case when the coefficients of the PDO have bounded fractional derivatives.

Lemma 2.3 (Kato-Ponce inequality). *Let $s \geq 0$, $1 < r < \infty$, $1 < q_1 \leq \infty$ and $1 < p_2 \leq \infty$ such that $\frac{1}{r} = \frac{1}{p_1} + \frac{1}{q_1} = \frac{1}{p_2} + \frac{1}{q_2}$. If $f \in L^{p_2}(\mathbb{R}^n)$, $J^s f \in L^{p_1}(\mathbb{R}^n)$, $g \in L^{q_1}(\mathbb{R}^n)$ and $J^s g \in L^{q_2}(\mathbb{R}^n)$, then $J^s(fg) \in L^r(\mathbb{R}^n)$ and*

$$\|J^s(fg)\|_{L^r(\mathbb{R}^n)} \leq C(\|J^s f\|_{L^{p_1}(\mathbb{R}^n)} \|g\|_{L^{q_1}(\mathbb{R}^n)} + \|f\|_{L^{p_2}(\mathbb{R}^n)} \|J^s g\|_{L^{q_2}(\mathbb{R}^n)})$$

where J^s is the Bessel potential of order s and $C = C(s, n, r, p_1, p_2, q_1, q_2)$.

The proof of lemma 2.3 can be found in [17] (see also [16, 18]).

2.3. Spaces of rough coefficients. Following [28, Ch. 3], we introduce the space of multipliers $M(H^r \rightarrow H^t)$ between pairs of Sobolev spaces. Here we are assuming that $r, t \in \mathbb{R}$. The coefficients of $P(x, D)$ in theorem 1.1 will be picked from such spaces of multipliers.

If $f \in \mathcal{D}'(\mathbb{R}^n)$ is a distribution, we say that $f \in M(H^r \rightarrow H^t)$ whenever the norm

$$\|f\|_{r,t} := \sup\{|\langle f, uv \rangle| ; u, v \in C_c^\infty(\mathbb{R}^n), \|u\|_{H^r(\mathbb{R}^n)} = \|v\|_{H^{-t}(\mathbb{R}^n)} = 1\}$$

is finite. Here $\langle \cdot, \cdot \rangle$ is the duality pairing. By $M_0(H^r \rightarrow H^t)$ we indicate the closure of $C_c^\infty(\mathbb{R}^n)$ in $M(H^r \rightarrow H^t) \subset \mathcal{D}'(\mathbb{R}^n)$. If $f \in M(H^r \rightarrow H^t)$ and $u, v \in C_c^\infty(\mathbb{R}^n)$ are both non-vanishing, we have the multiplier inequality

(3)

$$|\langle f, uv \rangle| = \left| \left\langle f, \frac{u}{\|u\|_{H^r(\mathbb{R}^n)}} \frac{v}{\|v\|_{H^{-t}(\mathbb{R}^n)}} \right\rangle \right| \|u\|_{H^r(\mathbb{R}^n)} \|v\|_{H^{-t}(\mathbb{R}^n)} \leq \|f\|_{r,t} \|u\|_{H^r(\mathbb{R}^n)} \|v\|_{H^{-t}(\mathbb{R}^n)}.$$

By density (2.3) can be extended to act over $u \in H^r(\mathbb{R}^n), v \in H^{-t}(\mathbb{R}^n)$. Moreover, each $f \in M(H^r \rightarrow H^t)$ gives rise to a multiplication map $m_f : H^r(\mathbb{R}^n) \rightarrow H^t(\mathbb{R}^n)$ defined as

$$\langle m_f(u), v \rangle := \langle f, uv \rangle \quad \text{for all } u \in H^r(\mathbb{R}^n), v \in H^{-t}(\mathbb{R}^n).$$

We have as well the unique adjoint multiplication map $m_f^* : H^{-t}(\mathbb{R}^n) \rightarrow H^{-r}(\mathbb{R}^n)$ such that

$$\langle m_f^*(v), u \rangle := \langle f, uv \rangle \quad \text{for all } u \in H^r(\mathbb{R}^n), v \in H^{-t}(\mathbb{R}^n).$$

Since one sees that the adjoint of m_f is m_f^* , the chosen notation is justified. For convenience, in the rest of the paper we will just write $f u$ for both $m_f(u)$ and $m_f^*(u)$.

Remark 2.4. *The spaces of rough coefficients we use are generalizations of the ones considered in [36]. In fact, the space $Z^{-s}(\mathbb{R}^n)$ used there coincides with our space $M(H^s \rightarrow H^{-s})$.*

In the next lemma we state some elementary properties of the spaces of multipliers. Other interesting properties may be found in [28].

Lemma 2.5. *Let $\lambda, \mu \geq 0$ and $r, t \in \mathbb{R}$. Then*

- (i) $M(H^r \rightarrow H^t) = M(H^{-t} \rightarrow H^{-r})$, and the norms associated to the two spaces also coincide.
- (ii) $M(H^{r-\lambda} \rightarrow H^{t+\mu}) \hookrightarrow M(H^r \rightarrow H^t)$ continuously.
- (iii) $M(H^r \rightarrow H^t) = \{0\}$ whenever $r < t$.

Proof. (i) Let $f \in \mathcal{D}'(\mathbb{R}^n)$ be a distribution. Then by just using the definition we see that

$$\begin{aligned} \|f\|_{r,t} &= \sup\{|\langle f, uv \rangle| ; u, v \in C_c^\infty(\mathbb{R}^n), \|u\|_{H^r(\mathbb{R}^n)} = \|v\|_{H^{-t}(\mathbb{R}^n)} = 1\} \\ &= \sup\{|\langle f, vu \rangle| ; v, u \in C_c^\infty(\mathbb{R}^n), \|v\|_{H^{-t}(\mathbb{R}^n)} = \|u\|_{H^{-(-r)}(\mathbb{R}^n)} = 1\} = \|f\|_{-t, -r}. \end{aligned}$$

(ii) Observe that the given definition of $\|f\|_{r,t}$ is equivalent to the following:

$$\|f\|_{r,t} = \sup\{|\langle f, uv \rangle| ; u, v \in C_c^\infty(\mathbb{R}^n), \|u\|_{H^r(\mathbb{R}^n)} \leq 1, \|v\|_{H^{-t}(\mathbb{R}^n)} \leq 1\}.$$

Since $\lambda, \mu \geq 0$, we also have

$$\|u\|_{H^{r-\lambda}(\mathbb{R}^n)} \leq \|u\|_{H^r(\mathbb{R}^n)}, \quad \|v\|_{H^{-(t+\mu)}(\mathbb{R}^n)} \leq \|v\|_{H^{-t}(\mathbb{R}^n)}.$$

This implies $\|f\|_{r,t} \leq \|f\|_{r-\lambda, t+\mu}$, which in turn gives the wanted inclusion.

(iii) If $0 \leq r < t$, then this was considered in [28, Ch. 3]. The proof given there recalls the easier one for Sobolev spaces ([28, Sec. 2.1]), which is based on the explicit computation of derivatives of aptly chosen exponential functions.

If $r < t \leq 0$, then by point (i) we have $M(H^r \rightarrow H^t) = M(H^{-t} \rightarrow H^{-r})$. We need to show that $M(H^{-t} \rightarrow H^{-r}) = \{0\}$ whenever $0 \leq -t < -r$. This reduces the problem back to the case of non-negative Sobolev scales.

If $r \leq 0 < t$, then $-r \geq 0$. Now by point (ii), we have $M(H^r \rightarrow H^t) \subseteq M(H^{r+(-r)} \rightarrow H^t) = M(L^2 \rightarrow H^t)$. It is therefore enough to show that this last space is trivial, which again immediately follows from the case of non-negative Sobolev scales.

If $r < 0 \leq t$, then the problem can be reduced again to the earlier cases. \square

Remark 2.6. *We also have $M_0(H^{r-\lambda} \rightarrow H^{t+\mu}) \subseteq M_0(H^r \rightarrow H^t)$ whenever $\lambda, \mu \geq 0$, since the inclusion in (ii) is continuous.*

Remark 2.7. *In light of lemma 2.5 (ii) we are only interested in $M(H^r \rightarrow H^t)$ in the case $r \geq t$, the case $r < t$ being trivial. For our theorem 1.1, this translates into the condition $m \leq 2s$. We decided not to consider the limit case $m = 2s$ in this work, as our machinery (in particular, the coercivity estimate (4.1)) breaks down in this case. However, it should be noted that since by assumption we have $m \in \mathbb{Z}$ and $s \notin \mathbb{Z}$, the equality $m = 2s$ can only arise if m is odd, which forces $s = 1/2 + k$ with $k \in \mathbb{Z}$. This case was excluded in [7, 15] as well.*

The next lemmas relate our spaces of multipliers with some special Bessel potential spaces. This is interesting since in the coming section 3 we will consider the inverse problem for coefficients coming from such spaces. We start with a general result.

Lemma 2.8. *Let $\Omega \subset \mathbb{R}^n$ be an open set and let $t \in \mathbb{R}$ and $r \in \mathbb{R}$ be such that $t > \max\{0, r\}$. The following inclusions hold:*

- (i) $\tilde{H}^{r', \infty}(\Omega) \subset M_0(H^{-r} \rightarrow H^{-t})$ whenever $r' \geq \max\{0, r\}$.
- (ii) $H_0^{r', \infty}(\Omega) \subset M_0(H^{-r} \rightarrow H^{-t})$ whenever $r' \geq \max\{0, r\}$ such that $r' \notin \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\}$ and Ω is a Lipschitz domain.
- (iii) $\tilde{H}^{r'}(\Omega) \subset M_0(H^{-r} \rightarrow H^{-t})$ whenever $r' \geq t$ and $r' > n/2$. The same holds true for $H_{\Omega}^{r'}(\mathbb{R}^n)$ if Ω is a Lipschitz domain, and for $H_0^{r'}(\Omega)$ when Ω is a Lipschitz domain and $r' \notin \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\}$.

Proof. Throughout the proof we assume that $u, v \in C_c^\infty(\mathbb{R}^n)$ such that $\|u\|_{H^{-r}(\mathbb{R}^n)} = \|v\|_{H^t(\mathbb{R}^n)} = 1$. In parts (i) and (ii) we can assume that $r' < t$ since if $r' \geq t$, then we have the continuous inclusion $H^{r', \infty}(\Omega) \hookrightarrow H^{r'', \infty}(\Omega)$ where $\max\{0, r\} \leq r'' < t$ (such r'' always exists since $t > \max\{0, r\}$).

(i) Let $f \in \tilde{H}^{r', \infty}(\Omega)$. Now $f = f_1 + f_2$ where $f_1 \in C_c^\infty(\Omega)$ and $\|f_2\|_{H^{r', \infty}(\mathbb{R}^n)} \leq \epsilon$. Then

$$\begin{aligned} |\langle f_2, uv \rangle| &\leq \|f_2 v\|_{H^{r'}(\mathbb{R}^n)} \|u\|_{H^{-r'}(\mathbb{R}^n)} \leq C \|f_2\|_{H^{r', \infty}(\mathbb{R}^n)} \|v\|_{H^{r'}(\mathbb{R}^n)} \|u\|_{H^{-r}(\mathbb{R}^n)} \\ &\leq C \epsilon \|v\|_{H^t(\mathbb{R}^n)} = C \epsilon. \end{aligned}$$

Here we used the Kato-Ponce inequality (lemma 2.3)

$$\begin{aligned} \|J^{r'}(f_2 v)\|_{L^2(\mathbb{R}^n)} &\leq C(\|f_2\|_{L^\infty(\mathbb{R}^n)} \|J^{r'} v\|_{L^2(\mathbb{R}^n)} + \|J^{r'} f_2\|_{L^\infty(\mathbb{R}^n)} \|v\|_{L^2(\mathbb{R}^n)}) \\ &\leq C \|f_2\|_{H^{r', \infty}(\mathbb{R}^n)} \|v\|_{H^{r'}(\mathbb{R}^n)} \end{aligned}$$

and the assumption $\max\{0, r\} \leq r' < t$. Therefore $\|f - f_1\|_{-r, -t} = \|f_2\|_{-r, -t} \leq C \epsilon$ which shows that $f \in M_0(H^{-r} \rightarrow H^{-t})$.

(ii) Let $f \in H_0^{r', \infty}(\Omega)$. Now $f = f_1 + f_2$ where $f_1 \in C_c^\infty(\Omega)$ and $\|f_2\|_{H^{r', \infty}(\Omega)} \leq \epsilon$. By the definition of the quotient norm $\|\cdot\|_{H^{r', \infty}(\Omega)}$ we can take $F \in H^{r', \infty}(\mathbb{R}^n)$ such that $F|_{\Omega} = f_2$ and $\|F\|_{H^{r', \infty}(\mathbb{R}^n)} \leq 2 \|f_2\|_{H^{r', \infty}(\Omega)}$. The assumptions imply the duality $(H^{-r'}(\Omega))^* = H_0^{r'}(\Omega) \subset$

$H^{r'}(\Omega)$. Using the Kato-Ponce inequality for the extension F we obtain as in the proof of part (i) that

$$\left\| J^{r'}(Fv) \right\|_{L^2(\mathbb{R}^n)} \leq C \|F\|_{H^{r',\infty}(\mathbb{R}^n)} \|v\|_{H^{r'}(\mathbb{R}^n)} \leq 2C \|f_2\|_{H^{r',\infty}(\Omega)} \|v\|_{H^t(\mathbb{R}^n)} \leq 2C\epsilon$$

and hence

$$\begin{aligned} |\langle f_2, uv \rangle| &\leq \|f_2 v\|_{(H^{-r'}(\Omega))^*} \|u\|_{H^{-r'}(\Omega)} \leq \|f_2 v\|_{H^{r'}(\Omega)} \|u\|_{H^{-r}(\mathbb{R}^n)} \\ &\leq \left\| J^{r'}(Fv) \right\|_{L^2(\mathbb{R}^n)} \leq 2C\epsilon. \end{aligned}$$

This shows that $f \in M_0(H^{-r} \rightarrow H^{-t})$.

(iii) Let $f \in \tilde{H}^{r'}(\Omega)$. Now $f = f_1 + f_2$ where $f_1 \in C_c^\infty(\Omega)$ and $\|f_2\|_{H^{r'}(\mathbb{R}^n)} \leq \epsilon$. Now [3, Theorem 7.3] implies the continuity of the multiplication $H^{r'}(\mathbb{R}^n) \times H^t(\mathbb{R}^n) \hookrightarrow H^t(\mathbb{R}^n)$ when $r' \geq t$ and $r' > n/2$. We obtain

$$|\langle f_2, uv \rangle| \leq \|f_2 v\|_{H^t(\mathbb{R}^n)} \|u\|_{H^{-t}(\mathbb{R}^n)} \leq C \|f_2\|_{H^{r'}(\mathbb{R}^n)} \|v\|_{H^t(\mathbb{R}^n)} \|u\|_{H^{-r}(\mathbb{R}^n)} \leq C\epsilon.$$

Hence $f \in M_0(H^{-r} \rightarrow H^{-t})$. If Ω is a Lipschitz domain, then $H_{\Omega}^{r'}(\mathbb{R}^n) = \tilde{H}^{r'}(\Omega)$. If in addition $r' \notin \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\}$, we also have $H_0^{r'}(\Omega) = \tilde{H}^{r'}(\Omega)$. \square

Note that the assumptions in theorem 1.1 satisfy the conditions of the previous lemma since then $r = |\alpha| - s$ and $t = s$. The following lemma gives examples of spaces of lower order coefficients ($|\alpha| \leq s$).

Lemma 2.9. *Let $\Omega \subset \mathbb{R}^n$ be an open set and $t > 0$. The following inclusions hold:*

- (i) $L^p(\Omega) \subset M_0(H^0 \rightarrow H^{-t})$ whenever $2 \leq p < \infty$ and $p > n/t$. Especially, if Ω is bounded, then $L^\infty(\Omega) \subset M_0(H^0 \rightarrow H^{-t})$.
- (ii) $\tilde{H}^r(\Omega) \subset M_0(H^0 \rightarrow H^{-t})$ whenever $r \geq 0$ and $r > n/2 - t$. The same holds true for $H_{\Omega}^r(\mathbb{R}^n)$ if Ω is a Lipschitz domain, and for $H_0^r(\Omega)$ when Ω is Lipschitz domain and $r \notin \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\}$.

Proof. Throughout the proof we assume that $u, v \in C_c^\infty(\mathbb{R}^n)$ such that $\|u\|_{L^2(\mathbb{R}^n)} = \|v\|_{H^t(\mathbb{R}^n)} = 1$.

(i) Let $f \in L^p(\Omega)$. By density of $C_c^\infty(\Omega)$ in $L^p(\Omega)$ we have $f = f_1 + f_2$ where $f_1 \in C_c^\infty(\Omega)$ and $\left\| \tilde{f}_2 \right\|_{L^p(\mathbb{R}^n)} \leq \epsilon$ where \tilde{f}_2 is the zero extension of $f_2 \in L^p(\Omega)$. The assumptions on p imply the continuity of the multiplication $L^p(\mathbb{R}^n) \times H^t(\mathbb{R}^n) \hookrightarrow L^2(\mathbb{R}^n)$ ([3, Theorem 7.3]) and we have

$$\left| \langle \tilde{f}_2, uv \rangle \right| \leq \left\| \tilde{f}_2 v \right\|_{L^2(\mathbb{R}^n)} \|u\|_{L^2(\mathbb{R}^n)} \leq C \left\| \tilde{f}_2 \right\|_{L^p(\mathbb{R}^n)} \|v\|_{H^t(\mathbb{R}^n)} \leq C\epsilon.$$

This gives that $f \in M_0(H^0 \rightarrow H^{-t})$. If Ω is bounded, we have $L^\infty(\Omega) \hookrightarrow L^p(\Omega)$ for all $1 \leq p < \infty$, giving the second claim.

(ii) Let $f \in \tilde{H}^r(\Omega)$. Now we have $f = f_1 + f_2$ where $f_1 \in C_c^\infty(\Omega)$ and $\|f_2\|_{H^r(\mathbb{R}^n)} \leq \epsilon$. The assumptions on r imply that the multiplication $H^r(\mathbb{R}^n) \times H^t(\mathbb{R}^n) \hookrightarrow L^2(\mathbb{R}^n)$ is continuous ([3, Theorem 7.3]). We obtain

$$|\langle f_2, uv \rangle| \leq \|f_2 v\|_{L^2(\mathbb{R}^n)} \|u\|_{L^2(\mathbb{R}^n)} \leq C \|f_2\|_{H^r(\mathbb{R}^n)} \|v\|_{H^t(\mathbb{R}^n)} \leq C\epsilon$$

and therefore $f \in M_0(H^0 \rightarrow H^{-t})$. The claims for $H_{\Omega}^r(\mathbb{R}^n)$ and $H_0^r(\Omega)$ follow as in the proof of part (iii) of lemma 2.8 from the usual identifications for Lipschitz domains. \square

As mentioned above we put $t = s > 0$ in theorem 1.1 and the condition in lemma 2.9 is satisfied. Note that under the assumption $|\alpha| \leq s$ we have $M_0(H^0 \rightarrow H^{-s}) \subset M_0(H^{s-|\alpha|} \rightarrow H^{-s})$. Hence we can choose the lower order coefficients from a less regular space in theorem 1.1 (compare to lemma 2.8).

3. MAIN THEOREM FOR SINGULAR COEFFICIENTS

In this section, to shorten the notation, we will write $\|\cdot\|_{H^s}$, $\|\cdot\|_{L^2}$ and so on for the global norms in \mathbb{R}^n when the base set is not written explicitly.

3.1. Well-posedness of the inverse problem. Consider the problem

$$(4) \quad \begin{aligned} (-\Delta)^s u + \sum_{|\alpha| \leq m} a_\alpha (D^\alpha u) &= F \quad \text{in } \Omega, \\ u &= f \quad \text{in } \Omega_e \end{aligned}$$

and the corresponding *adjoint-problem*

$$(5) \quad \begin{aligned} (-\Delta)^s u^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha u^*) &= F^* \quad \text{in } \Omega, \\ u^* &= f^* \quad \text{in } \Omega_e. \end{aligned}$$

Note that if $u, u^* \in H^s(\mathbb{R}^n)$ and $a_\alpha \in M(H^{s-|\alpha|} \rightarrow H^{-s}) = M(H^s \rightarrow H^{|\alpha|-s})$, then $a_\alpha (D^\alpha u) \in H^{-s}(\mathbb{R}^n)$ and $D^\alpha (a_\alpha u^*) \in H^{-s}(\mathbb{R}^n)$ matching with $(-\Delta)^s u, (-\Delta)^s u^* \in H^{-s}(\mathbb{R}^n)$.

The problems (3.1) and (3.1) are associated to the bilinear forms

$$(6) \quad B_P(v, w) := \langle (-\Delta)^{s/2} v, (-\Delta)^{s/2} w \rangle_{\mathbb{R}^n} + \sum_{|\alpha| \leq m} \langle a_\alpha, (D^\alpha v) w \rangle_{\mathbb{R}^n}$$

and

$$(7) \quad B_P^*(v, w) := \langle (-\Delta)^{s/2} v, (-\Delta)^{s/2} w \rangle_{\mathbb{R}^n} + \sum_{|\alpha| \leq m} \langle a_\alpha, v (D^\alpha w) \rangle_{\mathbb{R}^n},$$

defined on $v, w \in C_c^\infty(\mathbb{R}^n)$. In the latter terms of the bilinear forms we have written the dual pairing as $\langle \cdot, \cdot \rangle_{\mathbb{R}^n}$ since a_α is now a distribution in the whole space \mathbb{R}^n in contrast to section 4 where a_α is an object defined only in Ω .

Remark 3.1. *Observe that B_P is not symmetric, which motivates the introduction of the bilinear form B_P^* . Moreover, one sees by simple inspection that $B_P(v, w) = B_P^*(w, v)$ for all $v, w \in C_c^\infty(\mathbb{R}^n)$. This identity holds for $v, w \in H^s(\mathbb{R}^n)$ as well by density, thanks to the following boundedness lemma.*

Lemma 3.2 (Boundedness of the bilinear forms). *Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $m \in \mathbb{N}$ such that $2s \geq m$, and let $a_\alpha \in M(H^{s-|\alpha|} \rightarrow H^{-s})$. Then B_P and B_P^* extend as bounded bilinear forms on $H^s(\mathbb{R}^n) \times H^s(\mathbb{R}^n)$.*

Proof of lemma 3.2. We only prove the boundedness of B_P , as for B_P^* one can proceed in the same way. The proof is a simple calculation following from inequality (2.3). Let $u, v \in C_c^\infty(\mathbb{R}^n)$. We can then estimate that

$$\begin{aligned} |B_P(v, w)| &\leq |\langle (-\Delta)^{s/2} v, (-\Delta)^{s/2} w \rangle_{\mathbb{R}^n}| + \sum_{|\alpha| \leq m} |\langle a_\alpha, D^\alpha v w \rangle_{\mathbb{R}^n}| \\ &\leq \|w\|_{H^s(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)} + \sum_{|\alpha| \leq m} \|a_\alpha\|_{s-|\alpha|, -s} \|D^\alpha v\|_{H^{s-|\alpha|}(\mathbb{R}^n)} \|w\|_{H^s(\mathbb{R}^n)} \\ &\leq \left(1 + \sum_{|\alpha| \leq m} \|a_\alpha\|_{s-|\alpha|, -s} \right) \|w\|_{H^s(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)}. \end{aligned}$$

Now the claim follows from the density of $C_c^\infty(\mathbb{R}^n)$ in $H^s(\mathbb{R}^n)$. □

Next we shall prove existence and uniqueness of solutions for the problems (3.1) and (3.1). To this end, we will use the following form of Young's inequality, which holds for all $a, b, \eta \in \mathbb{R}^+$

and $p, q \in (1, \infty)$ such that $1/p + 1/q = 1$:

$$(8) \quad ab \leq \frac{(q\eta)^{-p/q}}{p} a^p + \eta b^q.$$

The validity of (3.1) is easily proved by choosing $a_1 = a(q\eta)^{-1/q}$ and $b_1 = b(q\eta)^{1/q}$ in Young's inequality $a_1 b_1 \leq a_1^p/p + b_1^q/q$.

Lemma 3.3 (Well-posedness). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set. Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $m \in \mathbb{N}$ be such that $2s > m$, and let $a_\alpha \in M_0(H^{s-|\alpha|} \rightarrow H^{-s})$. There exist a real number $\mu > 0$ and a countable set $\Sigma \subset (-\mu, \infty)$ of eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots \rightarrow \infty$ such that if $\lambda \in \mathbb{R} \setminus \Sigma$, for any $f \in H^s(\mathbb{R}^n)$ and $F \in (\tilde{H}^s(\Omega))^*$ there exists unique $u \in H^s(\mathbb{R}^n)$ such that $u - f \in \tilde{H}^s(\Omega)$ and*

$$B_P(u, v) - \lambda \langle u, v \rangle_\Omega = F(v) \quad \text{for all } v \in \tilde{H}^s(\Omega).$$

One has the estimate

$$\|u\|_{H^s(\mathbb{R}^n)} \leq C \left(\|f\|_{H^s(\mathbb{R}^n)} + \|F\|_{(\tilde{H}^s(\Omega))^*} \right).$$

The function u is also the unique $u \in H^s(\mathbb{R}^n)$ satisfying

$$r_\Omega \left((-\Delta)^s + \sum_{|\alpha| \leq m} a_\alpha D^\alpha - \lambda \right) u = F$$

in the sense of distributions in Ω and $u - f \in \tilde{H}^s(\Omega)$. Moreover, if (3.1) holds then $0 \notin \Sigma$.

Proof. Let $\tilde{u} := u - f$. The above problem is reduced to finding a unique $\tilde{u} \in \tilde{H}^s(\Omega)$ such that $B_P(\tilde{u}, v) - \lambda \langle \tilde{u}, v \rangle_\Omega = \tilde{F}(v)$, where $\tilde{F} := F - B_P(f, \cdot) + \lambda \langle f, \cdot \rangle_\Omega$. Observe that the modified functional \tilde{F} belongs to $(\tilde{H}^s(\Omega))^*$ as well, since by lemma 3.2 we have for all $v \in \tilde{H}^s(\Omega)$

$$|\tilde{F}(v)| \leq |F(v)| + |B_P(f, v)| + |\lambda| |\langle f, v \rangle_\Omega| \leq (\|F\|_{(\tilde{H}^s(\Omega))^*} + (C + |\lambda|) \|f\|_{H^s(\mathbb{R}^n)}) \|v\|_{H^s(\mathbb{R}^n)}.$$

Since $a_\alpha \in M_0(H^{s-|\alpha|} \rightarrow H^{-s})$, for any $\epsilon > 0$ we can write $a_\alpha = a_{\alpha,1} + a_{\alpha,2}$, where $a_{\alpha,1} \in C_c^\infty(\mathbb{R}^n) \cap M(H^{s-|\alpha|} \rightarrow H^{-s})$ and $\|a_{\alpha,2}\|_{s-|\alpha|, -s} < \epsilon$. Thus by formula (2.3), the continuity of the multiplication $H^r(\mathbb{R}^n) \times H^s(\mathbb{R}^n) \hookrightarrow H^s(\mathbb{R}^n)$ for large enough $r \in \mathbb{R}$ (see [3, Theorem 7.3]) and the fact that $a_{\alpha,1} \in C_c^\infty(\mathbb{R}^n) \subset H^r(\mathbb{R}^n)$ for all $r \in \mathbb{R}$ we obtain

(9)

$$\begin{aligned} |\langle a_\alpha, D^\alpha v w \rangle| &\leq |\langle a_{\alpha,1}, D^\alpha v w \rangle| + |\langle a_{\alpha,2}, D^\alpha v w \rangle| \\ &\leq \|a_{\alpha,1}\|_{H^r(\mathbb{R}^n)} \|D^\alpha v\|_{H^{-s}(\mathbb{R}^n)} \|w\|_{H^s(\mathbb{R}^n)} + \|a_{\alpha,2}\|_{s-|\alpha|, -s} \|D^\alpha v\|_{H^{s-|\alpha|}(\mathbb{R}^n)} \|w\|_{H^s(\mathbb{R}^n)} \\ &\leq c \|w\|_{H^s(\mathbb{R}^n)} \left(\|a_{\alpha,1}\|_{H^r(\mathbb{R}^n)} \|v\|_{H^{|\alpha|-s}(\mathbb{R}^n)} + \epsilon \|v\|_{H^s(\mathbb{R}^n)} \right) \end{aligned}$$

where $r \in \mathbb{R}$ is large enough ($r > \max\{s, n/2\}$ is sufficient). If $|\alpha| < s$, from formulas (3.1) and (3.1) with $p = q = 2$ we get directly

$$(10) \quad \begin{aligned} |\langle a_\alpha, D^\alpha v w \rangle| &\leq C \left(\|v\|_{H^s(\mathbb{R}^n)} \|v\|_{L^2(\mathbb{R}^n)} + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\leq C(\epsilon^{-1} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2) \end{aligned}$$

for a constant C independent of v, w, ϵ . If instead $|\alpha| > s$ (observe that we can not have $|\alpha| = s$, because s can not be an integer), we use the interpolation inequality

$$\|v\|_{H^{|\alpha|-s}(\mathbb{R}^n)} \leq C \|v\|_{L^2(\mathbb{R}^n)}^{1-(|\alpha|-s)/s} \|v\|_{H^s(\mathbb{R}^n)}^{(|\alpha|-s)/s} = C \|v\|_{L^2(\mathbb{R}^n)}^{2-|\alpha|/s} \|v\|_{H^s(\mathbb{R}^n)}^{|\alpha|/s-1}$$

in order to get

$$|\langle a_\alpha, D^\alpha v w \rangle| \leq C \|w\|_{H^s(\mathbb{R}^n)} \left(\|v\|_{L^2(\mathbb{R}^n)}^{2-|\alpha|/s} \|v\|_{H^s(\mathbb{R}^n)}^{|\alpha|/s-1} + \epsilon \|v\|_{H^s(\mathbb{R}^n)} \right).$$

Then by formula (3.1) with

$$a = \|v\|_{L^2(\mathbb{R}^n)}^{2-|\alpha|/s}, \quad b = \|v\|_{H^s(\mathbb{R}^n)}^{|\alpha|/s-1}, \quad p = \frac{s}{2s-|\alpha|}, \quad q = \frac{s}{|\alpha|-s}, \quad \eta = \epsilon$$

we obtain

$$|\langle a_\alpha, D^\alpha v w \rangle| \leq C \|w\|_{H^s(\mathbb{R}^n)} \left(\epsilon^{\frac{s-|\alpha|}{2s-|\alpha|}} \|v\|_{L^2(\mathbb{R}^n)} + \epsilon \|v\|_{H^s(\mathbb{R}^n)} \right)$$

for a constant C independent of v, w, ϵ . Now we use formula (3.1) again, but this time we choose

$$a = \|v\|_{L^2(\mathbb{R}^n)}, \quad b = \|v\|_{H^s(\mathbb{R}^n)}, \quad q = p = 2, \quad \eta = \epsilon^{s/(2s-|\alpha|)}.$$

This leads to

$$\begin{aligned} (11) \quad |\langle a_\alpha, D^\alpha v v \rangle| &\leq C \left(\epsilon^{\frac{s-|\alpha|}{2s-|\alpha|}} \|v\|_{L^2(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)} + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\leq C \left(\epsilon^{\frac{-|\alpha|}{2s-|\alpha|}} \|v\|_{L^2(\mathbb{R}^n)}^2 + 2\epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\leq C \left(\epsilon^{\frac{-|\alpha|}{2s-|\alpha|}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\leq C' \left(\epsilon^{\frac{-m}{2s-m}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \end{aligned}$$

where C, C' are constants changing from line to line. Observe that C' can be taken independent of α . Eventually, using (3.1) and (3.1) we get

$$\begin{aligned} (12) \quad B_P(v, v) &\geq \|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 - \sum_{|\alpha| \leq m} |\langle a_\alpha, D^\alpha v v \rangle| \\ &\geq \|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 - C' \left((\epsilon^{\frac{-m}{2s-m}} + \epsilon^{-1}) \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right). \end{aligned}$$

By the higher order Poincaré inequality (lemma 2.2) (4.1) turns into

$$\begin{aligned} B_P(v, v) &\geq c \left(\|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 + \|v\|_{L^2(\mathbb{R}^n)}^2 \right) - C' \left((\epsilon^{\frac{-m}{2s-m}} + \epsilon^{-1}) \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\geq c \|v\|_{H^s(\mathbb{R}^n)}^2 - C' \left((\epsilon^{\frac{-m}{2s-m}} + \epsilon^{-1}) \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \end{aligned}$$

for some constant $c = c(\Omega, n, s)$ changing from line to line. For ϵ small enough, this eventually gives the coercivity estimate

$$(13) \quad B_P(v, v) \geq c_0 \|v\|_{H^s(\mathbb{R}^n)}^2 - \mu \|v\|_{L^2(\mathbb{R}^n)}^2$$

for some constants $c_0, \mu > 0$ independent of v .

As a consequence of the coercivity estimate, $B_P(\cdot, \cdot) + \mu \langle \cdot, \cdot \rangle_{L^2(\mathbb{R}^n)}$ is an equivalent inner product on $\tilde{H}^s(\Omega)$. Therefore, by the Riesz representation theorem there exists a bounded linear operator $G_\mu : (\tilde{H}^s(\Omega))^* \rightarrow \tilde{H}^s(\Omega)$ associating each functional in $(\tilde{H}^s(\Omega))^*$ to its unique representative in the inner product $B_P(\cdot, \cdot) + \mu \langle \cdot, \cdot \rangle_{L^2(\mathbb{R}^n)}$ on $\tilde{H}^s(\Omega)$. Thus $\tilde{u} := G_\mu \tilde{F}$ verifies

$$B_P(\tilde{u}, v) + \mu \langle \tilde{u}, v \rangle_{L^2(\mathbb{R}^n)} = \tilde{F}(v) \quad \text{for all } v \in \tilde{H}^s(\Omega)$$

and it is the required unique solution $\tilde{u} \in \tilde{H}^s(\Omega)$. Moreover, G_μ induces a compact, self-adjoint and positive operator $\tilde{G}_\mu : L^2(\Omega) \rightarrow L^2(\Omega)$ by the compact Sobolev embedding theorem. The remaining claims follow from the spectral theorem for \tilde{G}_μ and from the Fredholm alternative as in [15]. \square

By the above lemma 3.3, both problems (3.1) and (3.1) have a countable set of Dirichlet eigenvalues. Throughout the paper we will assume that the coefficients a_α are such that 0 is

not a Dirichlet eigenvalue for either of the problems. That is, we assume that

$$(14) \quad \begin{cases} \text{if } u \in H^s(\mathbb{R}^n) \text{ solves } (-\Delta)^s u + \sum_{|\alpha| \leq m} a_\alpha D^\alpha u = 0 \text{ in } \Omega \text{ and } u|_{\Omega_e} = 0, \\ \text{then } u \equiv 0 \end{cases}$$

and

$$(15) \quad \begin{cases} \text{if } u^* \in H^s(\mathbb{R}^n) \text{ solves } (-\Delta)^s u^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha u^*) = 0 \text{ in } \Omega \text{ and } u^*|_{\Omega_e} = 0, \\ \text{then } u^* \equiv 0. \end{cases}$$

With this in mind, we shall define the DN maps. Consider the abstract trace space $X := H^s(\mathbb{R}^n)/\tilde{H}^s(\Omega)$ equipped with the quotient norm

$$\| [f] \|_X := \inf_{\phi \in \tilde{H}^s(\Omega)} \| f - \phi \|_{H^s(\mathbb{R}^n)}, \quad f \in H^s(\mathbb{R}^n)$$

and its dual space X^* . We use these in order to define the DN maps associated to the problems (3.1) and (3.1), which we study in the following lemma.

Lemma 3.4 (DN maps). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set. Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $m \in \mathbb{N}$ such that $2s > m$, and let $a_\alpha \in M_0(H^{s-|\alpha|} \rightarrow H^{-s})$. There exist two continuous linear maps*

$$\Lambda_P : X \rightarrow X^* \quad \text{defined by} \quad \langle \Lambda_P[f], [g] \rangle := B_P(u_f, g)$$

and

$$\Lambda_P^* : X \rightarrow X^* \quad \text{defined by} \quad \langle \Lambda_P^*[f], [g] \rangle := B_P^*(u_f^*, g)$$

where u_f, u_f^* are the unique solutions to the equations

$$(-\Delta)^s u + \sum_{|\alpha| \leq m} a_\alpha D^\alpha u = 0 \quad \text{in } \Omega, \quad u - f \in \tilde{H}^s(\Omega)$$

and

$$(-\Delta)^s u^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha u^*) = 0 \quad \text{in } \Omega, \quad u^* - f \in \tilde{H}^s(\Omega)$$

with $f, g \in H^s(\mathbb{R}^n)$. Moreover, the identity $\langle \Lambda_P[f], [g] \rangle = \langle [f], \Lambda_P^*[g] \rangle$ holds.

Proof. We show well-definedness and continuity only for Λ_P , the proof being similar for Λ_P^* . We note that such unique solutions exist by lemma 3.3.

If $\phi \in \tilde{H}^s(\Omega)$, then $u_f|_{\Omega_e} = f = u_{f+\phi}|_{\Omega_e}$, and also $u_f, u_{f+\phi}$ both solve $(-\Delta)^s u + Pu = 0$ in Ω . By unicity of solutions, we must then have that u_f and $u_{f+\phi}$ coincide. On the other hand, if $\psi \in \tilde{H}^s(\Omega)$, then $\psi|_{\Omega_e} = 0$. These two facts imply the well-definedness of Λ_P , since

$$B_P(u_{f+\phi}, g + \psi) = B_P(u_f, g) + B_P(u_f, \psi) = B_P(u_f, g).$$

The continuity of Λ_P is an easy consequence of lemma 3.2 and the estimate in lemma 3.3. If $f, g \in H^s(\mathbb{R}^n)$ and $\phi, \psi \in \tilde{H}^s(\Omega)$, then

$$|\langle \Lambda_P[f], [g] \rangle| = |B_P(u_{f-\phi}, g - \psi)| \leq C \|u_{f-\phi}\|_{H^s} \|g - \psi\|_{H^s} \leq C \|f - \phi\|_{H^s} \|g - \psi\|_{H^s}.$$

By taking the infimum on both sides with respect to ϕ and ψ , we end up with

$$|\langle \Lambda_P[f], [g] \rangle| \leq C \inf_{\phi \in \tilde{H}^s(\Omega)} \|f - \phi\|_{H^s} \inf_{\psi \in \tilde{H}^s(\Omega)} \|g - \psi\|_{H^s} = C \| [f] \|_X \| [g] \|_X.$$

The well-posedness result proved above implies that for all $f, g \in H^s(\mathbb{R}^n)$ we have $\langle \Lambda_P[f], [g] \rangle = B_P(u_f, e_g)$, where e_g is a generic extension of $g|_{\Omega_e}$ from Ω_e to \mathbb{R}^n . In particular, $\langle \Lambda_P[f], [g] \rangle = B_P(u_f, u_g^*)$. By lemma 3.2 this leads to

$$\langle \Lambda_P[f], [g] \rangle = B_P(u_f, u_g^*) = B_P^*(u_g^*, u_f) = \langle \Lambda_P^*[g], [f] \rangle$$

which concludes the proof. \square

Remark 3.5. *We should observe at this point that a priori Λ_P^* has no reason to be the adjoint of Λ_P , as the symbols would suggest. However, the identity we proved in lemma 3.4 shows that this is in fact true, and thus there is no abuse of notation.*

3.2. Proof of injectivity. The proof of injectivity is based on an Alessandrini identity and the Runge approximation property for our operator, following the scheme developed in [15].

Lemma 3.6 (Alessandrini identity). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set. Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $m \in \mathbb{N}$ such that $2s > m$. For $j = 1, 2$, let $a_{j,\alpha} \in M_0(H^{s-|\alpha|} \rightarrow H^{-s})$. For any $f_1, f_2 \in H^s(\mathbb{R}^n)$, let $u_1, u_2^* \in H^s(\mathbb{R}^n)$ respectively solve*

$$(-\Delta)^s u_1 + \sum_{|\alpha| \leq m} a_{1,\alpha} D^\alpha u_1 = 0 \quad \text{in } \Omega, \quad u_1 - f_1 \in \tilde{H}^s(\Omega)$$

and

$$(-\Delta)^s u_2^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_{2,\alpha} u_2^*) = 0 \quad \text{in } \Omega, \quad u_2^* - f_2 \in \tilde{H}^s(\Omega).$$

Then we have the integral identity

$$\langle (\Lambda_{P_1} - \Lambda_{P_2})[f_1], [f_2] \rangle = \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha u_1 u_2^* \rangle.$$

Proof. The proof is a simple computation following from lemma 3.4

$$\begin{aligned} \langle (\Lambda_{P_1} - \Lambda_{P_2})[f_1], [f_2] \rangle &= \langle \Lambda_{P_1}[f_1], [f_2] \rangle - \langle \Lambda_{P_2}[f_1], [f_2] \rangle = \langle \Lambda_{P_1}[f_1], [f_2] \rangle - \langle [f_1], \Lambda_{P_2}^*[f_2] \rangle \\ &= B_{P_1}(u_1, u_2^*) - B_{P_2}^*(u_2^*, u_1) = \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha u_1 u_2^* \rangle. \quad \square \end{aligned}$$

Lemma 3.7 (Runge approximation property). *Let $\Omega, W \subset \mathbb{R}^n$ respectively be a bounded open set and a non-empty open set such that $\bar{W} \cap \bar{\Omega} = \emptyset$. Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $m \in \mathbb{N}$ be such that $2s > m$, and let $a_\alpha \in M_0(H^{s-|\alpha|} \rightarrow H^{-s})$. Moreover, let $\mathcal{R} := \{u_f - f : f \in C_c^\infty(W)\} \subset \tilde{H}^s(\Omega)$ where u_f solves*

$$(-\Delta)^s u_f + \sum_{|\alpha| \leq m} a_\alpha D^\alpha u_f = 0 \quad \text{in } \Omega, \quad u_f - f \in \tilde{H}^s(\Omega)$$

and $\mathcal{R}^* := \{u_f^* - f : f \in C_c^\infty(W)\} \subset \tilde{H}^s(\Omega)$ where u_f^* solves

$$(-\Delta)^s u_f^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha u_f^*) = 0 \quad \text{in } \Omega, \quad u_f^* - f \in \tilde{H}^s(\Omega).$$

Then \mathcal{R} and \mathcal{R}^* are dense in $\tilde{H}^s(\Omega)$.

Proof. The proofs of the two statements are similar, so we show only the density of \mathcal{R} in $\tilde{H}^s(\Omega)$. By the Hahn-Banach theorem, it is enough to prove that any functional F acting on $\tilde{H}^s(\Omega)$ that vanishes on \mathcal{R} must be identically 0. Thus, let $F \in (\tilde{H}^s(\Omega))^*$ and assume $F(u_f - f) = 0$ for all $f \in C_c^\infty(W)$. Let ϕ be the unique solution of

$$(16) \quad (-\Delta)^s \phi + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha \phi) = -F \quad \text{in } \Omega, \quad \phi \in \tilde{H}^s(\Omega).$$

In other words, ϕ is the unique function in $\tilde{H}^s(\Omega)$ such that $B_P^*(\phi, w) = -F(w)$ for all $w \in \tilde{H}^s(\Omega)$. Then we can compute

$$\begin{aligned} (17) \quad 0 &= F(u_f - f) = -B_P^*(\phi, u_f - f) = B_P^*(\phi, f) \\ &= \langle (-\Delta)^{s/2} f, (-\Delta)^{s/2} \phi \rangle + \sum_{|\alpha| \leq m} \langle a_\alpha, D^\alpha f \phi \rangle \\ &= \langle f, (-\Delta)^s \phi \rangle. \end{aligned}$$

On the first line of (3.2) we used that $\phi \in \tilde{H}^s(\Omega)$ and u_f solves the equation in Ω , and on the last line we used the support condition for f . By the arbitrariness of $f \in C_c^\infty(W)$ we have obtained that $(-\Delta)^s \phi = 0$ in W , and on the same set we also have $\phi = 0$. Using the unique continuation

result for the higher order fractional Laplacian given in lemma 2.1 we deduce $\phi \equiv 0$ on all of \mathbb{R}^n . The vanishing of the functional F now follows easily from the definition of ϕ . \square

Remark 3.8. We remark that using the same proof one can show that $r_\Omega \mathcal{R} \subset L^2(\Omega)$ and $r_\Omega \mathcal{R}^* \subset L^2(\Omega)$ are dense in $L^2(\Omega)$, where r_Ω is the restriction to Ω . If $F \in L^2(\Omega)$, then F induces an element in $(\tilde{H}^s(\Omega))^*$ via the inner product $F(w) := \langle F, r_\Omega w \rangle_\Omega$, where $w \in \tilde{H}^s(\Omega)$. Hence one can choose the solution ϕ in equation (3.2) with F as a source term and complete the proof as in equation (3.2) showing that $(r_\Omega \mathcal{R})^\perp = \{0\}$ in $L^2(\Omega)$ (similarly $(r_\Omega \mathcal{R}^*)^\perp = \{0\}$).

We are ready to prove the main result of the paper.

Proof of theorem 1.1. Step 1. Since one can always shrink the sets W_1 and W_2 if necessary, we can assume without loss of generality that $\bar{W}_1 \cap \bar{W}_2 = \emptyset$. Let $v_1, v_2 \in C_c^\infty(\Omega)$. By the Runge approximation property proved in lemma 3.7 we can find two sequences of functions $\{f_{j,k}\}_{k \in \mathbb{N}} \subset C_c^\infty(W_j)$, $j = 1, 2$, such that

$$u_{1,k} = f_{1,k} + v_1 + r_{1,k}, \quad u_{2,k}^* = f_{2,k} + v_2 + r_{2,k}$$

where $u_{1,k}, u_{2,k}^* \in \tilde{H}^s(\Omega)$ respectively solve

$$(-\Delta)^s u_{1,k} + \sum_{|\alpha| \leq m} a_{1,\alpha} D^\alpha u_{1,k} = 0 \quad \text{in } \Omega, \quad u_{1,k} - f_{1,k} \in \tilde{H}^s(\Omega)$$

and

$$(-\Delta)^s u_{2,k}^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_{2,\alpha} u_{2,k}^*) = 0 \quad \text{in } \Omega, \quad u_{2,k}^* - f_{2,k} \in \tilde{H}^s(\Omega)$$

and $r_{1,k}, r_{2,k} \rightarrow 0$ in $\tilde{H}^s(\Omega)$ as $k \rightarrow \infty$. By the assumption on the DN maps and the Alessandrini identity from lemma 3.6 we have

$$\begin{aligned} (18) \quad 0 &= \langle (\Lambda_{P_1} - \Lambda_{P_2})[f_{1,k}], [f_{2,k}] \rangle = \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha u_{1,k} u_{2,k}^* \rangle \\ &= \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha r_{1,k} u_{2,k}^* \rangle + \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha v_1 r_{2,k} \rangle \\ &\quad + \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha v_1 v_2 \rangle. \end{aligned}$$

However, for the first two terms on the right hand side of (3.2) we can deduce

$$\begin{aligned} \left| \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha r_{1,k} u_{2,k}^* \rangle \right| &\leq \sum_{|\alpha| \leq m} |\langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha r_{1,k} u_{2,k}^* \rangle| \\ &\leq C \|u_{2,k}^*\|_{H^s} \|r_{1,k}\|_{H^s} \sum_{|\alpha| \leq m} \|a_{1,\alpha} - a_{2,\alpha}\|_{s-|\alpha|, -s} \rightarrow 0 \end{aligned}$$

and

$$\begin{aligned} \left| \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha v_1 r_{2,k} \rangle \right| &\leq \sum_{|\alpha| \leq m} |\langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha v_1 r_{2,k} \rangle| \\ &\leq C \|r_{2,k}\|_{H^s} \|v_1\|_{H^s} \sum_{|\alpha| \leq m} \|a_{1,\alpha} - a_{2,\alpha}\|_{s-|\alpha|, -s} \rightarrow 0 \end{aligned}$$

as $k \rightarrow \infty$. Thus by taking the limit in formula (3.2) we obtain

$$(19) \quad \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha v_1 v_2 \rangle = 0 \quad \text{for all } v_1, v_2 \in C_c^\infty(\Omega).$$

Step 2. Assume that we have $a_{1,\alpha}|_\Omega = a_{2,\alpha}|_\Omega$ for all α such that $|\alpha| < N$ for some $N \in \mathbb{N}$. We show that the equality of the coefficients also holds for α for which $|\alpha| = N$ and this will prove the theorem by the principle of complete induction.

To this end, consider $v_2 \in C_c^\infty(\Omega)$, and then take $v_1 \in C_c^\infty(\Omega)$ such that $v_1(x) = x^\alpha$ on $\text{supp}(v_2) \Subset \Omega$. Recall that since $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{N}^n$ is a multi-index and $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, the symbol x^α is intended to mean $x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}$. With this choice of v_1, v_2 , equation (3.2) becomes

$$\begin{aligned} (20) \quad 0 &= \sum_{|\beta| \leq m} \langle (a_{1,\beta} - a_{2,\beta}), D^\beta v_1 v_2 \rangle = \sum_{N \leq |\beta| \leq m} \langle (a_{1,\beta} - a_{2,\beta}), D^\beta (x^\alpha) v_2 \rangle \\ &= \sum_{N < |\beta| \leq m} \langle (a_{1,\beta} - a_{2,\beta}), D^\beta (x^\alpha) v_2 \rangle + \sum_{|\beta|=N, \beta \neq \alpha} \langle (a_{1,\beta} - a_{2,\beta}), D^\beta (x^\alpha) v_2 \rangle \\ &\quad + \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha (x^\alpha) v_2 \rangle. \end{aligned}$$

If $|\beta| > N = |\alpha|$, then there must exist $k \in \{1, 2, \dots, n\}$ such that $\beta_k > \alpha_k$. This is true also if $|\beta| = N$ with $\beta \neq \alpha$. In both cases we can compute

$$D^\beta (x^\alpha) = (\partial_{x_1}^{\beta_1} x_1^{\alpha_1}) (\partial_{x_2}^{\beta_2} x_2^{\alpha_2}) \dots (\partial_{x_n}^{\beta_n} x_n^{\alpha_n}) = 0$$

because $\partial_{x_k}^{\beta_k} x_k^{\alpha_k} = 0$. Therefore formula (3.2) becomes

$$0 = \langle (a_{1,\alpha} - a_{2,\alpha}), D^\alpha (x^\alpha) v_2 \rangle_{\mathbb{R}^n} = \alpha! \langle a_{1,\alpha} - a_{2,\alpha}, v_2 \rangle_{\mathbb{R}^n}$$

which by the arbitrariness of $v_2 \in C_c^\infty(\Omega)$ implies $a_{1,\alpha}|_\Omega = a_{2,\alpha}|_\Omega$ also for α for which $|\alpha| = N$.

Step 3. We have proved that $a_{1,\alpha}|_\Omega = a_{2,\alpha}|_\Omega$ for all α of the order $|\alpha| \leq m$. Since this entails $P_1|_\Omega = P_2|_\Omega$, the proof is complete. \square

4. MAIN THEOREM FOR BOUNDED COEFFICIENTS

We shall now study the case when the coefficients of PDOs are from the bounded spaces $H^{r_\alpha, \infty}(\Omega)$. It should be noted, however, that most of the considerations of the previous section still apply identically.

4.1. Well-posedness of the inverse problem. We shall define the bilinear forms for the problems (3.1) and (3.1) respectively by (3.1) and (3.1), just as in the case of singular coefficients. These will turn out to be bounded in $H^s(\mathbb{R}^n) \times H^s(\mathbb{R}^n)$ as well, but the proof we give of this fact is *a fortiori* different. Since now we assume that $a_\alpha \in H^{r_\alpha, \infty}(\Omega) \subset L^\infty(\Omega)$ for $r_\alpha \geq 0$, the duality pairing $\langle a_\alpha, D^\alpha v w \rangle$ becomes an inner product over Ω and we write $\langle a_\alpha(x) D^\alpha v, w \rangle_\Omega$ to emphasize that the coefficients $a_\alpha = a_\alpha(x)$ are now functions defined in Ω .

Lemma 4.1 (Boundedness of the bilinear forms). *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain and $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $m \in \mathbb{N}$ such that $2s > m$. Let $a_\alpha \in H^{r_\alpha, \infty}(\Omega)$, with r_α defined as in (1.2). Then B_P and B_P^* extend as bounded bilinear forms on $H^s(\mathbb{R}^n) \times H^s(\mathbb{R}^n)$.*

Remark 4.2. *Since $s \in \mathbb{R}^+ \setminus \mathbb{Z}$ and $|\alpha| \leq m < 2s$, we also have that $\max(0, |\alpha| - s) \leq r_\alpha < s$ for $\delta > 0$ small (see equation (1.2)).*

Proof of lemma 4.1. We only prove the boundedness of B_P , as for B_P^* one can proceed in the same way. If $v, w \in C_c^\infty(\mathbb{R}^n)$, then

$$|\langle a_\alpha(x) D^\alpha v, w \rangle_\Omega| = \left| \int_\Omega a_\alpha w D^\alpha v \, dx \right| \leq \|a_\alpha w\|_{(H^{-r_\alpha}(\Omega))^*} \|D^\alpha v\|_{H^{-r_\alpha}(\Omega)}.$$

Since Ω is a Lipschitz domain and $r_\alpha > -1/2$, $r_\alpha \notin \{\frac{1}{2}, \frac{3}{2}, \frac{5}{2}, \dots\}$, we have $(H^{-r_\alpha}(\Omega))^* = H_0^{r_\alpha}(\Omega) \subset H^{r_\alpha}(\Omega)$. Therefore

$$\begin{aligned} (21) \quad |\langle a_\alpha(x) D^\alpha v, w \rangle_\Omega| &\leq C \|a_\alpha w\|_{H^{r_\alpha}(\Omega)} \|D^\alpha v\|_{H^{-r_\alpha}(\Omega)} \leq C \|A_\alpha w\|_{H^{r_\alpha}(\mathbb{R}^n)} \|D^\alpha v\|_{H^{-r_\alpha}(\Omega)} \\ &\leq C \|J^{r_\alpha}(A_\alpha w)\|_{L^2(\mathbb{R}^n)} \|v\|_{H^{|\alpha| - r_\alpha}(\Omega)} \end{aligned}$$

where $J = (\text{Id} - \Delta)^{1/2}$ is the Bessel potential and A_α is an extension of a_α from Ω to \mathbb{R}^n such that $A_\alpha|_\Omega = a_\alpha$ and $\|A_\alpha\|_{H^{r_\alpha, \infty}(\mathbb{R}^n)} \leq 2\|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)}$. Since $r_\alpha \geq 0$, we may estimate the last term of (4.1) by the Kato-Ponce inequality given in lemma 2.3

$$\begin{aligned} \|J^{r_\alpha}(A_\alpha w)\|_{L^2(\mathbb{R}^n)} &\leq C \left(\|A_\alpha\|_{L^\infty(\mathbb{R}^n)} \|J^{r_\alpha} w\|_{L^2(\mathbb{R}^n)} + \|J^{r_\alpha} A_\alpha\|_{L^\infty(\mathbb{R}^n)} \|w\|_{L^2(\mathbb{R}^n)} \right) \\ &\leq C \|A_\alpha\|_{H^{r_\alpha, \infty}(\mathbb{R}^n)} \|w\|_{H^{r_\alpha}(\mathbb{R}^n)} \leq C \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|w\|_{H^{r_\alpha}(\mathbb{R}^n)}. \end{aligned}$$

Substituting this into (4.1) gives

$$(22) \quad \begin{aligned} |\langle a_\alpha(x) D^\alpha v, w \rangle_\Omega| &\leq C \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|w\|_{H^{r_\alpha}(\mathbb{R}^n)} \|v\|_{H^{|\alpha| - r_\alpha}(\Omega)} \\ &\leq C \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|w\|_{H^s(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)} \end{aligned}$$

given that both $r_\alpha < s$ and $|\alpha| - r_\alpha \leq s$ hold by remark 4.2. Eventually we obtain

$$\begin{aligned} |B_P(v, w)| &\leq | \langle (-\Delta)^{s/2} v, (-\Delta)^{s/2} w \rangle_{\mathbb{R}^n} | + \sum_{|\alpha| \leq m} | \langle a_\alpha D^\alpha v, w \rangle_{\mathbb{R}^n} | \\ &\leq \|w\|_{H^s(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)} + \sum_{|\alpha| \leq m} C \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|w\|_{H^s(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)} \\ &\leq C \|w\|_{H^s(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)}. \quad \square \end{aligned}$$

Next we shall prove existence and uniqueness of solutions for the problems (3.1) and (3.1). The reasoning is similar to the one for the proof of lemma 3.3, but the details of the computations are quite different.

Lemma 4.3 (Well-posedness). *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain and $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $m \in \mathbb{N}$ such that $2s > m$. Let $a_\alpha \in H^{r_\alpha, \infty}(\Omega)$, with r_α defined as in (1.2). There exist a real number $\mu > 0$ and a countable set $\Sigma \subset (-\mu, \infty)$ of eigenvalues $\lambda_1 \leq \lambda_2 \leq \dots \rightarrow \infty$ such that if $\lambda \in \mathbb{R} \setminus \Sigma$, for any $f \in H^s(\mathbb{R}^n)$ and $F \in (\tilde{H}^s(\Omega))^*$ there exists a unique $u \in H^s(\mathbb{R}^n)$ such that $u - f \in \tilde{H}^s(\Omega)$ and*

$$B_P(u, v) - \lambda \langle u, v \rangle_\Omega = F(v) \quad \text{for all } v \in \tilde{H}^s(\Omega).$$

One has the estimate

$$\|u\|_{H^s(\mathbb{R}^n)} \leq C \left(\|f\|_{H^s(\mathbb{R}^n)} + \|F\|_{(\tilde{H}^s(\Omega))^*} \right).$$

The function u is also the unique $u \in H^s(\mathbb{R}^n)$ satisfying

$$r_\Omega \left((-\Delta)^s + \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha - \lambda \right) u = F$$

in the sense of distributions in Ω and $u - f \in \tilde{H}^s(\Omega)$. Moreover, if (3.1) holds then $0 \notin \Sigma$.

Proof. Again it is enough to find unique $\tilde{u} \in \tilde{H}^s(\Omega)$ such that $B_P(\tilde{u}, v) - \lambda \langle \tilde{u}, v \rangle_\Omega = \tilde{F}(v)$, where $\tilde{F} := F - B_P(f, \cdot) + \lambda \langle f, \cdot \rangle_\Omega$. Consider $v, w \in C_c^\infty(\Omega)$ and $r_\alpha \neq 0$. Since $0 < r_\alpha < s$, the interpolation inequality

$$\|w\|_{H^{r_\alpha}(\mathbb{R}^n)} \leq C \|w\|_{L^2(\mathbb{R}^n)}^{1-r_\alpha/s} \|w\|_{H^s(\mathbb{R}^n)}^{r_\alpha/s}$$

holds. Using this and formula (4.1) we get, for a constant $C = C(\Omega, n, s, r_\alpha)$ which may change from line to line,

$$(23) \quad \begin{aligned} |\langle a_\alpha(x) D^\alpha v, w \rangle_\Omega| &\leq C \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|v\|_{H^s(\mathbb{R}^n)} \|w\|_{H^{r_\alpha}(\mathbb{R}^n)} \\ &\leq C \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|v\|_{H^s(\mathbb{R}^n)} \|w\|_{L^2(\mathbb{R}^n)}^{1-r_\alpha/s} \|w\|_{H^s(\mathbb{R}^n)}^{r_\alpha/s} \\ &\leq \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|v\|_{H^s(\mathbb{R}^n)} \left(C \epsilon^{r_\alpha/(r_\alpha-s)} \|w\|_{L^2(\mathbb{R}^n)} + \epsilon \|w\|_{H^s(\mathbb{R}^n)} \right). \end{aligned}$$

In the last step of (4.1) we used formula (3.1) with

$$q = \frac{s}{r_\alpha}, \quad p = \frac{s}{s - r_\alpha}, \quad b = \|w\|_{H^s(\mathbb{R}^n)}^{r_\alpha/s}, \quad a = C \|w\|_{L^2(\mathbb{R}^n)}^{1-r_\alpha/s}, \quad \eta = \epsilon.$$

If instead $r_\alpha = 0$, just by formula (4.1) we already have

$$|\langle a_\alpha(x)D^\alpha v, w \rangle_\Omega| \leq C \|a_\alpha\|_{L^\infty(\Omega)} \|v\|_{H^s(\mathbb{R}^n)} \|w\|_{L^2(\mathbb{R}^n)}.$$

Moreover, the two estimates above also hold for $v, w \in \tilde{H}^s(\Omega)$ by the density of $C_c^\infty(\Omega)$ in $\tilde{H}^s(\Omega)$. Now we use formula (3.1) again, but this time we choose

$$q = p = 2, \quad b = \|v\|_{H^s(\mathbb{R}^n)}, \quad a = \|v\|_{L^2(\mathbb{R}^n)}, \quad \eta = \epsilon^{s/(s-r_\alpha)}.$$

This leads to

$$\begin{aligned} |\langle a_\alpha(x)D^\alpha v, v \rangle_\Omega| &\leq \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \|v\|_{H^s(\mathbb{R}^n)} \left(C\epsilon^{r_\alpha/(r_\alpha-s)} \|v\|_{L^2(\mathbb{R}^n)} + \epsilon \|v\|_{H^s(\mathbb{R}^n)} \right) \\ &= \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \left(C\epsilon^{r_\alpha/(r_\alpha-s)} \|v\|_{L^2(\mathbb{R}^n)} \|v\|_{H^s(\mathbb{R}^n)} + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\leq \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \left(C\epsilon^{\frac{r_\alpha+s}{r_\alpha-s}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon(C+1) \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\leq C \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \left(\epsilon^{\frac{r_\alpha+s}{r_\alpha-s}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\leq C' \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \left(\epsilon^{\frac{M+s}{M-s}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \end{aligned}$$

where $C = C(\Omega, n, s, r_\alpha)$ and $C' = C'(\Omega, n, s)$ are constants changing from line to line and $M \in [0, s)$ is defined by $M := \max_{|\alpha| \leq m} r_\alpha$. Eventually

$$\begin{aligned} (24) \quad B_P(v, v) &\geq \|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 - \sum_{|\alpha| \leq m} |\langle a_\alpha(x)D^\alpha v, v \rangle_\Omega| \\ &\geq \|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 - C' \left(\epsilon^{\frac{M+s}{M-s}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \sum_{|\alpha| \leq m} \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)} \\ &= \|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 - C' C'' \left(\epsilon^{\frac{M+s}{M-s}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \end{aligned}$$

where $C'' := \sum_{|\alpha| \leq m} \|a_\alpha\|_{H^{r_\alpha, \infty}(\Omega)}$ is a constant independent of ϵ and v . By the higher order Poincaré inequality (lemma 2.2) (4.1) turns into

$$\begin{aligned} B_P(v, v) &\geq c \left(\|(-\Delta)^{s/2} v\|_{L^2(\mathbb{R}^n)}^2 + \|v\|_{L^2(\mathbb{R}^n)}^2 \right) - C' C'' \left(\epsilon^{\frac{M+s}{M-s}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \\ &\geq c \|v\|_{H^s(\mathbb{R}^n)}^2 - C' C'' \left(\epsilon^{\frac{M+s}{M-s}} \|v\|_{L^2(\mathbb{R}^n)}^2 + \epsilon \|v\|_{H^s(\mathbb{R}^n)}^2 \right) \end{aligned}$$

for some constant $c = c(\Omega, n, s)$ changing from line to line. For ϵ small enough (notice that $M - s < 0$), this eventually gives the coercivity estimate

$$(25) \quad B_P(v, v) \geq c_0 \|v\|_{H^s(\mathbb{R}^n)}^2 - \mu \|v\|_{L^2(\mathbb{R}^n)}^2$$

for some constants $c_0, \mu > 0$ independent of v . The proof is now concluded as in lemma 3.3. \square

Assuming as in Section 3 that both (3.1) and (3.1) hold, by means of the above lemma 4.3 we can define the DN-maps Λ_P, Λ_P^* just as in lemma 3.4. We also arrive at the same Alessandrini identity and Runge approximation property which we get in lemmas 3.6 and 3.7.

Lemma 4.4 (DN maps). *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain and $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $m \in \mathbb{N}$ such that $2s > m$. Let $a_\alpha \in H^{r_\alpha, \infty}(\Omega)$, with r_α defined as in (1.2). There exist two continuous linear maps*

$$\Lambda_P : X \rightarrow X^* \quad \text{defined by} \quad \langle \Lambda_P[f], [g] \rangle := B_P(u_f, g)$$

and

$$\Lambda_P^* : X \rightarrow X^* \quad \text{defined by} \quad \langle \Lambda_P^*[f], [g] \rangle := B_P^*(u_f^*, g)$$

where u_f, u_f^* are the unique solutions to the equations

$$(-\Delta)^s u + \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha u = 0 \quad \text{in } \Omega, \quad u - f \in \tilde{H}^s(\Omega)$$

and

$$(-\Delta)^s u^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha(x) u^*) = 0 \quad \text{in } \Omega, \quad u^* - f \in \tilde{H}^s(\Omega)$$

with $f, g \in H^s(\mathbb{R}^n)$. Moreover, the identity $\langle \Lambda_P[f], [g] \rangle = \langle [f], \Lambda_P^*[g] \rangle$ holds.

Lemma 4.5 (Alessandrini identity). *Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain and $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $m \in \mathbb{N}$ such that $2s > m$. Let $a_\alpha \in H^{r_\alpha, \infty}(\Omega)$, with r_α defined as in (1.2). For any $f_1, f_2 \in H^s(\mathbb{R}^n)$, let $u_1, u_2^* \in H^s(\mathbb{R}^n)$ respectively solve*

$$(-\Delta)^s u_1 + \sum_{|\alpha| \leq m} a_{1,\alpha}(x) D^\alpha u_1 = 0 \quad \text{in } \Omega, \quad u_1 - f_1 \in \tilde{H}^s(\Omega)$$

and

$$(-\Delta)^s u_2^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_{2,\alpha}(x) u_2^*) = 0 \quad \text{in } \Omega, \quad u_2^* - f_2 \in \tilde{H}^s(\Omega).$$

Then we have the integral identity

$$\langle (\Lambda_{P_1} - \Lambda_{P_2})[f_1], [f_2] \rangle = \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}) D^\alpha u_1, u_2^* \rangle_\Omega.$$

Lemma 4.6 (Runge approximation property). *Let $\Omega, W \subset \mathbb{R}^n$ respectively be a bounded Lipschitz domain and a non-empty open set such that $\overline{W} \cap \overline{\Omega} = \emptyset$. Let $s \in \mathbb{R}^+ \setminus \mathbb{Z}$, $m \in \mathbb{N}$ such that $2s > m$. Let $a_\alpha \in H^{r_\alpha, \infty}(\Omega)$, with r_α defined as in (1.2). Moreover, let $\mathcal{R} := \{u_f - f : f \in C_c^\infty(W)\} \subset \tilde{H}^s(\Omega)$, where u_f solves*

$$(-\Delta)^s u_f + \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha u_f = 0 \quad \text{in } \Omega, \quad u_f - f \in \tilde{H}^s(\Omega)$$

and $\mathcal{R}^* := \{u_f^* - f : f \in C_c^\infty(W)\} \subset \tilde{H}^s(\Omega)$, where u_f^* solves

$$(-\Delta)^s u_f^* + \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha(x) u_f^*) = 0 \quad \text{in } \Omega, \quad u_f^* - f \in \tilde{H}^s(\Omega).$$

Then \mathcal{R} and \mathcal{R}^* are dense in $\tilde{H}^s(\Omega)$.

4.2. Proof of injectivity.

Proof of theorem 1.2. The proof is virtually identical to the one of theorem 1.1, the unique difference being in the way the error terms of the Runge approximation are estimated. We make use of (4.1), which relied on the Kato-Ponce inequality instead of multiplier space estimates. In this way we get

$$\begin{aligned} \left| \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}) D^\alpha r_{1,k}, u_{2,k}^* \rangle_{\mathbb{R}^n} \right| &\leq \sum_{|\alpha| \leq m} | \langle (a_{1,\alpha} - a_{2,\alpha}) D^\alpha r_{1,k}, u_{2,k}^* \rangle_{\mathbb{R}^n} | \\ &\leq C \|u_{2,k}^*\|_{H^s(\mathbb{R}^n)} \|r_{1,k}\|_{H^s(\mathbb{R}^n)} \sum_{|\alpha| \leq m} \|a_{1,\alpha} - a_{2,\alpha}\|_{H^{r_\alpha, \infty}(\Omega)} \rightarrow 0 \end{aligned}$$

and

$$\begin{aligned} \left| \sum_{|\alpha| \leq m} \langle (a_{1,\alpha} - a_{2,\alpha}) D^\alpha v_1, r_{2,k} \rangle_{\mathbb{R}^n} \right| &\leq \sum_{|\alpha| \leq m} | \langle (a_{1,\alpha} - a_{2,\alpha}) D^\alpha v_1, r_{2,k} \rangle_{\mathbb{R}^n} | \\ &\leq C \|r_{2,k}\|_{H^s(\mathbb{R}^n)} \|v_1\|_{H^s(\mathbb{R}^n)} \sum_{|\alpha| \leq m} \|a_{1,\alpha} - a_{2,\alpha}\|_{H^{r_\alpha, \infty}(\Omega)} \rightarrow 0. \quad \square \end{aligned}$$

REFERENCES

- [1] H. Abels. Pseudodifferential and Singular Integral Operators. De Gruyter, First edition, 2012.
- [2] F. Andreu-Vailló, J. M. Mazón, J. D. Rossi, and J. J. Toledo-Melero. Nonlocal Diffusion Problems. American Mathematical Society, First edition, 2010.
- [3] A. Behzadan and M. Holst. Multiplication in Sobolev spaces, revisited. 2017. arXiv:1512.07379v2.
- [4] S. Bhattacharyya, T. Ghosh, and G. Uhlmann. Inverse Problem for Fractional-Laplacian with Lower Order Non-local Perturbations. *Transactions of the AMS*, 2020. To appear.
- [5] C. Bucur and E. Valdinoci. Nonlocal Diffusion and Applications. Springer, First edition, 2016.
- [6] X. Cao, Y.-H. Lin, and H. Liu. Simultaneously recovering potentials and embedded obstacles for anisotropic fractional Schrödinger operators. *Inverse Probl. Imaging*, 13(1):197–210, 2019.
- [7] M. Cekić, Y.-H. Lin, and A. Rüländ. The Calderón problem for the fractional Schrödinger equation with drift. *Calculus of Variations and Partial Differential Equations*, 59(3):91, 2020.
- [8] G. Covi. Inverse problems for a fractional conductivity equation. *Nonlinear Analysis*, 2018.
- [9] G. Covi. An inverse problem for the fractional Schrödinger equation in a magnetic field. *Inverse Problems*, 36(4):045004, 2020.
- [10] G. Covi, K. Mönkkönen, and J. Railo. Unique continuation property and Poincaré inequality for higher order fractional Laplacians with applications in inverse problems. 2020. arXiv:2001.06210.
- [11] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bulletin des Sciences Mathématiques*, 136(5):521–573, 2012.
- [12] Q. Du, M. Gunzburger, R. B. Lehoucq, and K. Zhou. Analysis and Approximation of Nonlocal Diffusion Problems with Volume Constraints. *SIAM Rev.*, 54, No. 4:667–696, 2012.
- [13] T. Ghosh, Y.-H. Lin, and J. Xiao. The Calderón problem for variable coefficients nonlocal elliptic operators. *Communications in Partial Differential Equations*, 42(12):1923–1961, 2017.
- [14] T. Ghosh, A. Rüländ, M. Salo, and G. Uhlmann. Uniqueness and reconstruction for the fractional Calderón problem with a single measurement. *Journal of Functional Analysis*, 279(1):108505, 2020.
- [15] T. Ghosh, M. Salo, and G. Uhlmann. The Calderón problem for the fractional Schrödinger equation. *Anal. PDE* 13(2):455-475, 2020.
- [16] L. Grafakos and S. Oh. The Kato-Ponce inequality. *Communications in Partial Differential Equations*, 39(6):1128-1157, 2014.
- [17] A. Gulisashvili and M. A. Kon. Exact Smoothing Properties of Schrödinger Semigroups. *American Journal of Mathematics*, 118(6):1215–1248, 1996.
- [18] T. Kato and G. Ponce. Commutator Estimates and the Euler and Navier-Stokes Equations. *Communications on Pure and Applied Mathematics*, 41(7):891–907, 1988.
- [19] R.-Y. Lai and Y.-H. Lin. Global uniqueness for the fractional semilinear Schrödinger equation. *Proc. Amer. Math. Soc.*, 147(3):1189–1199, 2019.
- [20] R.-Y. Lai and Y.-H. Lin. Inverse problems for fractional semilinear elliptic equations. 2020. arXiv:2004.00549.
- [21] R.-Y. Lai, Y.-H. Lin, and A. Rüländ. The Calderón Problem for a Space-Time Fractional Parabolic Equation, 2020.
- [22] N. Laskin. Fractional Quantum Mechanics and Lévy Path Integrals. *Physics Letters A*, 268(4):298 – 305, 2000.
- [23] N. Laskin. Fractional Quantum Mechanics. World Scientific, First edition, 2018.
- [24] L. Li. A Semilinear Inverse Problem For The Fractional Magnetic Laplacian. 2020. arXiv:2005.06714.
- [25] L. Li. Determining The Magnetic Potential In The Fractional Magnetic Calderón Problem. 2020. arXiv:2006.10150.
- [26] L. Li. The Calderón problem for the fractional magnetic operator. *Inverse Problems*, 36(7):075003, 2020.
- [27] A. Massaccesi and E. Valdinoci. Is a nonlocal diffusion strategy convenient for biological populations in competition? *Journal of Mathematical Biology*, 74(1):113–147, 2017.
- [28] V. G. Maz’ya and T. O. Shaposhnikova. Theory of Sobolev Multipliers. Springer, First edition, 2009.
- [29] W. McLean. Strongly Elliptic Systems and Boundary Integral Equations. Cambridge University Press, First edition, 2000.
- [30] M. Mišur. A Refinement of Peetre’s Theorem. *Results in Mathematics*, 74(4):199, 2019.
- [31] J. Navarro and J. B. Sancho. Peetre-Slovák’s theorem revisited. 2014. arXiv:1411.7499.
- [32] J. Peetre. Une caractérisation abstraite des opérateurs différentiels. *Mathematica Scandinavica*, 7:211–218, 1959.
- [33] X. Ros-Oton. Nonlocal elliptic equations in bounded domains: a survey. *Publicacions Matemàtiques*, 60:3 – 26, 2015.
- [34] A. Rüländ and M. Salo. Exponential instability in the fractional Calderón problem. *Inverse Problems*, 34(4):045003, 21, 2018.
- [35] A. Rüländ and M. Salo. Quantitative approximation properties for the fractional heat equation. *Mathematical Control & Related Fields*, pages 233–249, 2019.
- [36] A. Rüländ and M. Salo. The fractional Calderón problem: Low regularity and stability. *Nonlinear Analysis*, 2019.

- [37] M. Salo. The fractional Calderón problem. *Journées équations aux dérivées partielles*, Exp. No.(7), 2017.
- [38] H. Triebel. Interpolation Theory, Function Spaces, Differential Operators. North-Holland Publishing Company, 1978.
- [39] G. Uhlmann. Electrical impedance tomography and Calderón’s problem. *Inverse Problems*, 25(12):123011, 2009.
- [40] G. Uhlmann. Inverse problems: seeing the unseen. *Bulletin of Mathematical Sciences*, 4(2):209–279, 2014.
- [41] M. W. Wong. An Introduction to Pseudo-Differential Operators. World Scientific, Third edition, 2014.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ

E-mail address: `giovanni.g.covi@jyu.fi`

E-mail address: `kematamo@student.jyu.fi`

E-mail address: `jesse.t.railo@jyu.fi`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON / JOCKEY CLUB INSTITUTE FOR ADVANCED STUDY, HKUST, HONG KONG

E-mail address: `gunther@math.washington.edu`