# The Non-Abelian X-Ray Transform on Asymptotically Hyperbolic Spaces

## Haim Grebnev

## April 26, 2024

#### Abstract

In this paper we formulate and prove a gauge equivalence for connections and Higgs fields of suitable regularity that are mapped to the same function under the non-abelian X-ray transform on nontrapping asymptotically hyperbolic (AH) spaces with negative curvature and no nontrivial twisted conformal Killing tensor fields with certain regularity. If one furthermore fixes such a connection with zero curvature, a corollary provides an injectivity result for the non-abelian X-ray transform over Higgs fields.

# Contents

| 1        | Conv         | entions/notations                                     | <b>2</b> |
|----------|--------------|---|----------|
| <b>2</b> | Introduction |   |          |
|          | 2.1 I        | Motivation  | 2        |
|          | 2.2 A        | Asymptotically Hyperbolic Spaces                      | 4        |
|          | 2.3 I        | Results   | 4        |
|          | 2.4 I        | Existence of Examples                                 | 8        |
|          | 2.5 I        | Non-Abelian X-Ray Transform                           | 9        |
|          | 2.6 I        | Prior Research Discussion and Applications            | 11       |
|          | 2.7 A        | Acknowledgments                                       | 12       |
|          |              |   |          |
| 3        | Well-        | Definedness of the Non-abelian X-Ray Transform        | 12       |
| 4        | Geon         | netric Preliminaries                                  | 13       |
|          | 4.1          | The b and 0 Cotangent Bundles                         | 13       |
|          | 4.2          | The b and 0 Cosphere Bundles                          | 15       |
|          | 4.3 \$       | Splitting the Tangent Bundle                          | 15       |
|          | 4.4 I        | Integration on the (Co)Sphere Bundle                  | 17       |
|          | 4.5 \$       | Splitting the Connection Over the Unit Tangent Bundle | 17       |
|          | 4.6 0        | Curvatures  | 19       |

| <b>5</b> | Gauge Equivalence of Connections and Higgs fields     | <b>21</b> |
|----------|---|-----------|
|          | 5.1 The Regularity Spaces                             | 21        |
|          | 5.2 Pestov Identity                                   | 22        |
|          | 5.3 Finite Degree of Solutions to Transport Equations | 30        |
|          | 5.4 Regularity of Solutions to the Transport Equation | 40        |
|          | 5.5 Proof of Theorem 2.7                              | 45        |
| 6        | Proof of Theorem 2.14                                 | 48        |
| 7        | Injectivity over Higgs Fields                         | 49        |

# 1 Conventions/notations

In this paper we employ the following conventions/notations:

- 1. We employ the Einstein summation convention.
- 2. The dimension of our manifold will be n + 1, and we'll denote all indices related to its dimension by  $0, 1, \ldots, n$ . When using the Einstein summation convention on indices related to the manifold's dimension, we employ the convention that Latin indices run from 0 to n while Greek indices run from 1 to n.
- 3. The notation  $C^k$  denotes k times continuously differentiable objects. If the object is scalar valued, we always assume that it is complex valued (e.g.  $C^k(\overline{M}) = C^k(\overline{M}; \mathbb{C})$ ).
- 4. Whenever we say "smooth," we mean " $C^{\infty}$ ." All diffeomorphisms are smooth.
- 5. If  $\pi: E \to N$  is a vector bundle over a manifold N and  $S \subseteq N$  is a subset of N, then we let  $E|_S$  denote the restriction of E to the fibers over S (i.e. more precise notation would be  $E|_{\pi^{-1}[S]}$ ).
- 6. Continuing off of point 5), we write  $C^{\infty}(N; E)$  for smooth sections of E (i.e. not simply smooth maps from N to E).
- 7. We denote the geodesic vector field over the tangent and cotangent bundles (i.e. infinitesimal generator of the geodesic flow) by X. We will also let X denote its restriction to the sphere and cosphere bundles SM and  $S^*M$  (described below), which makes sense because X is tangent to it.

# 2 Introduction

## 2.1 Motivation

We begin by providing motivation for the non-abelian X-ray transform - the object of our interest, delaying precise definitions for a later section. This transform is a generalization of the so-called "scalar X-ray transform," the latter of which is used in reconstructing images

of the internals of patients after irradiating them with X-rays at various angles. The typical mathematical problem for the scalar X-ray transform is the following: suppose that we have a bounded subset  $D \subseteq \mathbb{R}^n$  with smooth boundary and a continuous function  $\phi : D \to (0, \infty)$ over it. In our analogy, D represents the shape of our patient and  $\phi$  the body's X-ray absorption coefficient at various points. Suppose we have a parametrized line l(t) that enters D at t = 0 and leaves at  $t = t_{\text{exit}}$ , which represents the motion of an X-ray moving through the body. The ray's intensity I(t) along this path decays according to the law

$$\frac{dI}{dt} = -\phi I, \qquad I(0) = I_0, \qquad (2.1)$$

where  $I_0$  represents the initial intensity of the ray. We record  $I(t_{\text{exit}})$  (i.e. the intensity of the ray when it exits) and then repeat this procedure for all possible lines l that pass through D, using the same value for  $I_0$  every time. The inverse problem is then to recover the coefficient  $\phi$  from knowing such data, which is equivalent to recovering a gray-scale image of the patient's internals. Immediately we may note that a necessary condition for this to be possible to do is that two different  $\phi$ 's cannot generate the same observation data. If this is the case, we say that the scalar X-ray transform that takes  $\phi$  to the observed data is "injective." It turns out that the answer to this problem is yes: one can indeed recover  $\phi$ from the mentioned data - see for instance Theorem 1.1.6 in [40].

The non-abelian X-ray transform is defined similarly, except that we turn (2.1) into a system of equations by letting I by a column vector and  $\phi$  a square matrix. The question is also if one can recover the matrix  $\phi$  from the collected data, or in other words if the operator involved is injective. One application of this is in the recently introduced *polarimetric neutron* tomography which attempts to reconstruct the structure of magnetic fields inside materials after sending neutron beams through them - see for instance [20] and [8]. We will mention a few more applications of this problem with references in Section 2.6 below.

We will actually be interested in a more sophisticated generalization of the transform, for instance by allowing the paths "l" to be geodesics with respect to some Riemannian metric g on D where D is now a smooth manifold. We will formulate I and  $\phi$  to be a section and endomorphism field respectively of a smooth vector bundle over M and formulate the time derivative to be a connection (which may also be unknown) in the direction of the curve's velocity. It turns out that in this more general setting it is not possible to recover all of the coefficients involved except in a special case when an additional assumption is made on the connection - see for instance Theorem 2.7 and Corollary 2.10 below. However, in the case when we can't recover the coefficients, we do have a "gauge equivalence" understanding of coefficients that produce the same data.

Such transforms have been well studied in the context of compact domains, and so a direction of research that has received significant attention in recent years is whether it's possible to generalize X-ray results to noncompact domains. We in particular will be pursuing this direction of research in this paper, in particular generalizing known results for the non-abelian X-ray transform to a particular class of noncompact manifolds called "asymptotically hyperbolic spaces."

### 2.2 Asymptotically Hyperbolic Spaces

In this section we introduce the geometry on which our transform will be defined. In this paper we let  $\overline{M}$  be a compact smooth manifold with smooth boundary of dimension n+1 with  $n \geq 1$ , whose interior we denote by M. We fix a boundary defining function  $\rho : \overline{M} \to [0, \infty)$  (i.e.  $\rho$  is smooth,  $\rho = 0$  on and only on  $\partial \overline{M}$ , and  $d\rho|_{\partial \overline{M}} \neq 0$ ). We also fix an **asymptotically hyperbolic (AH) Riemannian metric** g on M, which is defined as a metric such that the tensor  $\overline{g} = \rho^2 g$  extends to a smooth Riemannian metric on all of  $\overline{M}$  with  $|d\rho|_{\rho^2 g}^2 \equiv 1$  along  $\partial \overline{M}$ . The boundary of  $\partial \overline{M}$  is thought of as the "infinity" where the metric g blows up. Hence recalling that hyperbolic space has constant sectional curvature -1, the known fact that the sectional curvatures of g tend to -1 as one approaches  $\partial \overline{M}$  explains why such metrics are given the name "AH.<sup>1</sup>"

In fact, the Poincaré ball model of hyperbolic space is the archetypical example of an AH space. It is given by  $\overline{M} = \{|x| \leq 1\} \subseteq \mathbb{R}^{n+1}$ , where |x| denotes the Euclidean length, and

$$g = 4 \frac{(dx^{1})^{2} + \ldots + (dx^{n+1})^{2}}{(1 - |x|^{2})^{2}}.$$

Indeed if one takes the boundary defining function  $\rho = 1 - |x|^2$ , then an elementary exercise shows that  $|d\rho|^2_{\rho^2 g} \equiv 1$  along  $\{|x| = 1\}$ .

Taking our general AH space, we note that we can always construct boundary coordinates of  $\overline{M}$  of the form  $(\rho, y^1, \ldots, y^n)$  in which the metric g takes the following convenient form. Let  $\varepsilon > 0$  be such that the flowout of the gradient of  $\rho$  with respect to the (smooth) metric  $\overline{g} = \rho^2 g$  is a diffeomorphism from  $(0, \varepsilon) \times \partial \overline{M}$  onto a collar neighborhood  $\mathcal{C}_{\varepsilon}$  of  $\partial \overline{M}$ . Then, fixing coordinates  $(y^1, \ldots, y^n)$  of  $\partial \overline{M}$ , this flowout provides us boundary coordinates  $(\rho, y^{\mu})$ of  $\overline{M}$  in which the metric takes the form

$$g = \frac{d\rho^2 + h_{\mu\nu}dy^{\mu}dy^{\nu}}{\rho^2}.$$
 (2.2)

We call such boundary coordinates  $(\rho, y^{\mu})$  asymptotic boundary normal coordinates. We will often assume that our boundary coordinates are in this form because several results that we cite from [14] are only stated in such coordinates. However, assuming this extra assumption on the coordinates will not cause us any issues.

By Proposition 1.8 in [29], AH spaces are complete. Furthermore, in some cases we will assume that g is also **nontrapping** which means that for any complete g-geodesic  $\gamma : (-\infty, \infty) \to M$ ,  $\liminf_{t \to \pm \infty} \rho(\gamma(t)) = 0$ . Intuitively speaking, this condition requires that  $\gamma$  eventually "escapes to infinity."

#### 2.3 Results

We now state our results. As mentioned in Section 2.2 above we assume in this paper that  $(M \subseteq \overline{M}, g)$  is an AH space and that  $\rho$  is a boundary defining function. Throughout this

<sup>&</sup>lt;sup>1</sup>More generally, the sectional curvatures approaches  $-|d\rho|^2$  restricted to the boundary - see [29] for a precise statement.

paper we also assume that we have a smooth complex *d*-dimensional vector bundle  $\mathcal{E}$  over  $\overline{M}$  equipped with a smooth Hermitian inner product  $\langle \cdot, \cdot \rangle_{\mathcal{E}}$ . Moreover, we assume that we have a smooth section of the endomorphism bundle  $\Phi : \overline{M} \to \operatorname{End}_{\operatorname{Sk}} \mathcal{E}$  that is skew-Hermitian with respect to  $\langle \cdot, \cdot \rangle_{\mathcal{E}}$  (i.e.  $\langle \Phi u, v \rangle_{\mathcal{E}} = -\langle u, \Phi v \rangle_{\mathcal{E}}$ ). Lastly, we assume that we have a unitary smooth connection  $\nabla^{\mathcal{E}}$  in  $\mathcal{E}$  over  $\overline{M}$  with respect to  $\langle \cdot, \cdot \rangle_{\mathcal{E}}$  - meaning that

$$V\langle u, v \rangle_{\mathcal{E}} = \langle \nabla_V^{\mathcal{E}} u, v \rangle_{\mathcal{E}} + \langle u, \nabla_V^{\mathcal{E}} v \rangle_{\mathcal{E}},$$

when V is any smooth vector field over  $\overline{M}$  and u, v are any smooth sections of  $\mathcal{E}$ .

Take any unit-speed complete geodesic  $\gamma : (-\infty, \infty) \to M$  such that  $\liminf_{t \to \pm \infty} \rho(\gamma(t)) = 0$ . It follows from Lemma 2.3 in [14] that the limit of  $\gamma(t)$  in  $\overline{M}$  exists as  $t \to \pm \infty$ . The analog of the (2.1) that we will be considering is the following initial value problem for a section  $u : (-\infty, \infty) \to \mathcal{E}$ :

$$\nabla_{\dot{\gamma}(t)}^{\mathcal{E}} u(t) + \Phi(\gamma(t))u(t) = 0, \quad \lim_{t \to -\infty} u(t) = e, \tag{2.3}$$

where e is any element in  $\mathcal{E}_{x_0}$  where  $x_0 \in \partial \overline{M}$  is the limit of  $\gamma(t)$  as  $t \to -\infty$ . The data point that we "record" is

$$\lim_{t \to \infty} u(t). \tag{2.4}$$

The question that we are interested then becomes whether we can recover  $\Phi$  and  $\nabla^{\mathcal{E}}$  from the data recorded for all such possible pairs  $\gamma$  and e.

A bit of vocabulary: (2.3) is a type of differential equation called a **transport equation**, and  $\Phi$  is called a **Higgs field**. Going from the pair  $(\nabla^{\mathcal{E}}, \Phi)$  to the map that takes every  $(\gamma, e)$  as above to its associated data (2.4) is called the **non-abelian X-ray transform**, of which we give a more precise definition in Section 2.5 below.

To make rigorous sense of our problem however, we need to establish the well-definedness of the solution to (2.3) and the data (2.4). Considering that we're making use of the values of the solution to (2.3) at plus or minus infinities, we accomplish this by imposing a decay condition on  $\Phi$ . The following is our main existence lemma:

**Lemma 2.5.** Suppose that  $\Phi \in \rho C^{\infty}(\overline{M}; \operatorname{End}_{\operatorname{Sk}} \mathcal{E})$  and that  $\gamma : (-\infty, \infty) \to M$  is a complete geodesic such that  $\liminf_{t \to \pm \infty} \rho(\gamma(t)) = 0$ . Then for any  $e \in \mathcal{E}_{x_0}$  where  $x_0 = \lim_{t \to -\infty} \gamma(t) \in \partial \overline{M}$ , the solution to (2.3) exists and so does the limit (2.4).

For future use, we remark that the above lemma and its proof work equally well if one changes " $t \to \pm \infty$ " to " $t \to \mp \infty$ " in its statement and in (2.3) and (2.4).

Before we state our main result, we establish a way of talking about the decay regularity of the connection  $\nabla^{\mathcal{E}}$ :

**Definition 2.6.** If  $N \ge 0$  is an integer, we say that the "connection symbols of  $\nabla^{\mathcal{E}}$  are in  $\rho^N C^{\infty}(\overline{M})$  in any boundary coordinates (of  $\overline{M}$ ) and frame (for  $\mathcal{E}$ )" if the following holds. For any boundary coordinates  $(x^i)$  of  $\overline{M}$  and any frame  $(b_k)$  of  $\mathcal{E}$  over these coordinates' domain, the connection symbols  ${}^{\mathcal{E}}\Gamma^k_{ij}$  in the expression

$$\nabla_v^{\mathcal{E}} u = {}^{\mathcal{E}} \Gamma_{ij}^k v^i u^k b_k$$

satisfy  ${}^{\mathcal{E}}\Gamma_{ij}^k \in \rho^N C^{\infty}(\overline{M})$ , where  $v^i$  are the components of v with respect to  $(\partial/\partial x^i)$  and  $u^k$  are the components of u with respect to  $b_k$ .

As we mentioned earlier, the answer to our main problem is that we cannot recover the connection and Higgs field from the data (2.4) because such data can come from two distinct pairs  $(\nabla^{\mathcal{E}}, \Phi)$  and  $(\widetilde{\nabla}^{\mathcal{E}}, \widetilde{\Phi})$ . However if that is the case, the two pairs  $(\nabla^{\mathcal{E}}, \Phi)$  and  $(\widetilde{\nabla}^{\mathcal{E}}, \widetilde{\Phi})$  are related by a well understood gauge relation. The following is our main result on the matter. To state it, we use the regularity spaces  $\mathcal{R}^k(SM; \pi^*\mathcal{E})$  and the notion of *nontrivial twisted conformal Killing tensor fields* (CKTs for short) which are defined in Section 5.1 and at the end of Section 5.3 respectively. Their role of the latter are to allow us to formulate a technical assumption the connections that we need to make the vertical Fourier analysis in the proof to work. To state the regularity conclusion at the end, we also use the unit tangent bundle  $SM = \{v \in TM : |v|_g = 1\}$  with projection map  $\pi : SM \to M$  and the notion of pullback bundles " $\pi^* \dots$ " (see Section 4.5 below).

**Theorem 2.7.** Assume that (M, g) is nontrapping and that the sectional curvatures of g are negative. There exists an integer  $N_0 \geq 0$  big enough dependent only on (M, g) such that the following holds. Suppose that  $\Phi \in \rho^{N_0+1}C^{\infty}(\overline{M}; \operatorname{End}_{\operatorname{Sk}}\mathcal{E})$  and that the connection symbols of  $\nabla^{\mathcal{E}}$  are in  $\rho^{N_0}C^{\infty}(\overline{M})$  in any boundary coordinates and frame (in the sense of Definition 2.6), and that we have another pair  $\widetilde{\Phi}$  and  $\widetilde{\nabla}^{\mathcal{E}}$  satisfying the same conditions. Suppose that the data (2.4) for all possible  $\gamma$  and e as above are the same for (2.3), and (2.3) with  $\Phi$  and  $\nabla^{\mathcal{E}}$  replaced by  $\widetilde{\Phi}$  and  $\widetilde{\nabla}^{\mathcal{E}}$  respectively. Then there exists an everywhere invertible  $Q \in C^0(\overline{M}; \operatorname{End} \mathcal{E}) \cap C^{\infty}(M; \operatorname{End} \mathcal{E}|_M)$  such that  $Q|_{\partial\overline{M}} = \operatorname{id}$  and over  $\overline{M}$  satisfies

$$\widetilde{\nabla}^{\mathcal{E}} = Q^{-1} \nabla^{\mathcal{E}} Q, \qquad \qquad \widetilde{\Phi} = Q^{-1} \Phi Q. \tag{2.8}$$

Furthermore,  $(Q - \mathrm{id}) \in \mathcal{R}^3(SM; \pi^* \operatorname{End} \mathcal{E})$  when  $Q - \mathrm{id}$  is lifted from M to SM by setting  $(Q - \mathrm{id})(v) = (Q - \mathrm{id})(x)$  for any  $v \in S_x M$  using the canonical identification  $(\pi^* \operatorname{End} \mathcal{E})_v \cong (\operatorname{End} \mathcal{E})_x$ .

**Remark 2.9.** The notation  $\widetilde{\nabla}^{\mathcal{E}} = Q^{-1} \nabla^{\mathcal{E}} Q$  means

$$\widetilde{\nabla}_v^{\mathcal{E}} u = Q^{-1} \nabla_v^{\mathcal{E}} (Q u)$$

for any tangent vector  $v \in TM$  and any section  $u \in C^{\infty}(\overline{M}; \mathcal{E})$ . The equations (2.8) are called the **gauge relation** between the pairs  $(\nabla^{\mathcal{E}}, \Phi)$  and  $(\widetilde{\nabla}^{\mathcal{E}}, \widetilde{\Phi})$ , and we will provide intuition below for where it comes from.

The natural question arises of what the value of  $N_0$  is in the above theorem that determines the decay rate required of both the Higgs fields (e.g.  $\Phi$ ) and the connection symbols of the connections (e.g. of  $\nabla^{\mathcal{E}}$ ). We do not attempt to answer this question or to prove an upper bound. In this paper we prove that it only depends on the geometry of the space (M, g). Its size comes up when proving the regularity theorem for the transport equation, and in particular we will need it to be big enough so that the decay rate of  $\Phi$  and the connection symbols will overpower the growth of the derivatives of the geodesic flow. Since the latter are difficult to compute, albeit perhaps in some special cases such as the hyperbolic space, our approach does not indicate exactly how big  $N_0$  needs to be for the proof to work. We point out that in the special case of when the connection is known and has zero curvature (see Section 4.6 below for the latter), then it is possible to recover the Higgs field. Furthermore, in this case we can also drop the assumption about the nonexistence of twisted CKTs with suitable regularity because we'll get it for free from the zero-curvature assumption. Here is the precise statement:

**Corollary 2.10.** Assume that (M, g) is nontrapping, the sectional curvatures of g are negative, and that the curvature of  $\nabla^{\mathcal{E}}$  is zero everywhere. There exists an integer  $N_0 \geq 0$  big enough dependent only on (M, g) such that the following holds. Suppose that the connection symbols of  $\nabla^{\mathcal{E}}$  are in  $\rho^{N_0}C^{\infty}(\overline{M})$  in any boundary coordinates and frame (in the sense of Definition 2.6). Suppose also that  $\Phi, \widetilde{\Phi} \in \rho^{N_0+1}C^{\infty}(\overline{M}; \operatorname{End}_{\operatorname{Sk}}\mathcal{E})$  are such that the data (2.4) for all possible  $\gamma$  and e as above are the same for (2.3), and (2.3) with  $\Phi$  replaced by  $\widetilde{\Phi}$ . Then  $\Phi = \widetilde{\Phi}$ .

A simple example of when the corollary applies is if  $\overline{M}$  is a subset of  $\mathbb{R}^{n+1}$ ,  $\mathcal{E} = \overline{M} \times \mathbb{C}^d$ is the trivial bundle whose sections we write as column vectors, and the connection  $\nabla^{\mathcal{E}}$  is simply given by  $\nabla_v^{\mathcal{E}} u = [v(u^1), \dots, v(u^d)]$ .

We end this section with discussion on intuition and outline of the proofs. The relation (2.8) may look mysterious at first, so let us give intuition for it. The proof of Theorem 2.7 essentially begins with taking (2.3) and rewriting it in terms of endomorphism fields  $U : \mathbb{R} \to \text{End} \mathcal{E}$  in a way that it encodes the same data:

$$\nabla_{\dot{\gamma}(t)}^{\mathrm{En}} U(t) + \Phi(\gamma(t))U(t) = 0, \quad \lim_{t \to -\infty} U(t) = \mathrm{id}, \quad (2.11)$$

where " $\nabla^{\text{En}}$ " is a natural connection on the space of endomorphism fields  $\text{End} \mathcal{E}$ . Then we ask the following question. Suppose that we have a connection  $\nabla^{\mathcal{E}}$ , Higgs field  $\Phi$ , and the data (2.4) that it generates as above. How can we "come up" with another pair  $\widetilde{\nabla}^{\mathcal{E}}$  and  $\widetilde{\Phi}$  that generates the same data set? To do this, we take an arbitrary endomorphism field Q and manipulate the above equation as follows:

$$\nabla_{\dot{\gamma}}^{\mathrm{En}}(QQ^{-1}U) + \Phi(QQ^{-1}U) = 0, \qquad \text{since } QQ^{-1} = \mathrm{id},$$
$$\nabla_{\dot{\gamma}}^{\mathrm{En}}(Q)Q^{-1}U + Q\nabla_{\dot{\gamma}}^{\mathrm{En}}(Q^{-1}U) + \Phi(QQ^{-1}U) = 0, \qquad \text{product rule},$$
$$Q^{-1}\nabla_{\dot{\gamma}}^{\mathrm{En}}(Q)\widetilde{U} + \nabla_{\dot{\gamma}}^{\mathrm{En}}\widetilde{U} + (Q^{-1}\Phi Q)\widetilde{U} = 0, \qquad \text{multiply by } Q^{-1} \text{ and set } \widetilde{U} = Q^{-1}U,$$
$$\widetilde{\nabla}_{v}^{\mathrm{En}}\widetilde{U} + \widetilde{\Phi}\widetilde{U} = 0,$$

where in the last step we set  $\widetilde{\nabla}_v^{\text{En}}\widetilde{U} := \nabla_v^{\text{En}}\widetilde{U} + Q^{-1}\nabla_v^{\text{En}}(Q)\widetilde{U}$  and  $\widetilde{\Phi} := Q^{-1}\Phi Q$ . It's quick to see that  $\widetilde{\nabla}_v^{\text{En}}$  and  $\widetilde{\Phi}$  will generate the same data set (2.4) if  $Q|_{\partial \overline{M}} =$  id and, that this Q satisfies (2.8). The main point of Theorem 2.7 is that this is the only way that we can produce another pair  $\widetilde{\nabla}^{\mathcal{E}}$  and  $\widetilde{\Phi}$  with the same data set when the Higgs fields are skew symmetric. This example is illustrative in the sense that from the above we see that the gauge matrix Q is given by

$$Q = U\widetilde{U}^{-1}. (2.12)$$

In particular, the theorem is proved by taking the solution U to (2.11) and the solution  $\tilde{U}$ when  $\nabla^{\mathcal{E}}$  and  $\Phi$  are replaced by  $\tilde{\nabla}^{\mathcal{E}}$  and  $\tilde{\Phi}$  respectively, and showing that (2.12) satisfies the desired gauge relation (2.8). To elaborate more on the outline of the proof, we will show that Q-id satisfies a transport equation of the specific form

$$\nabla_X^{\pi^* \text{En}} W + \Psi W = f \tag{2.13}$$

on the sphere bundle SM where X is the geodesic vector field (c.f. (5.83) below). The right-hand side f will be sufficiently regular at infinity and have Fourier modes of order no bigger than one with respect to the vertical Laplacian. In Section 5.4 we will prove a regularity theorem for transport equations that will imply that the solutions W is also sufficiently regular at infinity. In Section 5.3 we will conduct a Fourier study of transport equations that, combined with the just mentioned result, will imply that W, and hence Q, have Fourier degree zero (i.e. are of the form  $C^{\infty}(\overline{M}; \operatorname{End} \mathcal{E})$ ). From there it will quickly follow that Q satisfies the conclusions in Theorem 2.7.

One of the key steps in our investigation will be to generalize an identity called the "Pestov identity" to vector bundles over AH spaces (i.e. Theorem 5.7 below). This is similar to the Pestov identity for scalar functions that was generalized to nontrapping AH spaces in [14] (see bottom of page 2892 there) which the authors accomplished by also generalizing Santaló's formula to AH spaces which required the additional assumption that the manifold is nontrapping. We will take a different approach which will allow us to not assume that the manifold nontrapping to prove this intermediate step. Of course, this will be more than we need since in our final result Theorem 2.7 we do assume that (M, g) is nontrapping.

Another key feature of this paper is the formulation of regularity spaces to which solutions of transport equations over SM of the form (2.13) belong to and to which we can apply vertical differential operators while maintaining sufficient regularity at infinity. We've chosen the regularity spaces  $\mathcal{R}^k$  that we introduce in Section 5.1 below which will play a similar role to Sobolev spaces in our paper. The approach that we take to study the existence and regularity of solutions to transport equations (see Proposition 5.54) is to embed SM as a subset of the *b*-cosphere bundle  ${}^{b}S^*\overline{M}$  and the 0-cosphere bundle  ${}^{0}S^*\overline{M}$  (defined in Section 4.2). The bundle  ${}^{b}S^*\overline{M}$  provides a natural setting to prove the existence of solutions to transport equations with boundary conditions at infinity due to the nice behavior of the geodesic vector field X at the boundary  $\partial^{b}S^*\overline{M}$ . The bundle  ${}^{0}S^*\overline{M}$  on the other hand provides a convenient domain to prove the regularity of solutions at infinity due to its compactness.

### 2.4 Existence of Examples

Here we address the question of the existence of manifolds (M, g) and connections  $\nabla^{\mathcal{E}}$  that satisfy the assumptions of our main results Theorem 2.7 and Corollary 2.10. The existence of a nontrapping AH space (M, g) with negative sectional curvature is provided by the Poincaré ball model. More examples can be produced by (smoothly) deforming the metric slightly in small enough regions; we provide a brief argument for this.

Suppose that our AH space (M, g) is nontrapping and has negative sectional curvature. It's clear that if one takes any open  $U \subseteq M$  whose closure is compact and contained in the domain of local coordinates, then small enough deformations of the metric g over Uwith respect to the coordinates'  $C^2$ -norm will preserve its negative curvature. So, let us demonstrate that if this region and deformation are small enough, then we can also preserve the nontrapping property of (M, g). Taking our boundary defining function  $\rho$ , it follows from Lemma 2.3 in [14] that there exists an  $\varepsilon > 0$  such that if a g-geodesic  $\gamma$  makes its way into the region  $\{\rho < \varepsilon\}$  at some time  $t_0$ , then it will stay in  $\{\rho < \varepsilon\}$  for all  $t > t_0$  and will escape to infinity (i.e.  $\liminf_{t\to\infty} \rho(\gamma(t)) = 0$ ). Now, let us suppose that our U above is outside of this region:  $U \subseteq \{\rho \ge \varepsilon\}$ . Let  $\theta_g$  be the flow of the geodesic vector field  $X_g$  over TM (we wrote "g" here for emphasis). The continuous dependence of solutions to ordinary differential equations on parameters (e.g. see Theorem 7.4 of Chapter 1 in [5]) implies that the map  $\theta_g$  depends continuously on the values of g and its first and second order partials in U. By assumption g is nontrapping, and hence all of its geodesics coming out of U will eventually make their way to  $\{\rho < \varepsilon\}$  and escape to infinity. If we make U precompact in coordinates, it's not hard to see then that small enough deformations of the metric g over U with respect to the  $C^2$ -norm will also satisfy that all of their geodesics coming out of U will eventually make their way to  $\{\rho < \varepsilon\}$ , and hence such deformations will be nontrapping. This proves our claim.

Finally, the following result address the question of the existence of connections  $\nabla^{\mathcal{E}}$  on such AH spaces that satisfy the assumption of the nonexistence of nontrivial twisted CKTs in  $\mathcal{R}^3(SM; \pi^*\mathcal{E})$ . Similar results can be found in [39, Corollary 3.6] and [7, Theorem 1.6]. We refer the reader to Section 4.6 for the definition of the curvature operator  $F^{\mathcal{E}}$  of  $\nabla^{\mathcal{E}}$ .

**Theorem 2.14.** Assume that (M, g) is nontrapping and that the sectional curvatures of g are negative. Then the sectional curvatures are bounded above by  $-\kappa$  for some  $\kappa > 0$ . For any connection  $\nabla^{\mathcal{E}}$  whose curvature norm satisfies  $\|F^{\mathcal{E}}\|_{L^{\infty}} \leq \kappa \sqrt{n}$ , there are no nontrivial twisted CKTs in  $\mathcal{R}^3(SM; \pi^*\mathcal{E})$ .

Similar to the remark made after Corollary 2.10, an example of when this lemma applies is if  $\overline{M}$  is a subset of  $\mathbb{R}^{n+1}$ ,  $\mathcal{E} = \overline{M} \times \mathbb{C}^d$  is the trivial bundle whose sections we write as column vectors, and the connection  $\nabla^{\mathcal{E}}$  is given by  $\nabla_v^{\mathcal{E}} u = [v(u^1) + {}^{\mathcal{E}}\Gamma_{ij}^1 v^i u^j, \ldots, v(u^d) + {}^{\mathcal{E}}\Gamma_{ij}^d v^i u^j]$ where the connection symbols  ${}^{\mathcal{E}}\Gamma_{ij}^k$  decay fast enough at the boundary (i.e.  ${}^{\mathcal{E}}\Gamma_{ij}^k \in \rho^{N_0}C^{\infty}(\overline{M})$ for big enough  $N_0 \ge 0$ ), and such that they and their first partials are small enough so that the curvature estimate in the above lemma is satisfied.

### 2.5 Non-Abelian X-Ray Transform

We mention a way to formalize the operator that takes  $(\nabla, \Phi)$  to the map taking pairs  $(\gamma, e)$ as above to (2.4) using concepts that we introduce in Sections 4.1, 4.2, and 4.5 below. We will not make use of this formulation, and only the material up to (2.15) and the two sentences afterwards here will be used later in the paper. Throughout this section we assume that (M, g) is nontrapping,  $\Phi \in \rho C^{\infty}(\overline{M}; \operatorname{End} \mathcal{E})$ , and that the connection symbols of  $\nabla^{\mathcal{E}}$  are in  $C^{\infty}(\overline{M})$  in any boundary coordinates and frame (in the sense of Definition 2.6).

Consider the cotangent and *b*-cotangent bundles  $T^*M$  and  ${}^{b}T^*\overline{M}$  respectively, and their unit cosphere bundles  $S^*M$  and  ${}^{b}S^*\overline{M}$  respectively. Suppose  $(\rho, y^1, \ldots, y^n)$  are asymptotic boundary normal coordinates of  $\overline{M}$  and consider the frame  $({}^{d\rho}\!/\!\rho, dy^1, \ldots, dy^n)$  spanning covectors in  $T^*M$ . On page 2865 of [14] the authors remind the reader that this extends to the boundary to become a smooth frame of  ${}^{b}T^*\overline{M}$  and that furthermore if  $\zeta = \eta_0 {}^{d\rho}\!/\!\rho + \eta_\mu dy^\mu \in {}^{b}T^*\overline{M}|_{\partial\overline{M}}$  is over the boundary, then the map

$$\eta_0 \frac{d
ho}{
ho} + \eta_\mu dy^\mu \longmapsto \eta_0$$

is well defined (i.e. independent of the coordinates  $(\rho, y^{\mu})$  that we choose). The boundary of the unit cosphere bundle  ${}^{b}S^{*}\overline{M} \subseteq {}^{b}T^{*}\overline{M}$  turns out to have the following two components:

$$\partial_{-}{}^{b}S^{*}\overline{M} = \left\{ \zeta \in {}^{b}T^{*}\overline{M} \big|_{\partial \overline{M}} : \eta_{0} = 1 \right\}$$
 called the "incoming boundary,"  
$$\partial_{+}{}^{b}S^{*}\overline{M} = \left\{ \zeta \in {}^{b}T^{*}\overline{M} \big|_{\partial \overline{M}} : \eta_{0} = -1 \right\}$$
 called the "outgoing boundary."

Let  $\pi : S^*M \to M$  and  $\pi_b : {}^bS^*\overline{M} \to \overline{M}$  denote the natural projection maps. Recall that any unit-speed geodesic  $\gamma : (-\infty, \infty) \to M$  is the image under  $\pi$  of an integral curve  $\sigma : (-\infty, \infty) \to S^*M$  of the geodesic vector field X. Letting  $X_b$  denote the pushforward of X onto  ${}^bS^*\overline{M}|_M$  via the canonical identification between  $T^*M$  and  ${}^bT^*\overline{M}|_M$ , we have that  $\gamma$  is the image under  $\pi_b$  of an integral curve  $\sigma_b : (-\infty, \infty) \to {}^bS^*\overline{M}|_M$  of  $X_b$ . It follows from the proof of Corollary 2.5 in [14] that the limit of any such curve exists in  ${}^bS^*\overline{M}$ :

$$\lim_{t \to -\infty} \sigma_b = \partial_{-}{}^b S^* \overline{M},$$

$$\lim_{t \to \infty} \sigma_b = \partial_{+}{}^b S^* \overline{M}.$$
(2.15)

Intuitively, the first limit here can be thought of as the "initial velocity" of the geodesic as it "enters" the AH space at infinity, while the second its "exit velocity" as it "leaves" at infinity. Conversely, it follows from the same proof that every  $\zeta \in \partial_{-}{}^{b}S^{*}\overline{M}$  (resp.  $\zeta \in \partial_{+}{}^{b}S^{*}\overline{M}$ ) is the limit in  ${}^{b}S^{*}\overline{M}$  of a unique (up to reparameterization) such curve  $\sigma_{b}$  as  $t \to -\infty$  (resp.  $t \to \infty$ ).

Hence we may define the map

$$T^{(\nabla^{\mathcal{E}},\Phi)}: \pi^*\mathcal{E}|_{\partial_{-}{}^bS^*\overline{M}} \longrightarrow \pi^*\mathcal{E}|_{\partial_{+}{}^bS^*\overline{M}}$$

as follows. Take any  $e \in \pi^* \mathcal{E}|_{\partial_{-}{}^{b}S^*\overline{M}}$  whose base point we denote by  $\zeta \in \partial_{-}{}^{b}S^*\overline{M}$ . Let  $\sigma_b$  be an integral curve of  $X_b$  with  $\zeta = \lim_{t \to -\infty} \sigma_b$  and let  $\zeta_{\text{exit}} = \lim_{t \to \infty} \sigma_b$ . Take the geodesic  $\gamma = \pi_b \circ \sigma_b$  and let u be the solution to (2.3) where we let e also denote the element in  $\mathcal{E}_{\pi_b(\zeta)}$  that's canonically identified to  $e \in (\pi^* \mathcal{E})_{\zeta}$ . Then we set

$$T^{(\nabla^{\mathcal{E}},\Phi)}(e) = \lim_{t \to \infty} u(t)$$

making the similar canonical identification  $(\pi^* \mathcal{E})_{\zeta_{\text{exit}}} = \mathcal{E}_{\pi_b(\zeta_{\text{exit}})}$ . We point out that this limit exists by Lemma 2.5, and that the "T" here stands for "transport equation."

**Definition 2.16.** Over the set of all  $(\nabla^{\mathcal{E}}, \Phi)$  such that  $\Phi$  satisfies the decay conditions in Lemma 2.5 above, the operator

$$(\nabla^{\mathcal{E}}, \Phi) \longmapsto T^{(\nabla^{\mathcal{E}}, \Phi)}$$

is called the **non-abelian X-ray transform**.

For example, another way to formulate Corollary 2.10 above is that for any g and  $\nabla^{\mathcal{E}}$  satisfying the conditions there, the non-abelian X-ray transform is injective over the set of all Higgs field satisfying the decay condition also described there.

#### 2.6 Prior Research Discussion and Applications

A standard approach for studying injectivity properties of X-ray transforms is via energy identities that was first introduced in [33]. The type of energy estimate that's used in this approach is called the **Pestov identity** (or **Muhometov-Pestov identity**) which over the years has taken many forms as authors apply them in various contexts - see for instance [11], [38], [41], and [42]. The mentioned paper [11] furthermore explains the connection between X-ray transform over connections and inverse problems related to the wave equation. Of recent works, in dimension two the authors of [37] used a Pestov identity to prove solenoidal injectivity of the X-ray transform over tensors, and in their earlier work [36] they proved a "Pestov type identity" to study the attenuated ray transform with a connection and Higgs field.

The paper [39] proceeded to generalize these methods to manifolds of dimensions greater than two, but it didn't cover the case of connections. In [16] the authors generalized the setup in [39] where they studied the X-ray transform for connections and Higgs fields together. For instance, Theorem 2.7 above was proved in [16] in the case of when (M, g) is a compact Riemannian manifold, has strictly convex boundary, has negative sectional curvature, the boundary condition in (2.3) is changed to  $u(\gamma(a))$  where  $\gamma : [a, b] \to M$  is a unit-speed geodesic traveling between boundary points, and (2.4) is changed to "recording"  $u(\gamma(b))$ . In this paper we also generalize the Pestov identity proved in [16] to AH spaces, of which a similar formula also appears in [43].

We mention the early work [1] that studied the injectivity of the X-ray transform for oneforms. The work of [45] studied injectivity on tensor fields of rank  $m \leq 2$  for analytic simple metrics and a generic class of two-dimensional simple metrics, and proved a stability estimate for the normal operator. Later [44] proved injectivity on two-tensors for all two-dimensional simple metrics which was then extended to tensors of all rank in [37]. The papers [50] and [48] proved injectivity for functions and two-tensors respectively on Riemannian manifolds that admit convex foliations. The paper [15] proved injectivity on tensors of all ranks over Riemannian manifolds with negative curvature and strictly convex boundary. We mention that the work [4] characterized the range of the non-abelian X-ray transform on simple surfaces in terms of boundary quantities and that [3] and [31] proved stability estimates for it over Higgs fields. Microlocal techniques have also been applied to the study of the X-ray transform in the presence of conjugate points - we refer the reader to the works [21], [32], [46], and [47].

In the noncompact realm, injectivity for the scalar X-ray transform over hyperbolic spaces was proved in [19], and inversion formulas are given in [2] and [18]. In [25] the authors proved analogous injectivity over Cartan-Hadamard manifolds and in [26] their results were extended to higher dimensions and tensor fields. The paper [35] proved a gauge equivalence for the X-ray transform for connections on Euclidean space assuming a bound on the size of the connection in dimension two.

AH manifolds have gained interest in the past two decades partly due to their role in physics such as the AdS/CFT conjecture made in [27]. The work [6] for instance describes the role of integral geometry in the AdS/CFT correspondence. In this setting, the paper [14] proved injectivity of the X-ray transform for tensor of all orders on asymptotically hyperbolic spaces. On simple AH manifolds, the work [9] generalized their result for the scalar X-ray

transform by proving a stability estimate for the normal operator. Analogous to the local problem studied in [50], [10] proved a local injectivity result for the scalar X-ray transform on AH spaces.

Regarding applications of the non-abelian X-ray transform, we also mention its appearance in the theory of solitons when studying the Bogomolny equations in dimensions 2 + 1- see [28] and [51] for details. The paper [22] describes its applications to coherent quantum tomography. For a survey of the non-abelian X-ray transform and to read more about its applications, we refer the reader to [34].

#### 2.7 Acknowledgments

First of all, I would like to thank my advisor Gunther Uhlmann who from the beginning of my PhD studies has always supported and encouraged my curiosities, mathematical development, and growth as a human being and mathematician. I'm grateful for both his patience and the invaluable advice that he gave me during times of both progress and confusion, including identifying key references for inspiration on how to proceed forward whenever faced with difficulties. I would also like to thank Robin Graham for helping me navigate my graduate career and having many fascinating mathematical discussions with me over the years. I would like to thank Kelvin Lam for our regular meetings in which we discussed our work.

I'm grateful to everyone in the math department at the University of Washington for creating a welcoming and productive environment in which I could study. The author was partly supported by the NSF grant # 2105956.

# 3 Well-Definedness of the Non-abelian X-Ray Transform

In this section we prove Lemma 2.5. Take any complete geodesic  $\gamma : (-\infty, \infty) \to M$  and any  $e \in \mathcal{E}_{x_0}$  where  $x_0$  is the limit of  $\gamma(t)$  in  $\overline{M}$  as  $t \to -\infty$ . Our plan is to

- 1. prove the existence and uniqueness of the solution to the initial value problem (2.3), on an interval of the form  $(-\infty, t_0]$  for some  $t_0 \in \mathbb{R}$ ,
- 2. argue the existence and uniqueness on the rest of the interval  $[t_0, \infty)$  (and hence everywhere),
- 3. and finally prove that the limit (2.4) exists.

We begin with 1), which we prove by mapping the infinite interval to a bounded one and then applying standard existence and uniqueness results of ordinary differential equations (ODEs). Let  $(\rho, y^1, \ldots, y^n) = (x^i)$  be asymptotic boundary normal coordinates of  $\overline{M}$ containing  $x_0$  in their domain and let  $(b_j)$  be a frame for  $\mathcal{E}$  over the same domain. Let  ${}^{\mathcal{E}}\Gamma_{ij}^k$  denote the connection symbols of  $\nabla^{\mathcal{E}}$  with respect to  $(\partial/\partial x^i)$  and  $(b_k)$ . Let  $t_0 \in \mathbb{R}$  be a time such that the image of  $\gamma$  is contained in these coordinates for all times  $t \in (-\infty, t_0]$ . Then, writing  $u = u^k b_k$ , in these coordinates for  $t \in (-\infty, t_0]$  we have that (2.3) becomes the following system of ODEs

$$\frac{du^k}{dt} + {}^{\mathcal{E}}\Gamma^k_{ij}\dot{\gamma}^i u^j + \Phi^k_i u^i = 0, \qquad \lim_{t \to -\infty} u^k = e^k, \qquad k \in \{1, \dots, d\}, \qquad (3.1)$$

Let's look at the growth rate of the  $\dot{\gamma}^i$ 's. By definition,  $g = \overline{g}/\rho^2$  for some smooth metric  $\overline{g}$  on  $\overline{M}$ . Since  $\gamma$  has a constant speed one, we have that

$$\overline{g}_{ij}\dot{\gamma}^i\dot{\gamma}^j = \rho^2.$$

Clearly the image  $\gamma(-\infty, t_0]$  is a compact subset of our coordinates' domain, and so the matrix in the bilinear form  $v \mapsto \overline{g}_{ij}v^iv^j$  has a minimum positive eigenvalue along this set. Hence from the above we get that there exists a C > 0 such that each  $|\dot{\gamma}^i| \leq C\rho$ .

Now, take the diffeomorphism  $h: (-\pi/2, s_0] \to (-\infty, t_0]$  given by  $h(s) = \tan s$ . Making the change of variables t = h(s) in (3.1) gives that for each  $k \in \{1, \ldots, d\}$ 

$$\frac{du^k}{ds} + {}^{\mathcal{E}}\Gamma^k_{ij}\dot{\gamma}^i \frac{dh}{ds} u^j + \Phi^k_i \frac{dh}{ds} u^i = 0 \quad \text{on } s \in (-\pi/2, s_0), \qquad \qquad u^k(-\pi/2) = e^k$$

In other words, the existence and uniqueness of a continuous solution u to this system of initial value problems will prove 1). This in turn will follow from standard results on linear ODEs (see for instance [5]) if we show that the above coefficients  ${}^{\mathcal{E}}\Gamma^k_{ij}\dot{\gamma}^i h'$  and  $\Phi^k_i h'$  extend continuously to  $s = -\pi/2$ .

It follows from Lemma 2.3 in [14] (specifically (2.11) there) that there exists a constant C' > 0 such that for  $t \in (-\infty, t_0]$ ,

$$\rho \circ \gamma(t) < C'e^t. \tag{3.2}$$

Since by assumption  ${}^{\mathcal{E}}\Gamma_{ij}^k \in C^{\infty}(\overline{M})$ ,  $|\dot{\gamma}^i| \leq C\rho$ , and  $\Phi_i^k \in \rho C^{\infty}(\overline{M})$ , we have that there exists a constant C'' > 0 such that for  $s \in (-\pi/2, s_0]$  both  $|{}^{\mathcal{E}}\Gamma_{ij}^k \dot{\gamma}^i h'|$  and  $|\Phi_i^k h'|$  are bounded above by

$$C''(\rho \circ \gamma(h(s)))h'(s) \le C''C'e^{\tan(s)}\sec^2(s) \to 0 \quad \text{as } s \to -\pi/2^+$$

Hence indeed  ${}^{\mathcal{E}}\Gamma^k_{ij}\dot{\gamma}^i h'$  and  $\Phi^k_i h'$  extend continuously to  $s = -\pi/2$ .

Item 2) follows by applying standard existence and uniqueness theory of ODEs in coordinates and frames of  $\mathcal{E}$  as one travels along the geodesic. Item 3) is proved similarly to 1) except one uses Lemma 2.3 in [14] in forward time (for instance, the  $e^t$  in (3.2) will change to  $e^{-t}$ ).

## 4 Geometric Preliminaries

#### 4.1 The b and 0 Cotangent Bundles

In this section we introduce the b and 0 cotangent bundles. We will only state their definitions and properties, referring the reader to Section 2.2 in [30] for more details. We begin by recalling that lowering and raising an index with respect to g provides a bundle isomorphism between the tangent and cotangent bundles over the interior:

$$\flat: TM \longrightarrow T^*M, \qquad \qquad \sharp: T^*M \longrightarrow TM.$$

The first bundle that we introduce is the *b***-tangent bundle** " ${}^{b}T\overline{M}$ ," which comes with a canonical smooth map  $F: {}^{b}T\overline{M} \to T\overline{M}$  that has the following two properties:

- 1. F is a bijection between smooth sections of  ${}^{b}T\overline{M}$  and smooth sections of  $T\overline{M}$  that are tangent to the boundary  $\partial \overline{M}$ .
- 2. For any fixed point  $x \in \overline{M}$ , F restricts to a linear homomorphism  $F_x : {}^{b}T_x\overline{M} \to T_x\overline{M}$  that is also an isomorphism when x is in the interior M.

The second is the **0-tangent bundle** " $^{0}T\overline{M}$ ," which is defined similarly as coming with a smooth map  $H: {}^{0}T\overline{M} \to T\overline{M}$  that has the following two properties:

- 1. *H* is a bijection between smooth sections of  ${}^{0}T\overline{M}$  and smooth sections of  $T\overline{M}$  that vanish at the boundary  $\partial \overline{M}$ .
- 2. For any fixed point  $x \in \overline{M}$ , H restricts to a linear homomorphism  $H_x : {}^{0}T_x\overline{M} \to T_x\overline{M}$  that is also an isomorphism when x is in the interior M.

Of more importance to us will be the dual bundles  ${}^{b}T^{*}\overline{M}$  and  ${}^{0}T^{*}\overline{M}$ , which are called the **b** and **0** cotangent bundles respectively. They naturally generate pullback maps  $F^{*}: T^{*}\overline{M} \to {}^{b}T^{*}\overline{M}$  and  $H^{*}: T^{*}\overline{M} \to {}^{0}T^{*}\overline{M}$  which are also bundle homomorphisms that are isomorphisms on fibers over the interior M (c.f. points 2) above)

**Remark 4.1.** Considering that  $\flat, \sharp, F, H, F^*, H^*$  are all isomorphisms (on fibers) over the interior M, we will often identify two points in TM,  $T^*M$ ,  ${}^bT\overline{M}|_M$ ,  ${}^bT^*\overline{M}|_M$ ,  ${}^0T\overline{M}|_M$ , and  ${}^0T^*\overline{M}|_M$  as being the same if it's possible to go from one to the other by a composition of the "canonical identification" maps mentioned above.

We mention important frames for the b and 0 cotangent bundles near the boundary  $\partial M$ . Suppose  $(\rho, y^1, \ldots, y^n) = (x^i)$  are boundary coordinates of  $\overline{M}$ . Then it turns out that

$$F^*\left(\frac{d\rho}{\rho}\right), F^*\left(dy^1\right), \dots, F^*\left(dy^n\right)$$
$$H^*\left(\frac{dx^0}{\rho}\right), \dots, H^*\left(\frac{dx^n}{\rho}\right)$$

extend smoothly to the boundary  $\partial M$  to be frames of  ${}^{b}T^{*}\overline{M}$  and  ${}^{0}T^{*}\overline{M}$  respectively. It's standard to abuse notation by simply writing that  ${}^{d\rho/\rho}, dy^{1}, \ldots, dy^{n}$  and  ${}^{dx^{0}/\rho}, \ldots, {}^{dx^{n}/\rho}$  are frames for  ${}^{b}T^{*}\overline{M}$  and  ${}^{0}T^{*}\overline{M}$  respectively. Hence we often write coordinates of  ${}^{b}T^{*}\overline{M}$  and  ${}^{0}T^{*}\overline{M}$  as  $\eta_{0}{}^{d\rho/\rho} + \eta_{\mu}dy^{\mu} \mapsto (\rho, y^{\mu}, \eta_{0}, \eta_{\mu})$  and  $\overline{\eta}_{i}{}^{dx^{i}/\rho} \mapsto (x^{i}, \overline{\eta}_{i})$  respectively. For example, if we consider the coordinates  $v^{i\partial/\partial x^{i}} \mapsto (x^{i}, v^{i})$  of TM, then the canonical identification  $H^{*} \circ \flat : TM|_{M} \to {}^{0}T^{*}\overline{M}$  is given by  $v^{i\partial/\partial x^{i}} \mapsto (\rho g_{ij}v^{i}){}^{dx^{j}/\rho}$ .

### 4.2 The b and 0 Cosphere Bundles

Suppose  $(\rho, y^1, \ldots, y^n)$  are asymptotic boundary normal coordinates of  $\overline{M}$  as described in Section 2.2 above. We know that  $T^*M$  has a fiber metric g. Thus the maps  $F^*: T^*\overline{M} \to {}^{b}T^*\overline{M}$  and  $H^*: T^*\overline{M} \to {}^{0}T^*\overline{M}$  push g to become fiber metrics on  ${}^{b}T^*\overline{M}|_{M}$  and  ${}^{0}T^*\overline{M}|_{M}$ , which we denote by  $g_b$  and  $g_0$  respectively. If we consider the boundary frames for  ${}^{b}T^*\overline{M}$ and  ${}^{0}T^*\overline{M}$  introduced at the end of Section 4.1 above, we have that these metrics are given by (c.f. (2.2))

$$\left| \eta_0 \frac{d\rho}{\rho} + \eta_\mu dy^\mu \right|_{g_b}^2 = \eta_0^2 + \rho^2 h^{\mu\nu} \eta_\mu \eta_\nu,$$

$$\left| \overline{\eta}_0 \frac{d\rho}{\rho} + \overline{\eta}_\mu \frac{dy^\mu}{\rho} \right|_{g_0}^2 = \overline{\eta}_0^2 + h^{\mu\nu} \overline{\eta}_\mu \overline{\eta}_\nu,$$

$$(4.2)$$

where  $(h^{\mu\nu})$  denotes the inverse matrix of  $(h_{\mu\nu})$ . From here we see that both  $g_b$  and  $g_0$  extend smoothly to all of  ${}^{b}T^*\overline{M}$  and  ${}^{0}T^*\overline{M}$  respectively. This allows us to define the **unit cosphere bundles** in  ${}^{b}T^*\overline{M}$  and  ${}^{0}T^*\overline{M}$ :

$${}^{b}S^{*}\overline{M} = \left\{ \zeta \in {}^{b}T^{*}\overline{M} : |\zeta|_{{}^{b}g} = 1 \right\},$$
$${}^{0}S^{*}\overline{M} = \left\{ \overline{\zeta} \in {}^{b}T^{*}\overline{M} : |\overline{\zeta}|_{{}^{0}g} = 1 \right\}.$$

We note that in [14], they use the notation " $\overline{S^*M}$ " for what we denote by " ${}^{b}S^*\overline{M}$ ." We let  $\pi_b: {}^{b}S^*\overline{M} \to \overline{M}$  and  $\pi_0: {}^{0}S^*\overline{M} \to \overline{M}$  denote the natural projection maps.

**Remark 4.3.** Similarly to the remark made in Remark 4.1, we will often identify two points in SM,  $S^*M$ ,  ${}^{b}S^*\overline{M}$ , and  ${}^{0}S^*\overline{M}$  to be the same if it's possible to go from one to the other by a composition of the maps mentioned there.

We point out that it's easy to see that both  ${}^{b}S^{*}\overline{M}$  and  ${}^{0}S^{*}\overline{M}$  are smooth embedded submanifolds with boundary of  ${}^{b}T^{*}\overline{M}$  and  ${}^{0}T^{*}\overline{M}$ . We also note that by (4.2),  $g_{b}$  degenerates over  $\partial \overline{M}$  (i.e. stops being positive definite) while  $g_{0}$  does not. In particular this implies that  ${}^{b}S^{*}\overline{M}$  is not compact while  ${}^{0}S^{*}\overline{M}$  is compact.

#### 4.3 Splitting the Tangent Bundle

Next we define a natural Riemannian metric on the tangent space TM, called the **Sasaki metric**, generated by g. We recommend that when checking many of the claims below, to check them above the center of normal coordinates since in many cases the expressions simplify considerably due to the vanishing of the Christoffel symbols and the first order partials of g. Consider the tangent bundle's projection map  $\pi : TM \to M$  and its differential  $d\pi : TTM \to TM$ . There is another natural map between these tangent spaces called the **connection map**:  $\mathcal{K} : TTM \to TM$ , which is defined as follows. Take any  $\omega \in T_vTM$  and let  $\alpha : (a, b) \to M$  be a smooth curve and  $V : (a, b) \to TM$  a smooth vector field along  $\alpha$ such that  $(\alpha, V)'(0) = \omega$ . Then we set  $\mathcal{K}(\omega)$  to be the covariant derivative

$$\mathcal{K}(\omega) := \frac{DV}{dt}(0).$$

To check that this is independent of the  $\alpha$  and V that we choose, a quick computation shows that taking coordinates  $(x^i)$  of M and the coordinates  $v^{i\partial}/\partial x^i \mapsto (x^i, v^i)$  of TM,  $\mathcal{K}$  is given by

$$\mathcal{K}\left(\alpha^{i}\frac{\partial}{\partial x^{i}}\Big|_{v^{i}\partial/\partial x^{i}}+\beta^{i}\frac{\partial}{\partial v^{i}}\Big|_{v^{i}\partial/\partial x^{i}}\right)=\left(\beta^{k}+\Gamma_{ij}^{k}\alpha^{i}v^{j}\right)\frac{\partial}{\partial x^{k}},$$

where  $\Gamma_{ij}^k$  are the Christoffel symbols of g with respect to  $(\partial/\partial x^i)$ . Next, an easy exercise shows that the kernels of  $d\pi$  and  $\mathcal{K}$  partition the tangent bundle's tangent space at any  $v \in T_x M$ :

$$T_v T M = \mathcal{H}_v \oplus \mathcal{V}_v \tag{4.4}$$

where

$$\widetilde{\mathcal{H}}_v = \ker \mathcal{K}|_{T_vTM}$$
 and  $\widetilde{\mathcal{V}}_v = \ker d\pi|_{T_vTM}$ 

The " $\widetilde{\mathcal{V}}$ " stands for "vertical" because it can be imagined as being a tangent subspace at v standing vertically above x, while the " $\widetilde{\mathcal{H}}$ " stands for "horizontal." As one can check, both spaces are canonically identified (i.e. isomorphically mapped to) with  $T_x M$  by the restricted maps

$$d\pi: \widetilde{\mathcal{H}}_v \longrightarrow T_x M,$$
$$\mathcal{K}: \widetilde{\mathcal{V}}_v \longrightarrow T_x M.$$

With this splitting in hand, the Sasaki metric G on TM is defined as follows: for any  $\omega, \varsigma \in T_v TM$ ,

$$\langle \omega, \varsigma \rangle_G = \langle d\pi(\omega), d\pi(\varsigma) \rangle_g + \langle \mathcal{K}(\omega), \mathcal{K}(\varsigma) \rangle_g.$$

It follows immediately that (4.4) is an orthogonal decomposition with respect to G.

We will only work with unit speed geodesics and hence most of our work will be done on the unit sphere bundle

$$SM = \Big\{ v \in TM : |v|_g = 1 \Big\}.$$

With a choice of unit normal, the Sasaki metric on TM induces a metric on SM which we'll also call the Sasaki metric and denote by G, relying on context to differentiate the two. It's not hard to see that at any  $v \in SM$  the tangent space of SM splits into the form

$$T_v SM = \mathcal{H}_v \oplus \mathcal{V}_v$$

where  $\mathcal{V}_v$  is the subspace of  $\widetilde{\mathcal{V}}_v$  that's *G*-perpendicular to the unit normals to the "sphere"  $S_x M$  above *x*. Now, take the geodesic vector field *X* over *SM*. It's easy to check that *X* always lies in  $\widetilde{\mathcal{H}}_v$  for all  $v \in SM$  and hence we obtain the splitting

$$T_v SM = (\mathbb{R}X_v) \oplus \mathcal{H}_v \oplus \mathcal{V}_v \tag{4.5}$$

where  $\mathcal{H}_v$  denotes the orthogonal complement of  $\mathbb{R}X_v$  in  $\mathcal{H}_v$ . For future use, we point out that restrictions of  $d\pi$  and  $\mathcal{K}$  map

$$\begin{aligned} d\pi : \mathcal{H}_v &\longrightarrow \left\{ v^{\perp} \right\} \subseteq T_x M, \\ \mathcal{K} : \mathcal{V}_v &\longrightarrow \left\{ v^{\perp} \right\} \subseteq T_x M. \end{aligned}$$
 (4.6)

#### 4.4 Integration on the (Co)Sphere Bundle

Since SM has a Riemannian metric G, it has a Riemannian density and hence the Lebesgue measure generated by it (the latter two are independent of orientation). Hence we may perform Lebesgue integration on SM with respect to G. If  $(x^i)$  are local coordinates of M,  $(r_i)$  is a frame of TM, and we take the coordinates  $v^i r_i \mapsto (x^i, v^i)$  of TM, then it turns out that the integral of any function  $f \in L^1(SM)$  supported over our coordinates is given by the iterated integral

$$\int f = \int f(x^1, \dots, x^n, v^1, \dots, v^n) dS_x(v^1, \dots, v^n) \sqrt{\det g} dx^0 \dots dx^n,$$
(4.7)

where  $(v^1, \ldots, v^n)$  are on the sphere  $|v|_g^2 = 1$  and  $dS_x$  is the Lebesgue measure on  $S_xM$ induced by  $T_xM$  with inner product  $g_x$ . We refer the reader to Section 3.6.2 in [40] for a proof. We point out that the (total) measure of  $S_xM$  is the Euclidean surface area of the Euclidean *n*-sphere for all  $x \in M$ , which for instance follows by looking at the center of normal coordinates.

One example of the usefulness of this observation is that since in boundary coordinates  $\sqrt{\det g}$  is  $\rho^{-(n+1)}$  times "something smooth" on  $\overline{M}$  we have that any function of the form  $\rho^{n+1}L^{\infty}(SM)$  is integrable.

### 4.5 Splitting the Connection Over the Unit Tangent Bundle

Let us take the natural projection map  $\pi : SM \to M$ . The pullback (vector) bundle  $\pi^* \mathcal{E}$ over SM is defined as the set obtained by taking any point  $x \in M$  and attaching a copy of  $\mathcal{E}_x$  to every point of the sphere  $S_xM$  above it. Formally,

$$\pi^* \mathcal{E} := \left\{ (v, e) : v \in SM, e \in \mathcal{E}_{\pi(v)} \right\}$$

We often canonically identify  $(v, e) \cong e$  for fixed v. To every fiber  $(\pi^* \mathcal{E})_v$  we impose the inner product space structure of  $(\mathcal{E}_x, \langle \cdot, \cdot \rangle_{\mathcal{E}_x})$ . If  $(b_i)$  is a smooth frame for  $\mathcal{E}$ , then we turn  $\pi^* \mathcal{E}$  into a smooth vector bundle over SM (with smooth inner product) by declaring<sup>2</sup>  $(\pi^* b_i)$  to be smooth frames for  $\pi^* \mathcal{E}$ . The pullback connection  $\nabla^{\pi^* \mathcal{E}} = \pi^* \nabla^{\mathcal{E}}$  in  $\pi^* \mathcal{E}$  is defined to be the unique connection so that if  $\omega \in TSM$  and  $u_0 : M \to \mathcal{E}$  is smooth, then

$$\nabla_{\omega}^{\pi^*\mathcal{E}}(\pi^*u_0) = \pi^* \big( \nabla_{d\pi(\omega)}^{\mathcal{E}} u_0 \big).$$
(4.8)

For a smooth section  $u = u^j \pi^* b_j : SM \to \pi^* \mathcal{E}$ , the pullback connection is explicitly given by

$$\nabla_{\omega}^{\pi^*\mathcal{E}}u = \omega(u^j)\pi^*b_j + u^j\pi^*(\nabla_{d\pi(\omega)}^{\mathcal{E}}b_j).$$
(4.9)

**Remark 4.10.** In the same way we can define the pullback bundles  $\pi_b^* \mathcal{E}$  and  $\pi_0^* \mathcal{E}$  on  ${}^bS^*\overline{M}$  and  ${}^0S^*\overline{M}$  respectively. In fact, over the interior we will often identify a section  $u \in C^{\infty}(SM; \pi^*\mathcal{E})$  as an element of  $C^{\infty}({}^bS^*\overline{M}|_M, \pi_b^*\mathcal{E}|_M)$  and  $C^{\infty}({}^0S^*\overline{M}|_M, \pi_0^*\mathcal{E}|_M)$  via the natural identification  $(\pi^*e)_v \cong (\pi_b^*e)_{\zeta} \cong (\pi_0^*e)_{\overline{\zeta}} \cong e$  where  $v \cong \zeta \cong \overline{\zeta}$  are identified points on  $S_xM$ ,  ${}^0S^*_x\overline{M}$ , and  ${}^bS^*_x\overline{M}$  respectively.

<sup>&</sup>lt;sup>2</sup>Applying  $\pi^*$  to  $b_i$  means " $b_i \circ \pi$ ."

Having defined the splitting of the unit tangent bundle in (4.5), we now define a natural splitting of the connection of any section  $u: SM \to \pi^* \mathcal{E}$  in the following form:

$$\nabla^{\pi^*\mathcal{E}} u = \nabla^{X} \pi^*\mathcal{E} u + \nabla^{h} \pi^*\mathcal{E} u + \nabla^{v} \pi^*\mathcal{E} u.$$
(4.11)

Let's start by defining  $\nabla^{\pi^*\mathcal{E}} u$ . We have that the full connection  $\nabla^{\pi^*\mathcal{E}} u$  is a tensor of the form  $C^{\infty}(SM; T^*SM \otimes \pi^*\mathcal{E})$ . Now, consider the same tensor but with the first index raised with respect to G:  $[\nabla^{\pi^*\mathcal{E}} u]^{\sharp} \in C^{\infty}(SM; TSM \otimes \pi^*\mathcal{E})$ . Next, it's an easy exercise to check that there exists a unique map

$$P_{\mathcal{H}}: C^{\infty}(SM; TSM \otimes \pi^* \mathcal{E}) \longrightarrow C^{\infty}(SM; TSM \otimes \pi^* \mathcal{E})$$
(4.12)

that satisfies

$$P_{\mathcal{H}}(\omega \otimes e) = (\operatorname{proj}_{\mathcal{H}}\omega) \otimes e$$

where  $\operatorname{proj}_{\mathcal{H}} : T_v TM \to \mathcal{H} \subseteq T_v TM$  is the orthogonal projection map onto  $\mathcal{H}$ . We then define

$$\stackrel{\mathbf{h}}{\nabla}^{\pi^*\mathcal{E}} u := P_{\mathcal{H}}\Big( \big[ \nabla^{\pi^*\mathcal{E}} u \big]^{\sharp} \Big).$$

We define  $\nabla^{X} \pi^* \mathcal{E} u$  and  $\nabla^{v} \pi^* \mathcal{E} u$  in the same way but instead use analogous map  $P_{\mathbb{R}X}$ ,  $\operatorname{proj}_{\mathbb{R}X}$ and  $P_{\mathcal{V}}$ ,  $\operatorname{proj}_{\mathcal{V}}$  respectively. However instead of using  $\nabla^{X} \pi^* \mathcal{E} u$ , it's more common to use the related quantity

$$\mathbb{X}u := \nabla_X^{\pi^*\mathcal{E}}u. \tag{4.13}$$

The quantities Xu and  $\nabla^{\pi^*\mathcal{E}}u$  are equivalent in the sense that knowing one allows you to compute the other. Hence we often record the decomposition (4.11) instead as

$$\nabla^{\pi^*\mathcal{E}}u = \left(\mathbb{X}u, \nabla^{\mathbf{h}}\pi^*\mathcal{E}u, \nabla^{\mathbf{v}}\pi^*\mathcal{E}u\right).$$
(4.14)

The second two components are called the **horizontal and vertical derivatives** of u respectively. However, it's convenient to change the interpretation of the latter two derivatives as follows.

We define the bundle N over SM by attaching to every  $v \in S_x M$  a copy of  $\{v^{\perp}\} \subseteq T_x M$ . Formally,

$$N = \left\{ (v, w) : v \in S_x M \text{ where } x \in M \text{ and } w \in \left\{ v^{\perp} \right\} \right\}.$$

To every fiber  $N_v$  we impose the inner product space structure of  $(\{v^{\perp}\}, g_x)$  which we denote by " $\langle \cdot, \cdot \rangle_{N_v}$ ." It's an easy exercise to show that this is a smooth subbundle of  $\pi^*TM$ . By (4.6) we can think of  $d\pi$  and  $\mathcal{K}$  as mapping

$$\begin{aligned}
d\pi : \mathcal{H}_v &\longrightarrow N_v, \\
\mathcal{K} : \mathcal{V}_v &\longrightarrow N_v.
\end{aligned} \tag{4.15}$$

Hence at every  $v \in TM$ , the maps  $P_{\mathcal{H}}$  and  $P_{\mathcal{V}}$  above map into  $\mathcal{H}_v \otimes (\pi^* \mathcal{E})_v$  and  $\mathcal{V}_v \otimes (\pi^* \mathcal{E})_v$ respectively. Thus using the identification (4.15) we can think of the horizontal and vertical derivatives as both being  $N \otimes \pi^* \mathcal{E}$ -valued:

$$\stackrel{\mathbf{h}}{\nabla}^{\pi^*\mathcal{E}} u \in C^{\infty}(SM; N \otimes \pi^*\mathcal{E}) \quad \text{and} \quad \stackrel{\mathbf{v}}{\nabla}^{\pi^*\mathcal{E}} u \in C^{\infty}(SM; N \otimes \pi^*\mathcal{E}).$$

We mention that we assign the natural inner product on  $N \otimes \pi^* \mathcal{E}$  (i.e. the unique one satisfying  $\langle z \otimes e, z' \otimes e' \rangle_{N \otimes \pi^* \mathcal{E}} = \langle z, z' \rangle_N \otimes \langle e, e' \rangle_{\pi^* \mathcal{E}}$ ). The reason that this interpretation is useful is that it becomes natural to apply well-known adjoint formulas for the horizontal and vertical derivatives over this space. In particular, it turns out that there are differential operators

$$\operatorname{div}^{\operatorname{n}} \pi^{*\mathcal{E}} : C^{\infty}(SM; N \otimes \pi^{*\mathcal{E}}) \longrightarrow C^{\infty}(SM; \pi^{*\mathcal{E}}),$$

$$\operatorname{div}^{\operatorname{v}} \pi^{*\mathcal{E}} : C^{\infty}(SM; N \otimes \pi^{*\mathcal{E}}) \longrightarrow C^{\infty}(SM; \pi^{*\mathcal{E}}),$$

$$(4.16)$$

with the property that if  $u \in C^{\infty}(SM; \pi^* \mathcal{E})$  and  $v \in C^{\infty}(SM; N \otimes \pi^* \mathcal{E})$  are such that at least one of them is of compact support, then

$$\langle \nabla^{\mathfrak{h}^{\pi^{*}\mathcal{E}}} u, v \rangle_{L^{2}(N \otimes \pi^{*}\mathcal{E})} = -\langle u, \operatorname{div}^{\mathfrak{h}} v^{\pi^{*}\mathcal{E}} v \rangle_{L^{2}(\pi^{*}\mathcal{E})} \text{ and } \langle \nabla^{\mathfrak{v}^{\pi^{*}\mathcal{E}}} u, v \rangle_{L^{2}(N \otimes \pi^{*}\mathcal{E})} = -\langle u, \operatorname{div}^{\mathsf{v}} v^{\pi^{*}\mathcal{E}} v \rangle_{L^{2}(\pi^{*}\mathcal{E})}.$$

We will discuss this in more detail in Section 5.2 below. The operators in (4.16) are naturally called the **horizontal and vertical divergences** respectively.

For future reference, we end this section with two more definitions. First, we define the differential operator

$$\mathbb{X}: C^{\infty}(SM; N \otimes \pi^* \mathcal{E}) \to C^{\infty}(SM; N \otimes \pi^* \mathcal{E}),$$

differentiated from the X introduced in (4.13) by context, to be the unique operator satisfying that for any  $Z \otimes b \in C^{\infty}(SM; N \otimes \pi^* \mathcal{E})$ ,

$$\mathbb{X}(Z \otimes b) = X(Z) \otimes b + Z \otimes \mathbb{X}(b), \tag{4.17}$$

where X(Z) at any point  $v \in SM$  denotes the covariant derivative of Z along the unit speed geodesic  $\gamma$  with initial velocity v at time t = 0:

$$X(Z)|_v = \frac{D_{\gamma}Z}{dt}(0). \tag{4.18}$$

Since  $Z \perp \dot{\gamma}$  implies that  $D_{\gamma} Z/dt \perp \dot{\gamma}$ , we see that X indeed maps into smooth sections of  $N \otimes \pi^* \mathcal{E}$ (i.e. not simply into  $\pi^* TM \otimes \pi^* \mathcal{E}$ ).

### 4.6 Curvatures

We now cover the curvature operators of  $\mathcal{E}$ ,  $\pi^* \mathcal{E}$ , and simply the metric g. We start with the first one. The operator  $\nabla^{\mathcal{E}}$  maps between the following spaces of sections:

$$\nabla^{\mathcal{E}}: C^{\infty}(\overline{M}; \mathcal{E}) \longrightarrow C^{\infty}(\overline{M}; T^*\overline{M} \otimes \mathcal{E})$$

Let  $\Lambda^k(T^*\overline{M})$  denote the bundle of covariant alternating k-tensors and let

$$C^{\infty}(\overline{M}; \Lambda^k(T^*\overline{M}) \otimes \mathcal{E})$$

denote the space of smooth sections of  $T^*\overline{M} \otimes \ldots \otimes T^*\overline{M} \otimes \mathcal{E}$  that are alternating in their first k arguments. The operators

$$\nabla^{\mathcal{E}}: C^{\infty}(\overline{M}; \Lambda^{k}(T^{*}\overline{M}) \otimes \mathcal{E}) \longrightarrow C^{\infty}(\overline{M}; \Lambda^{k+1}(T^{*}\overline{M}) \otimes \mathcal{E})$$

are defined to be the unique operators that satisfy that for any  $\theta \in C^{\infty}(\overline{M}; \Lambda^k(T^*\overline{M}))$  and any  $u \in C^{\infty}(M; \mathcal{E})$ 

$$\nabla^{\mathcal{E}}(\theta \otimes u) = d\theta \otimes u + (-1)^k \theta \wedge \nabla^{\mathcal{E}} u,$$

where  $\theta \wedge \nabla^{\mathcal{E}} u$  denotes the wedge-like product:

$$(\theta \wedge \nabla^{\mathcal{E}} u)(v_1, \dots, v_{k+1}, l) = \frac{1}{k! 1!} \sum_{\sigma \in S_{k+1}} \theta(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \nabla^{\mathcal{E}} u(v_{\sigma(k+1)}, l),$$

where  $S_{k+1}$  denotes the set of permutations of k+1 elements.

The **curvature** of  $\nabla^{\mathcal{E}}$  is defined to be

$$f^{\mathcal{E}} := \nabla^{\mathcal{E}} \circ \nabla^{\mathcal{E}} : C^{\infty}(\overline{M}; \mathcal{E}) \longrightarrow C^{\infty}(\overline{M}; \Lambda^{2}(T^{*}\overline{M}) \otimes \mathcal{E}).$$

A straightforward computation shows that in any coordinates  $(x^i)$  of  $\overline{M}$  and any frame  $(b_i)$  of  $\mathcal{E}$ , the curvature  $f^{\mathcal{E}}$  applied to any smooth section  $u \in C^{\infty}(\overline{M}; \mathcal{E})$  is given by

$$f^{\mathcal{E}}u = u^{l} \left( \frac{\partial^{\mathcal{E}}\Gamma_{jl}^{k}}{\partial x^{i}} - {}^{\mathcal{E}}\Gamma_{il}^{m\mathcal{E}}\Gamma_{jm}^{k} - \frac{\partial^{\mathcal{E}}\Gamma_{il}^{k}}{\partial x^{j}} + {}^{\mathcal{E}}\Gamma_{jl}^{m\mathcal{E}}\Gamma_{im}^{k} \right) dx^{i} \otimes dx^{j} \otimes b_{k}.$$
(4.19)

where  ${}^{\mathcal{E}}\Gamma_{ij}^k$  are the connection symbols of  $\nabla^{\mathcal{E}}$  with respect to  $(\partial/\partial x^i)$  and  $(b_k)$ . The resemblance of this tensor to the Riemann curvature tensor is the motivation for the name of  $f^{\mathcal{E}}$ .

Next we define a curvature operator associated to  $f^{\mathcal{E}}$  which acts over SM. Notice that  $f^{\mathcal{E}}$  can be viewed as a  $C^{\infty}(\overline{M}; \Lambda^2(T^*\overline{M}) \otimes \mathcal{E} \otimes \mathcal{E}^*)$  tensor field by thinking of the u in (4.19) as the fourth argument of  $f^{\mathcal{E}}$  (i.e. the argument of  $\mathcal{E}^*$ ). Hence it can also be canonically identified with a map, denoted by the same letter, of the form

$$f^{\mathcal{E}}: C^{\infty}(\overline{M}; T\overline{M}) \times C^{\infty}(\overline{M}; \mathcal{E}) \longrightarrow C^{\infty}(\overline{M}; T^*\overline{M} \otimes \mathcal{E}).$$

In our coordinates and frames it is given by the following: if  $f_{ij}{}^k{}_l$  denotes the tensor component written out in the parenthesis "(...)" in (4.19), then for any  $x \in \overline{M}, v \in T_x \overline{M}$ , and  $e \in \mathcal{E}_x$ ,

$$f_x^{\mathcal{E}}(v,e) = e^l f_{ij}{}^k{}_l v^i dx^j \otimes b_k.$$

$$(4.20)$$

We define the **curvature operator** associated to  $f^{\mathcal{E}}$  to be the map

$$F^{\mathcal{E}}: C^{\infty}(SM; \pi^*\mathcal{E}) \longrightarrow C^{\infty}(SM; N \otimes \pi^*\mathcal{E})$$

given by the following. For any  $x \in M$ ,  $v \in S_x M$ , and  $e \in (\pi^* \mathcal{E})_v$ ,

$$F_v^{\mathcal{E}}(e) := P_{N \otimes \pi^* \mathcal{E}} \left( \left[ f_x^{\mathcal{E}}(v, e) \right]^{\sharp} \right).$$
(4.21)

where  $\sharp$  raises the first index of  $f_x^{\mathcal{E}}(v, e)$  and  $P_{N \otimes \pi^* \mathcal{E}} : \pi^* TM \otimes \pi^* \mathcal{E} \to N \otimes \pi^* \mathcal{E}$  denotes projecting the first component of a tensor product perpendicularly onto the normal bundle (i.e.  $\pi^* TM$  onto N - c.f. (4.12)).

The last curvature quantity that we want to establish notation for is the ordinary curvature of g. Let

$$R: C^{\infty}(M;TM) \times C^{\infty}(M;TM) \times C^{\infty}(M;TM) \longrightarrow C^{\infty}(M;TM)$$

denote the Riemann curvature endomorphism given by

$$R(X,Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z,$$

where  $\nabla$  is the Levi-Civita connection and  $[\cdot, \cdot]$  is the Lie bracket. Recall that R is multilinear over  $C^{\infty}(M)$  and thus has well-defined restrictions to  $(T_x M)^3 \to T_x M$  for any fixed  $x \in M$ . We define operators, denoted by the same letter,

$$R: C^{\infty}(SM; N) \longrightarrow C^{\infty}(SM; N)$$
$$R: C^{\infty}(SM; N \otimes \pi^{*}\mathcal{E}) \longrightarrow C^{\infty}(SM; N \otimes \pi^{*}\mathcal{E})$$

given by the following, for any  $x \in M$ ,  $v \in S_x M$ ,  $w \in N_v$ , and  $e \in (\pi^* \mathcal{E})_v$ ,

 $R_v(w) := R_x(w, v)v$  and  $R_v(w \otimes e) := [R_x(w, v)v] \otimes e.$ 

## 5 Gauge Equivalence of Connections and Higgs fields

In this section we build the necessary tools to prove Theorem 2.7, and then prove it at the end.

#### 5.1 The Regularity Spaces

For the majority of this paper we will be working with the solution to a transport equation over SM. In particular, we will be making use of its  $L^2$  norm after we apply several differential operators to it that extends smoothly to the boundary of the 0-cosphere bundle. For this reason, we will make use of the following regularity spaces:

**Definition 5.1.** For any fixed integer  $k \geq 0$ , we define the spaces  $\mathcal{R}^k(SM; \pi^*\mathcal{E})$  and  $\mathcal{R}^k(SM; N \otimes \pi^*\mathcal{E})$  of order k to be the spaces of smooth sections  $u \in C^{\infty}(SM; \pi^*\mathcal{E})$  and  $w \in C^{\infty}(SM; N \otimes \pi^*\mathcal{E})$  respectively that satisfy the following.

We say that  $u \in \mathcal{R}^k(SM; \pi^*\mathcal{E})$  if for any smooth vector fields  $V_1, ..., V_k \in C^{\infty}({}^0S^*\overline{M}; T^0S^*\overline{M})$ over the 0-cosphere bundle, any frame  $(b_i)$  of  $\mathcal{E}$ , and any compact subset  $K \subseteq \text{dom}(b_i) \subseteq \overline{M}$ ,  $u = u^j \pi^* b_j$  must satisfy that

$$u^{j}, V_{1}u^{j}, V_{2}V_{1}u^{j}, \ldots, V_{k}\ldots V_{1}u^{j}$$
 are all in  $\rho^{(n+1)/2}L^{\infty}[K].$  (5.2)

We say that  $w \in \mathcal{R}^k(SM; N \otimes \pi^* \mathcal{E})$  if for any smooth vector fields  $V_1, ..., V_k \in C^{\infty}({}^0S^*\overline{M}; T^0S^*\overline{M})$ , any boundary coordinates  $(\rho, y^{\mu}) = (x^i)$  or interior coordinates  $(x^i)$  of  $\overline{M}$ , any frame  $(b_i)$  of  $\mathcal{E}$  over these coordinates' domain, w is of the form

$$w = \rho w^{ij} \pi^* \frac{\partial}{\partial x^i} \otimes \pi^* b_j$$

(note the  $\rho$  in front), and satisfies that for any compact subset  $K \subseteq \text{dom}(x^i) \subseteq \overline{M}$ ,

 $w^{ij}, V_1 w^{ij}, V_2 V_1 w^{ij}, \dots, V_k \dots V_1 w^{ij}$  are all in  $\rho^{(n+1)/2} L^{\infty}[K].$  (5.3)

We sometimes omit writing the domain of our space and simply write  $\mathcal{R}^k$  when either the domain is of no importance or clear from context. The power "(n+1)/2" above is chosen to ensure that both u and w as above are in  $L^2$  - see Remark 5.6 below.

**Remark 5.4.** We point out that (5.2) and (5.3) only need to be checked in an atlas of  $\overline{M}$ . To see why, take any pair of coordinates and frames  $(\rho, y^{\mu}) = (x^i), (b_j)$  and  $(\rho, \tilde{y}^{\mu}) = (\tilde{x}^i), (\tilde{b}_j)$  as above whose domains intersect. The coefficients  $\alpha_i^j$  and  $\beta_i^j$  in the transformation laws  $\partial/\partial x^i = \alpha_i^{j\partial}/\partial \tilde{x}^i$  and  $b_i = \beta_i^{j} \tilde{b}_j$  will be smooth on  $\overline{M}$  and hence smooth on  ${}^0S^*\overline{M}$  (when lifted). Thus  $\alpha_i^j$  and  $\beta_i^j$  will be bounded over any compact subset K of  $\overline{M}$ , and the same thing will hold if we apply smooth vectors  $V_1, \ldots, V_k$  to them as in the above definition. From here it quickly follows that (5.2) and (5.3) only need to be checked in an atlas of  $\overline{M}$ .

**Remark 5.5.** Regarding continuity at infinity, from (5.2) it follows that any element of  $\mathcal{R}^{k}(SM; \pi^{*}\mathcal{E})$  extends continuously to  ${}^{0}S^{*}\overline{M}$  if we identify it as an element of  $C^{\infty}({}^{0}S^{*}\overline{M}|_{M}, \pi^{*}\mathcal{E}|_{M})$  (see Remark 4.10) and set it to be zero on the boundary  $\partial^{0}S^{*}\overline{M}$ .

**Remark 5.6.** We make a remark regarding integrability. In Definition 5.1 we have that each  $|\pi^{*\partial}/\partial x^i|_N = |\partial/\partial x^i|_g$  is  $\rho^{-1}$  times something smooth on  $\overline{M}$  (since g is  $\rho^{-2}$  times something smooth on  $\overline{M}$ ). Furthermore, each  $|\pi^*b_j|_{\pi^*\mathcal{E}} = |b_j|_{\mathcal{E}}$  is bounded over K since  $\mathcal{E}$  is a smooth bundle over  $\overline{M}$  by our standing assumption. Hence, it follows that both  $|u|_{\pi^*\mathcal{E}}, |w|_{N\otimes\pi^*\mathcal{E}} \in \rho^{(n+1)/2}L^{\infty}[K]$ . Since  $\overline{M}$  is compact and hence can be covered by a finite collection of such sets K, it follows that  $\mathcal{R}^0 \subseteq L^2$  by the comment made at the end of Section 4.4 above. Furthermore, it follows straight from the definition that  $\mathcal{R}^k \subseteq \mathcal{R}^{k'}$  if  $k \geq k'$ . Hence,  $\mathcal{R}^k \subseteq L^2$  for all  $k \geq 0$ .

All of the differential operator that we will be using below will have the mapping properties  $\mathcal{R}^k \to \mathcal{R}^{k-1}$ . Because of this and the properties described in the above remark, the spaces  $\mathcal{R}^k$  will play the analogous role in our paper as Sobolev spaces in PDE theory.

#### 5.2 Pestov Identity

In this section we prove the following version of the **Pestov Identity** with a connection on asymptotically hyperbolic (AH) spaces. It will be used in the Fourier analysis study of transport equations.

**Theorem 5.7.** Suppose that  $u \in \mathcal{R}^2(SM; \pi^*\mathcal{E})$ . Then<sup>3</sup>

$$\left\| \stackrel{\mathbf{v}}{\nabla} \pi^{*\mathcal{E}} \mathbb{X} u \right\|_{L^{2}}^{2} = \left\| \mathbb{X} \stackrel{\mathbf{v}}{\nabla} \pi^{*\mathcal{E}} u \right\|_{L^{2}}^{2} - \langle R \stackrel{\mathbf{v}}{\nabla} \pi^{*\mathcal{E}} u, \stackrel{\mathbf{v}}{\nabla} \pi^{*\mathcal{E}} u \rangle_{L^{2}} - \langle F^{\mathcal{E}} u, \stackrel{\mathbf{v}}{\nabla} \pi^{*\mathcal{E}} u \rangle_{L^{2}} + n \| \mathbb{X} u \|_{L^{2}}^{2}$$

where  $L^2$  stands for  $L^2(SM; N \otimes \pi^* \mathcal{E})$  in the first four quantities and  $L^2(SM; \pi^* \mathcal{E})$  in the last one.

 $<sup>^{3}</sup>$ Here we're implicitly restricting to the interior so that we may apply the differential operators involved and integrate.

Intuitively speaking, the above Pestov identity studies how the "energy" (i.e.  $L^2$ -norm squared) changes when one switches the order of  $\nabla^{\pi^*\mathcal{E}}$  and X. We require that u is in  $\mathcal{R}^2(SM;\pi^*\mathcal{E})$  to ensure that all of the  $L^2$  norms and inner products in the above equation make sense. The order of the space  $\mathcal{R}^2$  is chosen to be "2" because both in the statement of the theorem and its proof we won't be applying more than two smooth vector fields over the 0-cosphere bundle to the components of u at any one time. The theorem is proved by simply starting with  $\|\nabla^{\pi^*\mathcal{E}} X u\|_{L^2}^2$  and then applying  $L^2$ -adjoint relations and commutator formulas until one arrives at  $\|X\nabla^{\pi^*\mathcal{E}} u\|_{L^2}^2$ . The following lemma provides us with the required set of adjoint relations.

**Lemma 5.8.** The following are true, where  $m, m' \ge 0$  are integers and all  $L^2$  stand for appropriate  $L^2(SM;...)$  spaces.

1. If  $u \in \mathcal{R}^m(SM; \pi^*\mathcal{E})$  with  $m \ge 1$ , then  $\mathbb{X}u \in \mathcal{R}^{m-1}(SM; \pi^*\mathcal{E})$ . Furthermore, if  $w \in \mathcal{R}^{m'}(SM; \pi^*\mathcal{E})$ , then

$$\langle \mathbb{X}u, w \rangle_{L^2} = -\langle u, \mathbb{X}w \rangle_{L^2}.$$

2. If  $u \in \mathcal{R}^m(SM; N \otimes \pi^* \mathcal{E})$  with  $m \ge 1$ , then  $\mathbb{X}u \in \mathcal{R}^{m-1}(SM; N \otimes \pi^* \mathcal{E})$ . Furthermore, if  $w \in \mathcal{R}^{m'}(SM; N \otimes \pi^* \mathcal{E})$ , then

$$\langle \mathbb{X}u, w \rangle_{L^2} = -\langle u, \mathbb{X}w \rangle_{L^2}$$

3. If  $u \in \mathcal{R}^m(SM; \pi^*\mathcal{E})$  and  $w \in \mathcal{R}^{m'}(SM; N \otimes \pi^*\mathcal{E})$  with  $m, m' \geq 1$ , then  $\stackrel{v}{\nabla} \pi^{*\mathcal{E}} u \in \mathcal{R}^{m-1}(SM; N \otimes \pi^*\mathcal{E})$  and  $\stackrel{v}{\operatorname{div}} \pi^{*\mathcal{E}} w \in \mathcal{R}^{m'-1}(SM; \pi^*\mathcal{E})$ . Furthermore,

$$\langle \nabla^{\mathbf{v}} \pi^* \mathcal{E} u, v \rangle_{L^2} = -\langle u, \operatorname{div}^{\mathbf{v}} \pi^* \mathcal{E} v \rangle_{L^2}.$$

In other words,  $\mathbb{X}$ ,  $\stackrel{\mathbf{v}}{\nabla} \pi^* \mathcal{E}$ , and  $\stackrel{\mathbf{v}}{\operatorname{div}} \pi^* \mathcal{E}$  map  $\mathcal{R}^m \to \mathcal{R}^{m-1}$  and their well-known adjoint relations are also satisfied on AH spaces as well. To prove the above lemma, we will use the following compactly supported version of it:

Lemma 5.9. The following are true.

1. If  $u, w \in C^{\infty}(SM; \pi^*\mathcal{E})$  are such that at least one of them is compactly supported in the interior M, then

$$\langle \mathbb{X}u, w \rangle_{L^2} = -\langle u, \mathbb{X}w \rangle_{L^2}.$$

2. If  $u, w \in C^{\infty}(SM; N \otimes \pi^* \mathcal{E})$  are such that at least one of them is compactly supported in the interior M, then

$$\langle \mathbb{X}u, w \rangle_{L^2} = -\langle u, \mathbb{X}w \rangle_{L^2}.$$

3. If  $u \in C^{\infty}(SM; \pi^* \mathcal{E})$  and  $w \in C^{\infty}(SM; N \otimes \pi^* \mathcal{E})$  are such that at least one of them is compactly supported in the interior M, then

$$\langle \nabla^{\mathbf{v}} \pi^* \mathcal{E} u, w \rangle_{L^2} = -\langle u, \operatorname{div}^{\mathbf{v}} \pi^* \mathcal{E} w \rangle_{L^2}.$$

**Proof:** The authors of [16] define  $\operatorname{div}^{\pi^*\mathcal{E}}$  as the  $L^2$  adjoint of  $\nabla^{\pi^*\mathcal{E}}$  in their Section 3.2 and then prove its existence in their Lemma 3.2, which proves our point 3).<sup>4</sup> To prove point 1), let  $(x^i)$  be coordinates of M, let  $(b_i)$  be a frame for  $\mathcal{E}$  over these coordinates' domain, and consider the coordinates  $v^{i\partial}/\partial x^i \mapsto (x^i, v^i)$  of TM. Let  ${}^{\mathcal{E}}\Gamma^k_{ij}$  denote the connection symbols of  $\nabla^{\mathcal{E}}$  with respect to  $(\partial/\partial x^i)$  and  $(b_k)$ . Now, suppose first that u or w is compactly supported over our coordinates' domain and write  $u = u^k \pi^* b_k$  and  $w = w^k \pi^* b_k$ . Then

$$\mathbb{X}u = \left[X(u^k) + {}^{\mathcal{E}}\Gamma^k_{ij}v^iu^j\right]\pi^*b_k$$

For convenience, assume that  $(b_i)$  is orthonormal so that we may write

$$\langle \mathbb{X}u, w \rangle_{L^2} = \sum_{k=1}^d \int_{SM} \left[ X(u^k) w^k + {}^{\mathcal{E}} \Gamma^k_{ij} v^i u^j w^k \right] dv_{SM}.$$

In Appendix A of [39] the authors prove that the  $L^2$  adjoint of  $X : C^{\infty}(SM) \to C^{\infty}(SM)$ is -X. Furthermore, since  $\nabla^{\mathcal{E}}$  is unitary it follows that the connection symbols are antisymmetric:  ${}^{\mathcal{E}}\Gamma_{ij}^k = -{}^{\mathcal{E}}\Gamma_{ik}^j$ . Applying these identities to the right-hand side above gives  $-\langle u, \mathbb{X}w \rangle_{L^2}$ . Point 1) then follows by a partition of unity argument. Point 3) is proved similarly where instead one uses the fact that the  $L^2$  adjoint of the operator  $X : C^{\infty}(SM; N) \to C^{\infty}(SM; N)$  defined in (4.18) is also -X, which is proved in Appendix A of [39].

#### Proof of Lemma 5.8 part 1):

Let u be as described in part 1). Let  $(\rho, y^{\mu}) = (x^i)$  be asymptotic boundary normal coordinates of  $\overline{M}$  as described in Section 2.2, let  $(b_i)$  be a frame for  $\mathcal{E}$  over these coordinates' domain, and consider the coordinates  $v^{i\partial}/\partial x^i \mapsto (x^i, v^i)$  of TM. Let  ${}^{\mathcal{E}}\Gamma^k_{ij}$  denote the connection symbols of  $\nabla^{\mathcal{E}}$  with respect to  $(\partial/\partial x^i)$  and  $(b_k)$  and let  $\Gamma^k_{ij}$  denote the Christoffel symbols with respect to  $(\partial/\partial x^i)$ . Consider also the coordinates  $\overline{\eta}_i dx^i / \rho \to (x^i, \overline{\eta}_i)$  of  ${}^0T^*\overline{M}$  and observe that the canonical identification  $H^* \circ \flat : TM \to {}^0T^*\overline{M}|_M$  is given by  $v^i = \rho^{-1}g^{ii'}\overline{\eta}_{i'}$ . Recall that  $g_{ij}$  and  $g^{ij}$  are respectively  $\rho^{-2}$  and  $\rho^2$  times something smooth on  $\overline{M}$ .

Writing  $u = u^k \pi^* b_k$ , we have that

$$\mathbb{X}u = \left[X(u^k) + {}^{\mathcal{E}}\Gamma^k_{ij}v^i u^j\right]\pi^* b_k.$$
(5.10)

Pulling  $v^i$  to  ${}^0S^*\overline{M}$  via the canonical identification gives  $\rho^{-1}g^{ii'}\overline{\eta}_{i'}$ , which is smooth over  ${}^0S^*\overline{M}$ . The terms  ${}^{\mathcal{E}}\Gamma^k_{ij}$  are smooth over  $\overline{M}$  by our standing assumption and hence on  ${}^0S^*\overline{M}$ 

<sup>&</sup>lt;sup>4</sup>Actually, the authors in the mentioned work are working over compact manifolds with boundary. But we can still apply their results by applying them to the compact manifold with boundary  $M_{\delta} = \{\rho \geq \delta\}$  for  $\delta > 0$  small enough so that  $M_{\delta}$  contains the (compact) intersection of the supports of u and w.

when lifted. Hence it quickly follows that the term  ${}^{\mathcal{E}}\Gamma^k_{ij}v^iu^j\pi^*b_k$  on the right-hand side of the above equation is in  $\mathcal{R}^m \subseteq \mathcal{R}^{m-1}$ . So, let's take a look at the other term:  $X(u^k)\pi^*b_k$ .

Consider the coordinates  $\xi_i dx^i \mapsto (x^i, \xi_i)$  of  $T^*\overline{M}$ . In (2.3) of [14] the authors write out an explicit equation for X over  $T^*M$  in asymptotic boundary normal coordinates (recall the convention about Greek and Latin indices):

$$X = \rho^2 \xi_0 \frac{\partial}{\partial \rho} + \rho^2 h^{\mu\nu} \xi_\mu \frac{\partial}{\partial y^\nu} - \left[ \rho \left( \xi_0^2 + |\xi|_h^2 \right) + \frac{1}{2} \rho^2 \partial_\rho |\xi|_h^2 \right] \frac{\partial}{\partial \xi_0} - \frac{1}{2} \rho^2 \partial_{y^k} |\xi|_h^2 \frac{\partial}{\partial \xi_k} + \frac{1}{2} \rho^2 \partial_\rho |\xi|_h^2 \frac{\partial}{\partial \xi_0} + \frac{1}{2} \rho^2 \partial_{y^k} |\xi|_h^2 \frac{\partial}{\partial \xi_k} + \frac{1}{2} \rho^2 \partial_\rho |\xi|_h^2 \frac{\partial}{\partial \xi_0} + \frac{1}{2} \rho^2 \partial_{y^k} |\xi|_h^2 \frac{\partial}{\partial \xi_k} + \frac{1}{2} \rho^2 \partial_\rho |\xi|_h^2 \frac{\partial}{\partial \xi_0} + \frac{1}{2} \rho^2 \partial_{y^k} + \frac{1}{$$

where  $|\xi|_h^2 = h^{\mu\nu}\xi_{\mu}\xi_{\nu}$ . Since canonical identification is given by  $\overline{\eta}_i = \rho\xi_i$ , it's a quick calculation to show that pushing X to  ${}^0S^*\overline{M}$  gives

$$X = \rho \overline{\eta}_0 \frac{\partial}{\partial \rho} + \rho h^{\mu\nu} \overline{\eta}_\mu \frac{\partial}{\partial y^\nu} - \left[ |\overline{\eta}|_h^2 + \frac{1}{2} \rho \partial_\rho |\overline{\eta}|_h^2 \right] \frac{\partial}{\partial \overline{\eta}_0} + \left[ \overline{\eta}_0 \overline{\eta}_\mu - \frac{1}{2} \rho \partial_{y^\mu} |\overline{\eta}|_h^2 \right] \frac{\partial}{\partial \overline{\eta}_\mu}.$$
 (5.11)

In particular, X extends to be a smooth vector on all of  ${}^{0}S^{*}\overline{M}$  and hence  $X(u^{k})\pi^{*}b_{k}$  is in  $\mathcal{R}^{m-1}$ . Thus by (5.10) we have that indeed  $\mathbb{X}u \in \mathcal{R}^{m-1}$ .

Now suppose that w is as in 1). We will prove the equality in 1) by multiplying w by a compactly supported (smooth) bump function, use Lemma 5.9 above, and then let the support of the bump function go out to infinity. To construct the suitable family of bump functions, let  $f_1 : [0, \infty) \to [0, \infty)$  be a smooth function that is identically zero on [0, 1/2], increasing on [1/2, 1], and then identically one on  $[1, \infty)$  (see Lemma 2.21 in [23] for an explicit construction). For any  $\delta > 0$ , let  $f_{\delta} : [0, \infty) \to [0, \infty)$  denote the function  $f_{\delta}(x) = f(x/\delta)$ . Finally, for  $\delta < \varepsilon$  let  $\phi_{\delta} : \overline{M} \to [0, \infty)$  denote the one parameter family of bump functions given by

$$\phi_{\delta}(x) = \begin{cases} f_{\delta} \circ \rho(x) & \rho(x) < \delta \\ 1 & \text{otherwise} \end{cases}$$

By Lemma 5.9 we have that

$$\langle \mathbb{X}u, \phi_{\delta}w \rangle_{L^2} = -\langle u, \mathbb{X}(\phi_{\delta}w) \rangle_{L^2}$$

since  $\phi_{\delta} w$  is compactly supported. Applying the product rule on the right-hand side gives

$$\langle \mathbb{X}u, \phi_{\delta}w \rangle_{L^2} = -\langle u, \phi_{\delta}\mathbb{X}w \rangle_{L^2} - \langle u, X(\phi_{\delta})w \rangle_{L^2}.$$
(5.12)

We now let  $\delta \to 0^+$  and show that this equation tends to the equality in 1). By what we proved above, we have that  $\langle \mathbb{X}u, w \rangle_{\pi^*\mathcal{E}}$  and  $\langle u, \mathbb{X}w \rangle_{\mathcal{E}}$  are the product of two  $L^2(SM)$  functions and hence also in  $L^2(SM)$ . Next, differentiating in  $\delta$  demonstrates that  $\phi_{\delta}$  monotonely increases to the identically one function as  $\delta \to 0^+$ . Hence by the dominated convergence theorem, we get that the first two terms in (5.12) tend to  $\langle \mathbb{X}u, w \rangle_{L^2}$  and  $-\langle u, \mathbb{X}w \rangle_{L^2}$  respectively as  $\delta \to 0^+$ .

Hence we will have proved 1) if we can show that the third term in (5.12) tends to zero as  $\delta \to 0^+$ . This will follow if we show that for any compact set  $K \subseteq \overline{M}$  contained in the domain of some interior coordinates  $(x^i)$  of  $\overline{M}$  or boundary coordinates  $(\rho, y^{\mu}) = (x^i)$  as above,

$$\int_{\pi^{-1}[K]} \langle u, X(\phi_{\delta})w \rangle_{\pi^*\mathcal{E}} \longrightarrow 0 \quad \text{as} \quad \delta \to 0^+$$

If K is contained in the domain of interior coordinates, then this follows immediately since  $\phi_{\delta} \equiv 1$  on K for sufficiently small  $\delta > 0$ . So suppose that K is contained in the domain of our boundary coordinates  $(\rho, y^{\mu}) = (x^i)$ . Writing the above integral in these coordinates as in Section 4.4 gives (here  $d\hat{x} = dx^0 \dots dx^n$ )

$$\left| \int_{K} \langle u, X(\phi_{\delta}) w \rangle_{\pi^{*} \mathcal{E}} dS_{x}(v) \sqrt{\det g} d\hat{x} \right| \leq \sup_{K} \left| \langle u, w \rangle_{\pi^{*} \mathcal{E}} \sqrt{\det g} \right| \omega_{n} \int_{K} |X(\phi_{\delta})| d\hat{x}.$$

where  $\omega_n$  denotes the surface area of the Euclidean *n*-sphere. The sup... is finite because  $|u|_{\pi^*\mathcal{E}}, |w|_{\pi^*\mathcal{E}} \in \rho^{(n+1)/2} L^{\infty}[K]$  and  $\sqrt{\det g} \in \rho^{-(n+1)} C^{\infty}(\overline{M})$ . Now, the explicit equation for X in coordinates of TM (e.g. see page 104 in [24]) gives that  $X(\phi_{\delta}) = v^0 f'_{\delta}(\rho)$ , which we note is supported in  $\{\rho \leq \delta\}$ . Since  $g = (d\rho^2 + h_{\mu\nu}dy^{\mu}dy^{\nu})/\rho^2$  and  $|v|_g = 1$ , we have that  $|v^0| \leq \rho$ . Letting  $K_y$  denote the (compact) ordinary projection of K onto the set  $\{(0, y^{\mu})\}$  in our coordinates, we can bound

$$\int_{K} |X(\phi_{\delta})| d\hat{x} \leq \int_{K_{y}} \int_{0}^{\delta} \delta f_{\delta}'(\rho) d\rho dy = \delta \int_{K_{y}} dy \longrightarrow 0 \quad \text{as} \quad \delta \to 0^{+}.$$

Hence the third term in (5.12) indeed tends to zero as  $\delta \to 0^+$ .

#### Proof of Lemma 5.8 parts 2), 3):

Let's begin with proving 2). Let u be as described there. We keep working in the same coordinates that we used in part 1) above. Writing  $u = \rho u^{ij} \pi^* \partial/\partial x^i \otimes \pi^* b_i$ , we have that

 $\mathbb{X}u$ 

$$= X \left(\rho u^{kj}\right) \pi^* \frac{\partial}{\partial x^k} \otimes \pi^* b_j \tag{5.13}$$

$$+\Gamma^{k}_{i'i}v^{i'}\rho u^{ij}\pi^*\frac{\partial}{\partial x^k}\otimes\pi^*b_j$$
(5.14)

$$+^{\mathcal{E}}\Gamma^{k}_{i'j}v^{i'}\rho u^{ij}\pi^{*}\frac{\partial}{\partial x^{i}}\otimes\pi^{*}b_{k}.$$
(5.15)

In part 1) we observed that  $v^{i'}$  is  $\rho$  times something smooth on  ${}^{0}S^*\overline{M}$  and hence it follows that the term (5.15) is in  $\mathcal{R}^m(SM; N \otimes \pi^*\mathcal{E}) \subseteq \mathcal{R}^{m-1}(SM; N \otimes \pi^*\mathcal{E})$ . Next, we have that since g is  $\rho^{-2}$  times a smooth metric on  $\overline{M}$ , the conformal transformation law of Christoffel symbols (e.g. see Proposition 7.29 in [24]) give that  $\Gamma_{ij}^k$  are  $\rho^{-1}$  times something smooth on  $\overline{M}$ . Thus the term (5.14) is also in  $\mathcal{R}^m \subseteq \mathcal{R}^{m-1}$ . Finally, from (5.11) we have that  $X(\rho)$  is  $\rho$  times something smooth on  ${}^{0}S^*\overline{M}$  and thus it follows from the product rule that the term (5.13) is in  $\mathcal{R}^{m-1}$ . Hence indeed  $\mathbb{X}u \in \mathcal{R}^{m-1}$ . The equality in 2) follows the same way that we proved the equality in 1).

Finally, let's prove 3). Let  $u = u^j \pi^* b_j$  and  $w = \rho w^{ij} \pi^* \partial / \partial x^i \otimes \pi^* b_j$  be as described there. In the proof of Lemma 3.2 in [16] the authors give an equation for the vertical derivative<sup>5</sup>

<sup>&</sup>lt;sup>5</sup>We remark that in their work they write " $\nabla^{\mathbf{v}}\mathcal{E}$ " for what we denote by " $\nabla^{\mathbf{v}\pi^*\mathcal{E}}$ ."

of u in terms of an operator " $\nabla$ " for which an explicit equation is given on page 350 of [39]. As the authors do in [16], we assume that  $(b_i)$  is orthonormal so that may use their formula to write that

$$\nabla^{\mathbf{v}} \pi^* \mathcal{E} u \text{ over } SM = \partial^i u^j \pi^* \frac{\partial}{\partial x^i} \otimes \pi^* b_j, \qquad (5.16)$$

where for any  $f \in C^{\infty}(SM)$ 

$$\partial_i f := \left[ \frac{\partial}{\partial v^i} (f \circ p) \right] \Big|_{SM},$$
$$\partial^i f := g^{ii'} \partial_{i'} f,$$

where  $p: TM \setminus \{0\} \to SM$  is the radial projection map  $v \mapsto v/|v|_g$  over the tangent bundle minus the zero section. Since the canonical identification  ${}^0S^*\overline{M} \cong SM$  is given by  $\overline{\eta}_i = \rho g_{ii'}v^{i'}$ , pushing  $\partial/\partial v^i$  to  ${}^0S^*\overline{M}$  gives

$$\frac{\partial}{\partial v^i}\longmapsto \rho g_{ii'}\frac{\partial}{\partial \overline{\eta}_{i'}}.$$

Thus  $\partial/\partial v^i$  extends to  $\rho^{-1}$  times a smooth vector field over  ${}^0S^*\overline{M}$  and hence so does  $\partial_i$ . Hence  $\partial^i$  is  $\rho$  times a smooth vector field over  ${}^0S^*\overline{M}$ . Thus by (5.16) we have that the vertical derivative of u is indeed in  $\mathcal{R}^{m-1}(SM; N \otimes \pi^* \mathcal{E})$ .

Next let's take a look at w. Using (5.16) above, a straightforward generalization of the derivation of the equation for "div Z" given on page 352 of [39] gives

$$\operatorname{div}^{^{\mathrm{v}}}\pi^{^{*}\mathcal{E}}w = \partial_{i}(\rho w^{ij})\pi^{*}b_{j}.$$
(5.17)

Since  $\partial_i$  only involves derivatives in  $v^i$ , we can pull  $\rho$  out of the derivative on the right-hand side. Since  $\partial_i$  is  $\rho^{-1}$  times a smooth vector field over  ${}^0S^*\overline{M}$ , it follows that this is indeed in  $\mathcal{R}^{m'-1}(SM;\pi^*\mathcal{E})$ .

The equality in 3) follow essentially the same way we proved the equality in 1). An example of a minor change that's needed is that the analog of (5.12) will be

$$\langle \nabla^{\mathbf{v}} \pi^* \mathcal{E} u, \phi_{\delta} w \rangle_{L^2} = -\langle u, \phi_{\delta} \operatorname{div}^{\mathbf{v}} \pi^* \mathcal{E} w \rangle_{L^2},$$

which we note doesn't have an analogous "third term" as in (5.12) because  $\phi_{\delta}$  only depends on position and thus isn't affected by the vertical divergence. From here one proceeds as before.

For use in Section 5.3 below, we record the  $\mathcal{R}^m$ -mapping property of the horizontal derivative as well.

**Lemma 5.18.** If 
$$u \in \mathcal{R}^m(SM; \pi^*\mathcal{E})$$
 with  $m \ge 1$ , then  $\stackrel{h}{\nabla}^{\pi^*\mathcal{E}} u \in \mathcal{R}^{m-1}(SM; N \otimes \pi^*\mathcal{E})$ .

**Proof:** Take any such u. Let  $(\rho, y^{\mu}) = (x^i)$ ,  $(b_i)$ ,  ${}^{\mathcal{E}}\Gamma^k_{ij}$ , and  $\Gamma^k_{ij}$  be as in the beginning of the proof of Lemma 5.8 part 1) and consider the coordinates  $v^{i\partial}/\partial x^i \mapsto (x^i, v^i)$  of TM and  $\overline{\eta}_i dx^i / \rho \to (x^i, \overline{\eta}_i)$  of  ${}^{0}T^*\overline{M}$  described there as well.

We write  $u = u^j \pi^* b_j$ . By the equations for the horizontal and vertical derivatives given in the proof of Lemma 3.2 in [16] and on page 350 of [39],

$$\nabla^{h} \nabla^{\pi^{*} \mathcal{E}} u$$

$$= \left( \delta^{i} u^{j} - \left( v^{k} \delta_{k} u^{j} \right) v^{i} \right) \pi^{*} \frac{\partial}{\partial x^{i}} \otimes \pi^{*} b_{j}$$
(5.19)

$$+u^{l} \nabla^{\pi^{*}\mathcal{E}} \left( {}^{\mathcal{E}} \Gamma^{j}_{kl} v^{k} \pi^{*} b_{j} \right)$$
(5.20)

where for any  $f \in C^{\infty}(SM)$ 

$$\delta_i f := \left[ \left( \frac{\partial}{\partial x^i} - \Gamma^k_{ij} v^j \frac{\partial}{\partial v^k} \right) (f \circ p) \right] \Big|_{SM}, \tag{5.21}$$

$$\delta^i f := g^{ii'} \delta_{i'} f, \tag{5.22}$$

where  $p: TM \setminus \{0\} \to SM$  is the radial projection map  $v \mapsto v/|v|_{q}$ .

Let's start by taking a look at the term (5.20). Recall our standing assumption that each  ${}^{\mathcal{E}}\Gamma^{j}_{kl}$  is smooth on  $\overline{M}$ . Next, we observed in the proof of Lemma 5.8 part 1) that each  $v^{k}$  is  $\rho$  times something smooth over  ${}^{0}S^{*}\overline{M}$ . By (5.16) the vertical derivative only involves derivatives in  $v^{i}$  and so we can pull the just mentioned factor of  $\rho$  out of the vertical derivative in (5.20). Thus the term (5.20) is equal to  $\rho u^{l}$  times  $\partial^{i}$  of something smooth on  ${}^{0}S^{*}\overline{M}$  times  $\pi^{*\partial}/\partial x^{i} \otimes \pi^{*}b_{j}$ . We observed in the proof of Lemma 5.8 parts 2) and 3) that  $\partial^{i}$  is a smooth vector field over  ${}^{0}S^{*}\overline{M}$  and thus  $\partial^{i}$  of something smooth on  ${}^{0}S^{*}\overline{M}$  is again smooth on  ${}^{0}S^{*}\overline{M}$ . From here it follows that the term (5.20) is in  $\mathcal{R}^{m} \subseteq \mathcal{R}^{m-1}$ . Finally let's take a look at the term (5.19).

Recall that  $g_{ij}$  and  $g^{ij}$  are respectively  $\rho^{-2}$  and  $\rho^2$  times something smooth on  $\overline{M}$ . We observed in the proof of Lemma 5.8 parts 2) and 3) that each  $\Gamma_{ij}^k v^j$  is smooth on  ${}^0S^*\overline{M}$ . Next, canonical identification is given by  $\overline{\eta}_i = \rho g_{ij} v^j$ , from which it's a quick computation to show that the differential of this canonical identification takes (recall that  $x^0 = \rho$ )

$$\frac{\partial}{\partial x^0} \longmapsto \frac{\partial}{\partial x^0} + \left(\rho^{-1}\overline{\eta}_i + \frac{\partial g_{ij}}{\partial \rho}g^{jj'}\overline{\eta}_{j'}\right)\frac{\partial}{\partial\overline{\eta}_i}$$
$$\frac{\partial}{\partial x^\lambda} \longmapsto \frac{\partial}{\partial x^\lambda} + \left(\frac{\partial g_{ij}}{\partial x^\lambda}g^{jj'}\overline{\eta}_{j'}\right)\frac{\partial}{\partial\overline{\eta}_i} \quad \text{for } \lambda = 1, \dots, n,$$
$$\frac{\partial}{\partial v^i} \longmapsto \rho g_{ii'}\frac{\partial}{\partial\overline{\eta}_{i'}}.$$

The important observation is that these are all  $\rho^{-1}$  times vector fields that are smooth over  ${}^{0}S^{*}\overline{M}$ . Thus by (5.21) and (5.22), the  $\delta_{i}$  and  $\delta^{i}$  are respectively  $\rho^{-1}$  and  $\rho$  times vector fields that are smooth over  ${}^{0}S^{*}\overline{M}$ . Plugging all of these observation into (5.19) finally gives that indeed the horizontal derivatives of u is in  $\mathcal{R}^{m-1}$ .

Next we need the following lemma that tells us that the curvature operators have the mapping property  $\mathcal{R}^m \to \mathcal{R}^m$ .

**Lemma 5.23.** Suppose that  $u \in \mathcal{R}^m(SM; N \otimes \pi^* \mathcal{E})$  and  $w \in \mathcal{R}^m(SM; \pi^* \mathcal{E})$  for integers  $m \ge 0$ . Then  $Ru \in \mathcal{R}^m(SM; N \otimes \pi^* \mathcal{E})$  and  $F^{\mathcal{E}}w \in \mathcal{R}^m(SM; N \otimes \pi^* \mathcal{E})$ .

**Proof:** Let  $(\rho, y^{\mu}) = (x^i)$ ,  $(b_i)$ ,  ${}^{\mathcal{E}}\Gamma^k_{ij}$ , and  $\Gamma^k_{ij}$  be as in the beginning of the proof of Lemma 5.8 part 1) and consider the coordinates  $v^{i\partial}/\partial x^i \mapsto (x^i, v^i)$  of TM and  $\overline{\eta}_i dx^i/\rho \to (x^i, \overline{\eta}_i)$  of  ${}^{0}T^*\overline{M}$  described there as well.

We write  $u = \rho u^{ij} \pi^* \partial / \partial x^i \otimes \pi^* b_j$  and  $w = w^j \pi^* b_j$ . We have that

$$Ru = \left(\frac{\partial\Gamma_{j'k}^l}{\partial x^{i}} - \frac{\partial\Gamma_{ik}^l}{\partial x^{j'}} + \Gamma_{j'k}^m\Gamma_{im}^l - \Gamma_{ik}^m\Gamma_{j'm}^l\right)\rho u^{ij}v^{j'}v^k\pi^*\frac{\partial}{\partial x^l}\otimes\pi^*b_j.$$

In the proof of Lemma 5.8 part 1) we observed that each  $v^i$  is  $\rho$  times something smooth on  ${}^{0}S^*\overline{M}$ . In the proof of Lemma 5.8 parts 2) and 3) we observed that each  $\Gamma_{ij}^k$  is  $\rho^{-1}$  times something smooth on  $\overline{M}$  and hence its first partials  $\partial \Gamma_{ij}^k / \partial x^i$  are  $\rho^{-2}$  times something smooth on  $\overline{M}$ . From this it follows that indeed  $Ru \in \mathcal{R}^m$ .

Next, looking at (4.20) and (4.21) we have that

$$F^{\mathcal{E}}w = P_{N\otimes\pi^{*}\mathcal{E}}\bigg(w^{l}g^{jj'}f_{ij}{}^{k}{}_{l}v^{i}\pi^{*}\frac{\partial}{\partial x^{j'}}\otimes\pi^{*}b_{k}\bigg).$$

By (4.19), we have that each  $f_{ij}{}^k{}_l$  is smooth on  $\overline{M}$ , and recall that each  $g^{ij}$  is  $\rho^2$  times something smooth on  $\overline{M}$ . The projection map here is given by

$$P_{N\otimes\pi^{*}\mathcal{E}}\left(\pi^{*}\frac{\partial}{\partial x^{j'}}\otimes\pi^{*}b_{k}\right) = \pi^{*}\left(\frac{\partial}{\partial x^{j'}}-\langle v,\frac{\partial}{\partial x^{j'}}\rangle v\right)\otimes\pi^{*}b_{k}$$
$$=\pi^{*}\left(\frac{\partial}{\partial x^{j'}}-g_{i'j'}v^{i'}v^{i''}\frac{\partial}{\partial x^{i''}}\right)\otimes\pi^{*}b_{k}$$

From here it follows that indeed  $F^{\mathcal{E}}w \in \mathcal{R}^m$ .

We need one final lemma that provides the needed commutator formulas to prove Theorem 5.7. The following lemma is Lemma 3.2 in [16], where one can also find a proof.

**Lemma 5.24.** The following are true, where  $[\ldots, \ldots]$  denotes the commutator bracket.

$$\left[\mathbb{X}, \overset{\mathbf{v}}{\nabla}^{\pi^* \mathcal{E}}\right] = -\overset{\mathbf{h}}{\nabla}^{\pi^* \mathcal{E}},\tag{5.25}$$

$$\left[\mathbb{X}, \stackrel{\mathbf{h}}{\nabla}^{\pi^* \mathcal{E}}\right] = R \stackrel{\mathbf{v}}{\nabla}^{\pi^* \mathcal{E}} + F^{\mathcal{E}}, \qquad (5.26)$$

$$\operatorname{div}^{\mathrm{h}} \pi^{*\mathcal{E}} \nabla^{\mathrm{v}} \pi^{*\mathcal{E}} - \operatorname{div}^{\mathrm{v}} \pi^{*\mathcal{E}} \nabla^{\mathrm{h}} \pi^{*\mathcal{E}} = n \mathbb{X}, \qquad (5.27)$$

$$\left[\mathbb{X}, \operatorname{div}^{\mathrm{v}}^{\pi^* \mathcal{E}}\right] = -\operatorname{div}^{\mathrm{h}}^{\pi^* \mathcal{E}}.$$
(5.28)

#### Proof of Theorem 5.7:

Let u be as described in the theorem. By Lemma 5.8 we have that

$$\langle \nabla^{\mathbf{v}} \pi^* \mathcal{E} \mathbb{X} u, \nabla^{\mathbf{v}} \pi^* \mathcal{E} \mathbb{X} u \rangle_{L^2} = \langle \mathbb{X} \mathrm{div}^{\mathbf{v}} \pi^* \mathcal{E} \nabla^{\mathbf{v}} \pi^* \mathcal{E} \mathbb{X} u, u \rangle_{L^2}.$$

We get that this is equal to (see right after for justifications)

$$\langle -\operatorname{div}^{h} \pi^{*\mathcal{E}} \nabla^{v} \pi^{*\mathcal{E}} \mathbb{X} u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \mathbb{X} \nabla^{\pi^{*\mathcal{E}}} \mathbb{X} u \rangle_{L^{2}},$$

$$= \langle -\operatorname{div}^{h} \pi^{*\mathcal{E}} \nabla^{v} \pi^{*\mathcal{E}} \mathbb{X} u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \mathbb{X} \nabla^{h} \pi^{*\mathcal{E}} u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \mathbb{X} \mathbb{X} \nabla^{\pi^{*\mathcal{E}}} u, u \rangle_{L^{2}},$$

$$= \langle -\operatorname{div}^{h} \pi^{*\mathcal{E}} \nabla^{v} \pi^{*\mathcal{E}} \mathbb{X} u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \left( R \nabla^{v} \pi^{*\mathcal{E}} + F^{\mathcal{E}} \right) u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \nabla^{h} \mathbb{X} u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \mathbb{X} \mathbb{X} \nabla^{\pi^{*\mathcal{E}}} u, u \rangle_{L^{2}},$$

$$= \langle -n \mathbb{X} \mathbb{X} u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \left( R \nabla^{v} \pi^{*\mathcal{E}} + F^{\mathcal{E}} \right) u + \operatorname{div}^{v} \pi^{*\mathcal{E}} \mathbb{X} \mathbb{X} \nabla^{v} \pi^{*\mathcal{E}} u, u \rangle_{L^{2}},$$

where in the above four lines we used respectively (5.28), (5.25), (5.26), and (5.27). Applying Lemma 5.8 again gives that this is equal to

$$n\langle \mathbb{X}u, \mathbb{X}u \rangle_{L^2} + \langle \operatorname{div}^{\mathsf{v}} \pi^* \mathcal{E} R \nabla^{\mathsf{v}} \pi^* \mathcal{E} u + \operatorname{div}^{\mathsf{v}} \pi^* \mathcal{E} F^{\mathcal{E}} u, u \rangle_{L^2} + \langle \mathbb{X} \nabla^{\mathsf{v}} \pi^* \mathcal{E} u, \mathbb{X} \nabla^{\mathsf{v}} \pi^* \mathcal{E} u \rangle_{L^2}$$

Splitting the second inner product over the "+" sign and then applying Lemma 5.8 to the resultant middle two terms proves the theorem.

## 5.3 Finite Degree of Solutions to Transport Equations

In the proof of Theorem 2.7 we will end up showing that Q – id satisfies an equation of a form similar to

$$Xu + \Phi u = f$$

over SM, which is called a "transport equation." It turns out that this equation has good behavior with respect to vertical Fourier analysis, which we now introduce. Consider the **vertical Laplacian**:

$$\Delta^{\pi^*\mathcal{E}} = -\operatorname{div}^{\operatorname{v}} \pi^{*\mathcal{E}} \nabla^{\pi^*\mathcal{E}} : C^{\infty}(SM; \pi^*\mathcal{E}) \longrightarrow C^{\infty}(SM; \pi^*\mathcal{E}).$$

By Lemma 5.8 3) this operator has the  $\mathcal{R}^l$ -mapping property  $\mathcal{R}^l(SM; \pi^*\mathcal{E}) \to \mathcal{R}^{l-2}(SM; \pi^*\mathcal{E})$ for  $l \geq 2$ . Let's see what this operator looks like in coordinates. Let  $(x^i)$  be coordinates of M, let  $(r_i)$  be an *orthonormal* frame of TM over their domain, let  $(b_j)$  denote a frame of  $\mathcal{E}$ over their domain, and consider the coordinates

$$v^i r_i \longmapsto \left(x^i, v^i\right) \tag{5.29}$$

of TM. Then we claim that for any smooth section  $u = u^j \pi^* b_j$ ,

$$\Delta^{\pi^* \mathcal{E}} u = \left( -\Delta^{\mathbb{S}^n} u^j \right) \pi^* b_j \tag{5.30}$$

where " $-\Delta^{\mathbb{S}^n v}$  is the negative Laplacian on the *n*-sphere in the variables  $v^i$ . This is most easily seen as follows. Pick an arbitrary point  $x_0 \in M$  in the domain of our coordinates, choose normal coordinates  $(\hat{x}^i)$  of M centered at  $x_0$ , and consider the coordinates  $\hat{v}^{i\partial}/\partial \hat{x}^i \mapsto$  $(\hat{x}^i, \hat{v}^i)$  of TM. Then observe that (5.16) and (5.17) tell us that on the sphere  $S_{x_0}M$ , the operator  $\Delta^{\pi^*\mathcal{E}}$  applied to  $u = \hat{u}^j \pi^* b_j$  is given by  $(-\Delta^{\mathbb{S}^n} \hat{u}^j) \pi^* b_j$ . The claim then follows by pushing this expression through the change of variables  $(\hat{x}^i, \hat{v}^i) \mapsto (x^i, v^i)$ .

From this observation and the theory of spherical harmonics (c.f. Section 2.H in [12] for the latter), we obtain several important implications regarding the vertical Laplacian. First, we get that the eigenvalues of  $\Delta^{\pi^*\mathcal{E}}$  match that of  $-\Delta^{\mathbb{S}^n}$ :

$$\lambda_m = m(m+n-1)$$
 for integers  $m \ge 0$ .

Furthermore, letting  $\Omega_m$  denote the set of smooth eigenfunctions of  $\Delta^{\pi^*\mathcal{E}}$  with eigenvalue  $\lambda_m$ , any  $u \in C^{\infty}(SM; \pi^*\mathcal{E})$  can be uniquely decomposed as the (pointwise converging) "Fourier series"

$$u = \sum_{m=0}^{\infty} u_m, \qquad u \in \Omega_m$$

Furthermore, for any fixed  $x \in M$  this convergence also holds in  $L^2(S_xM; (\pi^*\mathcal{E})_x)$ . The  $u_m$ 's are called u's **Fourier modes**. The maximum index m for which  $u_m \neq 0$  is called the **degree** of u and is denoted by "deg u" (which could be infinity). Naturally, we say that u is of **finite degree** if its degree is finite. We can write an explicit equation for the Fourier modes as follows. For each  $m \in \mathbb{Z}_+$  we let

$$\{Y_k^m: k=1,\ldots,l_m\}$$

denote a real-valued orthonormal basis of eigenfunctions of  $-\Delta^{\mathbb{S}^n}$  with eigenvalue  $\lambda_m$ . Then in the coordinates (5.29) and frame  $(b_i)$  there

$$u_{m}^{j}(x^{i}, v^{i}) = \sum_{k=1}^{l_{m}} \left[ \int_{\mathbb{S}^{n}} u^{j}(x^{i}, w^{i}) Y_{k}^{m}(w^{i}) dw_{\mathbb{S}^{n}} \right] Y_{k}^{m}(v^{i})$$
(5.31)

(no implicit summation meant in m here). An important property of the vertical Laplacian eigenspaces is that they are orthogonal:

**Proposition 5.32.** The spaces  $\Omega_m \cap L^2(SM; \pi^*\mathcal{E})$  and  $\Omega_{m'} \cap L^2(SM; \pi^*\mathcal{E})$  are orthogonal with respect to  $L^2$  when  $m \neq m'$ .

The above proposition follows directly from the theory of spherical harmonics and the fact that integrals over SM can be partitioned as described in Section 4.4 above. Another important property is that the vertical Laplacian commutes with taking the  $m^{\text{th}}$  Fourier mode:

**Proposition 5.33.** If  $u \in C^{\infty}(SM; \pi^*\mathcal{E})$ , then  $\Delta^{\pi^*\mathcal{E}}(u_m) = (\Delta^{\pi^*\mathcal{E}}u)_m$ .

**Proof:** Using the coordinate expression for the vertical Laplacian (5.30), this follows by taking (5.31) and integrating by parts:

$$\left[\Delta^{\pi^*\mathcal{E}}(u_m)\right]^j = \lambda_m u_m^j = \sum_{k=1}^{l_m} \left[\int_{\mathbb{S}^n} u^j \lambda_m Y_k^m dw_{\mathbb{S}^n}\right] Y_k^m = \sum_{k=1}^{l_m} \left[\int_{\mathbb{S}^n} u^j \left(-\Delta^{\mathbb{S}^n}\right) Y_k^m dw_{\mathbb{S}^n}\right] Y_k^m$$

$$=\sum_{k=1}^{l_m}\left[\int_{\mathbb{S}^n}\left(\left(-\Delta^{\mathbb{S}^n}\right)u^j\right)Y_k^mdw_{\mathbb{S}^n}\right]Y_k^m=\left(\Delta^{\pi^*\mathcal{E}}u\right)_m^j.$$

Next we'll need the fact that Fourier modes have the same regularity as their original section:

**Proposition 5.34.** If  $u \in \mathcal{R}^{l}(SM; \pi^{*}\mathcal{E})$  for  $l \geq 0$ , then each Fourier mode  $u_{m} \in \mathcal{R}^{l}(SM; \pi^{*}\mathcal{E})$  as well.

**Proof:** Take any  $u \in \mathcal{R}^{l}(SM; \pi^{*}\mathcal{E})$  with  $l \geq 0$  and fix  $m \geq 0$ . Let  $(x^{i})$  be coordinates of  $\overline{M}$ , let  $(r_{i})$  be an *orthonormal* frame of TM over dom  $(x^{i})$ , and consider the coordinates  $v^{i}r_{i} \mapsto (x^{i}, v^{i})$  of TM. Let  $(b_{j})$  denote a frame of  $\mathcal{E}$  over dom  $(x^{i})$ . We write  $u = u^{j}\pi^{*}b_{j}$ . Furthermore, let  $K \subseteq \text{dom}(x^{i}) \subseteq \overline{M}$  be any compact subset.

First suppose that l = 0. Since  $u^j \in \rho^{(n+1)/2} L^{\infty}[K]$ , (5.31) tells us that each  $u^j_m \in \rho^{(n+1)/2} L^{\infty}[K]$  as well and so indeed  $u_m \in \mathcal{R}^l$ .

Next suppose that l = 1. Suppose that the frame  $(r_i)$  was obtained by mapping a local orthonormal frame  $(\overline{\zeta}^i)$  of  ${}^0T^*\overline{M}$  via the canonical identification  $\sharp \circ H : {}^0T^*\overline{M}|_M \to TM$ , and consider the coordinates  $\overline{\eta}_i \overline{\zeta}^i \mapsto (x^i, \overline{\eta}_i)$  of  ${}^0T^*\overline{M}$ . Observe that this identification is given by  $\overline{\eta}_i = v^i$ . Now, take any smooth vector field  $V_1 \in C^{\infty}({}^0S^*\overline{M}; T^0S^*\overline{M})$  which we write as  $V = (V_1)^r \partial/\partial x^r + (V_1)_r \partial/\partial \overline{\eta}_r$ . Pushing V to SM gives  ${}^6V = (V_1)^r \partial/\partial x^r + \sum_{r=0}^n (V_1)_r \partial/\partial v^r$ . Thus by (5.31)

$$V_1(u_m^j) = (V_1)^r \sum_{k=1}^{l_m} \left[ \int_{\mathbb{S}^n} \frac{\partial u^j}{\partial x^r} Y_k^m dw_{\mathbb{S}^n} \right] Y_k^m + \sum_{k=1}^{l_m} \left[ \int_{\mathbb{S}^n} u^j Y_k^m dw_{\mathbb{S}^n} \right] \sum_{r=0}^n (V_1)_r \frac{\partial Y_k^m}{\partial v^r}.$$

Now, the components  $(V_1)^i$ ,  $(V_1)_i$  are bounded above K because they are smooth over the compact  ${}^0S^*\overline{M} \cap \pi_0^{-1}[K]$ . For future use, we point out that this also holds for their  $\partial/\partial x^i$ ,  $\partial/\partial v^i$  first and higher order partials as well. Furthermore, since  $\partial/\partial x^r$  is a smooth vector field over the 0-cosphere bundle, we have that each  $\partial u^j/\partial x^r$  in the above expression is in  $\rho^{(n+1)/2}L^{\infty}$  by assumption. Hence the above expression tells us that  $V_1(u_m^j) \in \rho^{(n+1)/2}L^{\infty}[K]$  and so indeed  $u_m \in \mathcal{R}^l$ . The cases  $l \geq 2$  are handled similarly.

One of the central properties of X is that it maps

$$\mathbb{X}: \Omega_m \longrightarrow \Omega_{m-1} \oplus \Omega_{m+1}. \tag{5.35}$$

This is proven in Section 3.4 of [16]. Similarly, multiplication on the left by  $\Phi$  maps  $\Omega_m \to \Omega_m$ since  $\Phi$  has no dependence on the vertical variable "v." In particular, we see that the operator in the transport equation " $\mathbb{X} + \Phi$ " maps sections of finite degree to sections of finite degree. The converse is also true, which is the main result of this section:

<sup>&</sup>lt;sup>6</sup>We can't use the Einstein summation convention on  $(V_1)_r^{\partial}/\partial v^r$  because it has two lower indices.

**Theorem 5.36.** Assume that the sectional curvatures of g are negative.<sup>7</sup> If  $u \in \mathcal{R}^3(SM; \pi^*\mathcal{E})$  solves

$$Xu + \Phi u = f \tag{5.37}$$

for some  $f \in C^{\infty}(SM; \pi^* \mathcal{E})$  of finite degree, then u is also of finite degree.

To prove this, we need several preliminary results.

Lemma 5.38. It holds that

$$\left[\mathbb{X}, \Delta^{\pi^* \mathcal{E}}\right] = 2 \operatorname{div}^{\mathsf{v}} \pi^* \mathcal{E} \stackrel{\mathsf{h}}{\nabla} \pi^* \mathcal{E} + n \mathbb{X}.$$

The above lemma is stated as Lemma 3.4 of [16], whose proof is essentially identical to that of Lemma 3.5 in [39].

To state the next preliminary result we observe that because of (5.35), over each  $\Omega_m$  we can decompose  $\mathbb{X} = \mathbb{X}_- + \mathbb{X}_+$  where

$$\mathbb{X}_{\pm}:\Omega_m\longrightarrow\Omega_{m\pm 1}.$$

We point out that the maps  $\mathbb{X}_{\pm}$  are distinct for different  $\Omega_m$  even though we use the same notation to denote them.

**Remark 5.39.** Like X, the operators  $\mathbb{X}_{\pm}$  map  $\mathcal{R}^l \to \mathcal{R}^{l-1}$  for  $l \geq 1$  by Lemma 5.8 and Proposition 5.34 above.

We mention that the idea of splitting the action of the geodesic vector field as above was first introduced by Guillemin and Kazhdan - see [17]. The following preliminary result is a special case of the Pestov identity with a connection (Theorem 5.7):

**Proposition 5.40.** Suppose that  $u \in \Omega_m \cap \mathcal{R}^2(SM; \pi^*\mathcal{E})$ . Then

$$= \left\| \nabla^{\pi^* \mathcal{E}} u \right\|_{L^2}^2 + (2m+n-2) \|\mathbb{X}_{-}u\|_{L^2}^2 - \langle R \nabla^{\pi^* \mathcal{E}} u, \nabla^{\pi^* \mathcal{E}} u \rangle_{L^2} - \langle F^{\mathcal{E}} u, \nabla^{\pi^* \mathcal{E}} u \rangle_{L^2}.$$

 $(2m+n)\|X_{+}u\|_{\infty}^{2}$ 

**Proof:** We have that u satisfies the equation in Theorem 5.7. Let's take a look at the term

$$\begin{aligned} \left\| \mathbb{X} \overset{\mathbf{v}}{\nabla}^{\pi^* \mathcal{E}} u \right\|_{L^2}^2 &= \langle -\overset{\mathbf{h}}{\nabla}^{\pi^* \mathcal{E}} u + \overset{\mathbf{v}}{\nabla}^{\pi^* \mathcal{E}} \mathbb{X} u, -\overset{\mathbf{h}}{\nabla}^{\pi^* \mathcal{E}} u + \overset{\mathbf{v}}{\nabla}^{\pi^* \mathcal{E}} \mathbb{X} u \rangle_{L^2} \qquad \text{by (5.25),} \\ &= \left\| \overset{\mathbf{h}}{\nabla}^{\pi^* \mathcal{E}} u \right\|_{L^2}^2 + 2 \langle \mathbb{X} u, \overset{\mathbf{v}}{\operatorname{div}}^{\pi^* \mathcal{E}} \overset{\mathbf{h}}{\nabla}^{\pi^* \mathcal{E}} u \rangle_{L^2} + \left\| \overset{\mathbf{v}}{\nabla}^{\pi^* \mathcal{E}} \mathbb{X} u \right\|_{L^2}^2 \qquad \text{Lemma 5.8 3) and Lemma 5.18.} \end{aligned}$$

Applying Lemma 5.38, we see that the middle term in the last quantity is equal to

$$\langle \mathbb{X}u, \mathbb{X}\Delta^{\pi^*\mathcal{E}}u - \Delta^{\pi^*\mathcal{E}}\mathbb{X}u - n\mathbb{X}u \rangle_{L^2}.$$

Splitting  $Xu = X_{-}u + X_{+}u \in \Omega_{m-1} \oplus \Omega_{m+1}$ , using that

$$\Delta^{\pi^*\mathcal{E}} u = \lambda_m u, \qquad \Delta^{\pi^*\mathcal{E}} \mathbb{X}_- u = \lambda_{m-1} \mathbb{X}_- u, \qquad \Delta^{\pi^*\mathcal{E}} \mathbb{X}_+ u = \lambda_{m+1} \mathbb{X}_+ u,$$

using the orthogonality of the vertical Laplacian eigenspaces, and then plugging the result into the equation in Theorem 5.7 proves the proposition after several cancellations.

<sup>&</sup>lt;sup>7</sup>Recall our standing assumption that  $\Phi$  is skew-Hermitian.

The following lemma provides the contraction property that's needed in the proof of Theorem 5.36.

**Lemma 5.41.** Suppose that the sectional curvatures of g are negative. Then there exist real constants  $c_m \to \infty$  such that for sufficiently large m,

$$\begin{cases} \|\mathbb{X}_{-}u\|_{L^{2}}^{2} + c_{m}\|u\|_{L^{2}}^{2} \leq \|\mathbb{X}_{+}u\|_{L^{2}}^{2} & \text{if } n \neq 2, \\ \|\mathbb{X}_{-}u\|_{L^{2}}^{2} + c_{m}\|u\|_{L^{2}}^{2} \leq d_{m}\|\mathbb{X}_{+}u\|_{L^{2}}^{2} & \text{if } n = 2, \end{cases}$$

$$(5.42)$$

for all  $u \in \Omega_m \cap \mathcal{R}^2(SM; \pi^*\mathcal{E})$  where  $d_m = 1 + 1/[(2m-1)(m+1)^2]$ .

**Proof:** We begin by using the fact that the sectional curvatures of g tend to -1 at  $\partial M$ . Precisely, by the remark after Proposition 1.10 in [29] there exists an  $\varepsilon > 0$  so that the sectional curvatures of g are less than  $-\kappa'$  for some  $\kappa' > 0$  over the region  $\{\rho < \varepsilon\}$ . Hence this, the compactness of  $\{\rho \ge \varepsilon\}$ , and the negative curvature assumption imply that there exists a  $\kappa > 0$  such that the sectional curvatures of g are bounded above by  $-\kappa$  on all of M.

Now, take any u as in the statement of the lemma. We have that u satisfies the equation in Proposition 5.40 above. We begin by estimating the  $L^2$  norm of the term  $\nabla^{\pi^* \mathcal{E}} u$  by utilizing the trick of looking at its vertical divergence. By Lemma 5.38 we have that

$$\begin{aligned} \operatorname{div}^{\mathsf{v}} \pi^{*\mathcal{E}} \overset{\mathsf{h}}{\nabla} \pi^{*\mathcal{E}} u &= \frac{1}{2} \mathbb{X} \Delta^{\pi^{*\mathcal{E}}} u - \frac{1}{2} \Delta^{\pi^{*\mathcal{E}}} \mathbb{X} u - \frac{n}{2} \mathbb{X} u, \\ &= \frac{1}{2} (\mathbb{X}_{+} \lambda_{m} u + \mathbb{X}_{-} \lambda_{m} u) - \frac{1}{2} (\lambda_{m+1} \mathbb{X}_{+} u + \lambda_{m-1} \mathbb{X}_{-} u) - \frac{n}{2} (\mathbb{X}_{+} u + \mathbb{X}_{-} u), \\ &= -(m+n) \mathbb{X}_{+} u + (m-1) \mathbb{X}_{-} u, \\ &= -\operatorname{div}^{\mathsf{v}} \pi^{*\mathcal{E}} \left( -\frac{m+n}{\lambda_{m+1}} \overset{\mathsf{v}}{\nabla} \pi^{*\mathcal{E}} \mathbb{X}_{+} u + \frac{m-1}{\lambda_{m-1}} \overset{\mathsf{v}}{\nabla} \pi^{*\mathcal{E}} \mathbb{X}_{-} u \right). \end{aligned}$$

Plugging the expression for the  $\lambda_k$ 's into this, we conclude that

$$\overset{\mathbf{h}}{\nabla}^{\pi^{*}\mathcal{E}}u = \frac{1}{m+1}\overset{\mathbf{v}}{\nabla}^{\pi^{*}\mathcal{E}}\mathbb{X}_{+}u - \frac{1}{m+n-2}\overset{\mathbf{v}}{\nabla}^{\pi^{*}\mathcal{E}}\mathbb{X}_{-}u + Z$$

where  $Z \in C^{\infty}(SM; N \otimes \pi^* \mathcal{E})$  is such that  $\operatorname{div}^{v} \pi^* \mathcal{E} Z = 0$  and hence perpendicular to the other two terms on the right-hand side with respect to  $L^2$ . Using the orthogonality of the vertical Laplacian eigenspaces and Lemma 5.8 3), this gives us the  $L^2$  estimate

$$\left\| \nabla^{\mathbf{h}^{*}\mathcal{E}} u \right\|_{L^{2}}^{2} \geq \frac{m+n}{m+1} \| \mathbb{X}_{+} u \|_{L^{2}}^{2} + \frac{m-1}{m+n-2} \| \mathbb{X}_{-} u \|_{L^{2}}^{2}.$$

Now, plugging this into the equation in Proposition 5.40 gives

$$\left(2m+n-\frac{m+n}{m+1}\right)\|\mathbb{X}_{+}u\|_{L^{2}}^{2}$$
(5.43)

$$\geq \left(2m+n-2+\frac{m-1}{m+n-2}\right) \|\mathbb{X}_{-}u\|_{L^{2}}^{2} - \langle R \nabla^{\mathsf{v}}\pi^{*}\mathcal{E}u, \nabla^{\mathsf{v}}\pi^{*}\mathcal{E}u \rangle_{L^{2}} - \langle F^{\mathcal{E}}u, \nabla^{\mathsf{v}}\pi^{*}\mathcal{E}u \rangle_{L^{2}}.$$

We have that the term

$$-\langle R \nabla^{\mathsf{v}} \pi^* \mathcal{E} u, \nabla^{\mathsf{v}} \pi^* \mathcal{E} u \rangle_{L^2} \ge \kappa \left\| \nabla^{\mathsf{v}} \pi^* \mathcal{E} u \right\|_{L^2}^2 = \kappa \lambda_m \|u\|_{L^2}^2.$$
(5.44)

Furthermore,

$$-\langle F^{\mathcal{E}}u, \nabla^{\mathbf{v}\pi^{*}\mathcal{E}}u \rangle_{L^{2}} \geq -\left\| F^{\mathcal{E}} \right\|_{L^{\infty}} \|u\|_{L^{2}} \left\| \nabla^{\mathbf{v}\pi^{*}\mathcal{E}}u \right\|_{L^{2}} = -\left\| F^{\mathcal{E}} \right\|_{L^{\infty}} \lambda_{m}^{1/2} \|u\|_{L^{2}}^{2}.$$
(5.45)

Hence, letting  $a_{m,n}$  and  $b_{m,n}$  denote the coefficients of  $\|X_+u\|_{L^2}^2$  and  $\|X_-u\|_{L^2}^2$  in (5.43) above respectively we get that

$$\frac{a_{m,n}}{b_{m,n}} \|\mathbb{X}_{+}u\|_{L^{2}}^{2} \geq \|\mathbb{X}_{-}u\|_{L^{2}}^{2} + \frac{\kappa\lambda_{m} - \left\|F^{\mathcal{E}}\right\|_{L^{\infty}}\lambda_{m}^{\frac{1}{2}}}{b_{m,n}} \|u\|_{L^{2}}^{2}.$$

Elementary algebra shows that  $a_{m,n}/b_{m,n}$  is less than or equal to 1 if  $n \neq 2$  and m > 1, and is equal to  $d_m$  if n = 2. Since  $\lambda_m = O(m^2)$ , the lemma follows.

We need one last technical lemma:

**Lemma 5.46.** If  $u \in \mathcal{R}^3(SM; \pi^*\mathcal{E})$ , then  $\|\mathbb{X}_+u_m\|_{L^2} \to 0$  as  $m \to \infty$ .

**Proof:** Take any  $u \in \mathcal{R}^3(SM; \pi^*\mathcal{E})$ . Let  $(x^i)$  be coordinates of  $\overline{M}$ , let  $(b_j)$  denote a frame of  $\mathcal{E}$  over dom  $(x^i)$ , and consider the coordinates  $v^{i\partial}/\partial x^i \mapsto (x^i, v^i)$  of TM. Take any compact subset  $K \subseteq \text{dom}(x^i) \subseteq \overline{M}$  and suppose for convenience that  $(b_j)$  is orthonormal. We will show that  $\|X_+u_m\|_{L^2(\pi^{-1}[K])} \to 0$  as  $m \to \infty$ , from which the lemma will follow by covering the compact  $\overline{M}$  by a finite number of such sets K.

We have that (we omit writing  $\pi^{-1}[K]$  for the rest of the proof)

$$\|\mathbb{X}_{+}u_{m}\|_{L^{2}} = \left\| \left[ X\left(u_{m}^{k}\right) + {}^{\mathcal{E}}\Gamma_{ij}^{k}v^{i}u_{m}^{j} \right]\pi^{*}b_{k} \right\|_{L^{2}}$$

$$\leq \sum_{k=1}^{d} \left\| X\left(u_{m}^{k}\right) \right\|_{L^{2}} + \sup_{v \in \pi^{-1}[K]} \left| {}^{\mathcal{E}}\Gamma_{ij}^{k}v^{i} \right| \left\| u_{m}^{j} \right\|_{L^{2}}.$$
(5.47)

Since  $g_{ij}v^iv^j \equiv 1$  and  $g_{ij}$  blows up like  $\rho^{-2}$  at the boundary, we have that each  $v^i$  is bounded over the compact K and hence the sup... above is finite. From Proposition 5.33 it follows that the last  $L^2$ -norm satisfies

$$\left\|u_{m}^{j}\right\|_{L^{2}} \leq \left\|u_{m}\right\|_{L^{2}} = \frac{1}{\lambda_{m}} \left\|\left(\Delta^{\pi^{*}\mathcal{E}}u\right)_{m}\right\|_{L^{2}} \leq \frac{1}{\lambda_{m}} \left\|\Delta^{\pi^{*}\mathcal{E}}u\right\|_{L^{2}} \to 0 \quad \text{as } m \to \infty,$$

where we've used that  $\Delta^{\pi^*\mathcal{E}} u \in \mathcal{R}^{3-2}(SM; \pi^*\mathcal{E}) \subseteq L^2(SM; \pi^*\mathcal{E})$  and so the last  $L^2$ -norm is finite. Thus by (5.47), the lemma will be proved if we show that  $\|X(u_m^k)\|_{L^2} \to 0$  as  $m \to \infty$ . Let  $(r_i)$  be an orthonormal frame of TM over dom  $(x^i)$  and now consider the coordinates  $v^i r_i \mapsto (x^i, v^i)$  of TM. Suppose furthermore that the frame  $(r_i)$  was obtained by mapping a local orthonormal frame  $(\overline{\zeta}^i)$  of  ${}^0T^*\overline{M}$  via the canonical identification  $\sharp \circ H : {}^0T^*\overline{M}|_M \to TM$ , and consider the coordinates  $\overline{\eta}_i \overline{\zeta}^i \mapsto (x^i, \overline{\eta}_i)$  of  ${}^0T^*\overline{M}$ . Recall that this identification is given by  $\overline{\eta}_i = v^i$ . In (5.11) we pointed out that X extends to a smooth vector field  $X_0$  over  ${}^0S^*\overline{M}$ . Let us write  $X_0 = (X_0)^i \partial/\partial x^i + (X_0)_i \partial/\partial \overline{\eta}_i$  over  ${}^0S^*\overline{M}$ , which if we push to SM we get  $X = (X_0)^i \partial/\partial x^i + \sum_{r=0}^n (X_0)_i \partial/\partial v^i$ . Thus

$$Xu_m^k = (X_0)^i \frac{\partial u_m^k}{\partial x^i} + \sum_{i=0}^n (X_0)_i \frac{\partial u_m^k}{\partial v^i}.$$

where  $\partial u_m^k / \partial v^i$  denotes  $\partial / \partial v^i$  applied to any smooth extension of  $u_m^k$  from SM. For definitiveness, let us say that we extend  $u_m^k$  to TM minus the zero section by making  $u_m^k$  constant along the radial lines  $t \mapsto tv$  for  $t \in (0, \infty)$  and  $v \in SM$ . Now, the components  $(X_0)^i$ ,  $(X_0)_i$  are bounded above K because they are smooth over the compact  ${}^0S^*\overline{M} \cap \pi_0^{-1}[K]$ . Hence by the above equation, the lemma will be proved if we show that both  $\|\partial u_m^k / \partial x^i\|_{L^2}$ ,  $\|\partial u_m^k / \partial v^i\|_{L^2} \to 0$ as  $m \to \infty$ .

First let's take a look at  $\|\partial u_m^k/\partial x^i\|_{L^2}$ . By tucking  $\partial/\partial x^i$  under the integral sign in (5.31), it follows that

$$\frac{\partial u_m^k}{\partial x^i} = \left(\frac{\partial}{\partial x^i}u\right)_m^k.$$

where  $\frac{\partial u}{\partial x^i}$  denotes  $(\frac{\partial u^k}{\partial x^i})\pi^* b_k$ . Thus

$$\left\|\frac{\partial u_m^k}{\partial x^i}\right\|_{L^2}^2 = \left\|\left(\frac{\partial}{\partial x^i}u\right)_m^k\right\|_{L^2}^2 \le \left\|\frac{\partial}{\partial x^i}u\right\|_{L^2}^2 = \frac{1}{\lambda_m^2}\left\|\frac{\partial}{\partial x^i}\Delta^{\pi^*\mathcal{E}}u\right\|_{L^2}^2 \to 0 \qquad \text{as } m \to \infty,$$

where we've used the fact that  $\partial/\partial x^i$  is a smooth vector field over the 0-cosphere bundle and hence  $\partial/\partial x^i (\Delta^{\pi^* \mathcal{E}} u) \in \mathcal{R}^{3-3} \subseteq L^2$ .

Finally, let's take a look at  $\|\partial u_m^k / \partial v^i\|_{L^2}$ . Since we extended  $u_m^k$  to be constant radially, for any fixed  $x \in M$  we have that

$$\int_{S_xM} \left| \frac{\partial u_m^k}{\partial v^i} \right|^2 dS_xM = \int_{\mathbb{S}^n} \left| \frac{\partial u_m^k}{\partial v^i} (x^i, v^i) \right|^2 dv_{\mathbb{S}^n} \le \int_{\mathbb{S}^n} \left| \operatorname{grad}_{v \in \mathbb{S}^n} u_m^k \right|^2 dv_{\mathbb{S}^n}$$

where  $\operatorname{grad}_{v\in\mathbb{S}^n}$  means the gradient in  $(v^i)$  over the Euclidean sphere  $\{(v^1)^2 + \dots (v^n)^2 = 1\}$  with respect to usual spherical metric. Looking at the last integral, integrating by parts gives

$$\int_{\mathbb{S}^n} \left| \operatorname{grad}_{v \in \mathbb{S}^n} u_m^k \right|^2 dv_{\mathbb{S}^n} = \int_{\mathbb{S}^n} u_m^k \left( -\Delta^{\mathbb{S}^n} u_m^k \right) dv_{\mathbb{S}^n} = \lambda_m \int_{S_x M} \left| u_m^k \right|^2 dS_x M.$$

Integrating in x then gives that

$$\left\|\frac{\partial u_m^k}{\partial v^i}\right\|_{L^2}^2 \le \lambda_m \left\|u_m^k\right\|_{L^2}^2 \le \frac{1}{\lambda_m} \left\|\Delta^{\pi^* \mathcal{E}} u\right\|_{L^2}^2 \to 0 \quad \text{as } m \to \infty.$$

As discussed above, this proves the lemma.

#### Proof of Theorem 5.36:

We start by assuming that  $n \neq 2$  since the proof of the case n = 2 requires a slight modification. Comparing Fourier modes of order  $m > \deg f$  in (5.37) gives that

$$X_{+}u_{m-1} + X_{-}u_{m+1} + \Phi u_{m} = 0, (5.48)$$

Using this relation and plugging  $u_{m+1}$  into u in (5.42) gives that

$$\|\mathbb{X}_{+}u_{m+1}\|_{L^{2}}^{2} \ge \|\mathbb{X}_{+}u_{m-1}\|_{L^{2}}^{2} + \|\Phi u_{m}\|_{L^{2}}^{2} + c_{m+1}\|u_{m+1}\|_{L^{2}}^{2} + 2\operatorname{Re}\langle\Phi u_{m}, \mathbb{X}_{+}u_{m-1}\rangle_{L^{2}}.$$
 (5.49)

The idea of the proof is the following. One can plug the relation (5.48) with "m+1" replaced by "m-1" into  $\|\mathbb{X}_+ u_{m-1}\|_{L^2}^2$  in (5.49), use (5.42) again, and then proceed recursively. One will get a long expression on the right which one needs to cleverly manipulated to bound  $\|u_{m_0}\|_{L^2}^2$  for some fixed index  $m_0$ . Then using that  $\|\mathbb{X}_+ u_{m+1}\|_{L^2}^2$  goes to zero as  $m \to \infty$  by Lemma 5.46 will force  $\|u_{m_0}\|_{L^2}^2 = 0$ . We will show that this holds for large enough  $m_0$ , from which the theorem will follow. The first obstacle to accomplishing this is the inner product term  $2\text{Re}\langle \Phi u_m, \mathbb{X}_+ u_{m-1} \rangle_{L^2}$  above. The following claim helps resolve this.

To state the claim, we introduce the following notation. For any  $U \in C^{\infty}(SM; \operatorname{End} \pi^* \mathcal{E})$ (such as  $\Phi$  up to identification), we let  $\mathbb{X}U \in C^{\infty}(SM; \operatorname{End} \pi^* \mathcal{E})$  denote the unique endomorphism field satisfying  $(\mathbb{X}U)h = [\mathbb{X}, U]h$  for any  $h \in C^{\infty}(SM; \pi^* \mathcal{E})$ . We will show in Section 5.5 below that this new operator  $\mathbb{X}$  comes from a connection on endomorphism fields applied to U. In particular, since  $\Phi$  is smooth on  $\overline{M}$  and hence on  ${}^{0}S^*\overline{M}$  (when lifted), it will follow from (5.10) and (5.11) that the quantity  $\mathbb{X}\Phi$  that we use below extends smoothly to  ${}^{0}S^*\overline{M}$ . *Claim:* The following identity is true (because  $\Phi$  is skew-Hermitian):

$$\langle \mathbb{X}_{+}u_{m-1}, \Phi u_{m} \rangle_{L^{2}} + \overline{\langle \mathbb{X}_{+}u_{m-2}, \Phi u_{m-1} \rangle}_{L^{2}} = -\langle u_{m-1}, (\mathbb{X}\Phi)u_{m} \rangle_{L^{2}} - \|\Phi u_{m-1}\|_{L^{2}}^{2}.$$

*Proof of claim:* This is simply a computation:

$$\langle \mathbb{X}_{+}u_{m-1}, \Phi u_{m} \rangle_{L^{2}} = \langle \mathbb{X}u_{m-1}, \Phi u_{m} \rangle_{L^{2}} \quad \text{use } \mathbb{X} = \mathbb{X}_{-} + \mathbb{X}_{+} \text{ and Proposition 5.32}, = -\langle u_{m-1}, \mathbb{X}(\Phi u_{m}) \rangle_{L^{2}} \qquad \text{Lemma 5.8}, \\ = -\langle u_{m-1}, (\mathbb{X}\Phi)u_{m} + \Phi \mathbb{X}u_{m} \rangle_{L^{2}} \qquad \text{definition of } \mathbb{X}\Phi, \\ = -\langle u_{m-1}, (\mathbb{X}\Phi)u_{m} \rangle_{L^{2}} - \langle u_{m-1}, \Phi \mathbb{X}_{-}u_{m} \rangle_{L^{2}} \qquad \mathbb{X} = \mathbb{X}_{-} + \mathbb{X}_{+} \text{ and Proposition 5.32}, \\ \langle u_{m-1}, (\mathbb{X}\Phi)u_{m} \rangle_{L^{2}} + \langle u_{m-1}, \Phi [\Phi u_{m-1} + \mathbb{X}_{+}u_{m-2}] \rangle_{L^{2}} \qquad \text{used (5.48)}.$$

From here the claim follows by rearranging and using that  $\Phi$  is skew-Hermitian.

End of proof of claim.

We return to proving the theorem. The above claim tells us that to get rid of the last term in (5.49), we can add (5.49) and (5.49) with "m" replaced by "m - 1":

$$\begin{aligned} \|\mathbb{X}_{+}u_{m+1}\|_{L^{2}}^{2} + \|\mathbb{X}_{+}u_{m}\|_{L^{2}}^{2} \geq \|\mathbb{X}_{+}u_{m-1}\|_{L^{2}}^{2} + \|\mathbb{X}_{+}u_{m-2}\|_{L^{2}}^{2} + \|\Phi u_{m}\|_{L^{2}}^{2} + \|\Phi u_{m-1}\|_{L^{2}}^{2} \\ + c_{m+1}\|u_{m+1}\|_{L^{2}}^{2} + c_{m}\|u_{m}\|_{L^{2}}^{2} + 2\operatorname{Re}\langle\Phi u_{m},\mathbb{X}_{+}u_{m-1}\rangle_{L^{2}} + 2\operatorname{Re}\langle\Phi u_{m-1},\mathbb{X}_{+}u_{m-2}\rangle_{L^{2}}, \end{aligned}$$

and then apply the equation in the claim to get the following inequality, where for brevity  $a_m = \|X_+ u_m\|_{L^2}^2 + \|X_+ u_{m-1}\|_{L^2}^2$ :

$$a_{m+1} \ge a_{m-1} + \|\Phi u_m\|_{L^2}^2 + \|\Phi u_{m-1}\|_{L^2}^2 + c_{m+1}\|u_{m+1}\|_{L^2}^2 + c_m\|u_m\|_{L^2}^2$$
$$-2\operatorname{Re}\langle u_{m-1}, (\mathbb{X}\Phi)u_m\rangle_{L^2} - 2\|\Phi u_{m-1}\|_{L^2}^2$$
$$\ge a_{m-1} + c_{m+1}\|u_{m+1}\|_{L^2}^2 + c_m\|u_m\|_{L^2}^2 - \|u_{m-1}\|_{L^2}^2 - \|(\mathbb{X}\Phi)u_m\|_{L^2}^2 - \|\Phi u_{m-1}\|_{L^2}^2.$$

Since  $\Phi$  and  $\mathbb{X}\Phi$  are smooth (and hence continuous) on the compact manifold  ${}^{0}S^{*}\overline{M}$ , there exist constants B, C > 0 such that

$$\|\Phi h\|_{L^2}^2 \le B \|h\|_{L^2}^2,$$
$$\|(\mathbb{X}\Phi)h\|_{L^2}^2 \le C \|h\|_{L^2}^2.$$

for any  $h \in C^{\infty}(SM; \pi^* \mathcal{E}) \cap L^2(SM; \pi^* \mathcal{E})$ . Hence

$$a_{m+1} \ge a_{m-1} + r_m \tag{5.50}$$

where

$$r_m := c_{m+1} \|u_{m+1}\|_{L^2}^2 + (c_m - C) \|u_m\|_{L^2}^2 - (1+B) \|u_{m-1}\|_{L^2}^2.$$

Applying (5.50) recursively gives

$$a_{m+1} = a_{m_0-1} + r_{m_0} + r_{m_0+2} + \ldots + r_m$$

for any pair of indices  $m, m_0 > \deg f$  such that  $m = m_0 + 2k$  for some integer  $k \ge 0$ . We seek to bound the resultant tail of  $r_i$ 's. To do so, choose  $m_0$  big enough so that for  $m \ge m_0$ ,  $c_m$  is bigger than both C and B + 1. Hence if  $m = m_0 + 2k$ ,

$$r_{m_0} + r_{m_0+2} + \ldots + r_m = c_{m+1} \|u_{m+1}\|_{L^2}^2 + \sum_{i=0}^{k-1} (c_{m_0+1+2i} - (1+B)) \|u_{m_0+1+2i}\|_{L^2}^2 + \sum_{i=0}^k (c_{m_0+2i} - C) \|u_{m_0+2i}\|_{L^2}^2 - (1+B) \|u_{m_0-1}\| \ge c_{m+1} \|u_{m+1}\|_{L^2}^2 - (1+B) \|u_{m_0-1}\|_{L^2}^2.$$

Hence for any such  $m = m_0 + 2k$  we get that

$$a_{m+1} \ge a_{m_0-1} - (1+B) \|u_{m_0-1}\|_{L^2}^2$$

By the definition of  $a_{m_0-1}$  and (5.42) we have that  $a_{m_0-1} \ge c_{m_0-1} ||u_{m_0-1}||^2_{L^2}$ , and hence we finally arrive at

$$a_{m+1} \ge (c_{m_0-1} - (1+B)) \|u_{m_0-1}\|_{L^2}^2.$$

Assume we defined  $m_0$  before so that  $c_{m_0-1} \ge 1 + B$  as well. Observe that  $a_{m+1} \to 0$  as  $m \to \infty$  by Lemma 5.46. Hence we get that  $||u_{m_0-1}||_{L^2}^2 = 0$  and thus  $u_{m_0-1} = 0$  for all such large enough  $m_0$ . This proves the theorem in the case  $n \neq 2$ .

Finally, let's discuss the modification needed in the case n = 2. In this case we instead use the second equality in (5.42) and hence proceeding as above arrive at that for sufficiently large m

$$d_m a_{m+1} \ge a_{m-1} + r_m,$$

where we've used that  $d_m \ge d_{m+1}$  and that all  $d_i \ge 1$  for  $i \ge 1$ . Multiplying through by  $d_{m-2}$ , applying the same inequality with "m" replaced by "m - 2" on the right-hand side, and then repeating recursively gives

$$(d_m \cdot \ldots \cdot d_{m_0})a_{m+1} = a_{m_0-1} + r_{m_0} + (d_{m_0})r_{m_0+2} + \ldots + (d_{m-2} \cdot \ldots \cdot d_{m_0})r_m$$

for any pair of  $m, m_0 > \deg f$  such that  $m = m_0 + 2k$  for some integer k. Since the  $d_i \ge 1$ , we get the inequality

$$\left(\prod_{i=0}^{k} d_{m-2i}\right)a_{m+1} \ge a_{m_0-1} + r_{m_0} + r_{m_0+2} + \ldots + r_m.$$

Again by the definition of  $a_{m_0-1}$  and (5.42) we have that  $a_{m_0} \ge (c_{m_0-1}/d_{m_0-1}) ||u_{m_0-1}||_{L^2}^2$  and so

$$\left(\prod_{i=0}^{k} d_{m-2i}\right) a_{m+1} \ge \left(\frac{c_{m_0-1}}{d_{m_0-1}} - (1+B)\right) \|u_{m_0-1}\|_{L^2}^2.$$

Since the  $d_m \to 1$  as  $m \to \infty$ , we can assume that we defined  $m_0$  before so that  $c_{m_0-1}/d_{m_0-1} \ge B + 1$  as well. Furthermore,  $\prod_{m=1}^{\infty} d_m$  converges by the infinite product criteria since  $\sum_{m=1}^{\infty} (d_m - 1) < \infty$  and so the coefficient on the left-hand side is bounded by some fixed constant. Hence, as before, the theorem follows from the fact that  $a_{m+1} \to 0$  as  $m \to \infty$ .

Theorem 5.36 has one disadvantage. Though it tells us that f being of finite degree implies that the solution u is also of finite degree, it gives no information about the degree of u itself. The following proposition remedies this by assuming an additional condition on the metric g and connection  $\nabla^{\mathcal{E}}$ .

**Definition 5.51.** For any index  $m \ge 0$ , we call elements of ker  $\mathbb{X}_+|_{\Omega_m}$  twisted conformal *Killing tensors (CKTs)* of degree m. A nontrivial twisted CKT is a twisted CKT of degree  $m \ge 1$  that is not identically zero.

**Proposition 5.52.** Assume that the sectional curvatures of g are negative. Suppose also that there are no nontrivial twisted CKTs in  $\mathcal{R}^3(SM; \pi^*\mathcal{E})$ . If  $u \in \mathcal{R}^3(SM; \pi^*\mathcal{E})$  solves

$$Xu + \Phi u = f$$

for some  $f \in C^{\infty}(SM; \pi^* \mathcal{E})$  of finite degree m, then u is of degree max  $\{m - 1, 0\}$ .

**Proof:** We know by Theorem 5.36 that u has finite degree, call it  $m' \ge 0$ . If m' = 0, then we're done. So suppose that  $m' \ge 1$ . We will prove this theorem by contradiction: suppose

that  $m' \ge m$ . Then comparing the Fourier modes of order m' + 1 of both sides of the above equation gives

$$\mathbb{X}_+ u_{m'} = 0.$$

Hence,  $u_{m'}$  is a twisted CKT that is in  $\mathcal{R}^3(SM; \pi^*\mathcal{E})$  by Proposition 5.34. Since we assumed that there are no such nontrivial CKTs, we conclude that  $u_m$  is identically zero. But this contradicts that the degree of u is m', and hence proves the proposition.

### 5.4 Regularity of Solutions to the Transport Equation

Before we prove the main result of our paper, we need to establish the regularity of solutions to the transport equation of a specific form. Here we use the material that we introduced in Section 2.5 up to (2.15) there and the two sentences after. We remind the reader that a map  $A: T\overline{M} \to \mathcal{E}$  is called a *bundle homomorphism* if for all  $x \in \overline{M}$ , A restricts to a linear map  $A: T_x\overline{M} \to \mathcal{E}_x$ . For such bundle homomorphisms A, we often let A also denote its restriction  $A: SM \to \mathcal{E}_{|SM|}$ .

**Definition 5.53.** If  $N \ge 0$  is an integer, we say that a "bundle homomorphism  $A : T\overline{M} \to \mathcal{E}$ is such that its entries are in  $\rho^N C^{\infty}(\overline{M})$  in any boundary coordinates (of  $\overline{M}$ ) and frame (for  $\mathcal{E}$ )" if the following holds. If one takes any boundary coordinates  $(x^i)$  of  $\overline{M}$  and any frame  $(b_k)$  of  $\mathcal{E}$  over these coordinates' domain, locally  $Av = A_i^k v^i b_k$  with each  $A_{ij}^k \in \rho^N C^{\infty}(\overline{M})$ where the  $v^i$  are v's components with respect to  $(\partial/\partial x^i)$ .

The following is our regularity theorem:

**Proposition 5.54.** Assume that (M, g) is nontrapping and that  $N \ge 0$  is an integer. Suppose that  $\Phi \in \rho^{N+1}C^{\infty}(\overline{M}; \operatorname{End}_{\operatorname{Sk}}\mathcal{E})$  and that we have a  $\phi \in \rho^{N+1}C^{\infty}(\overline{M}; \mathcal{E})$  and a bundle homomorphism  $A: TM \to \mathcal{E}$  such that its entries are in  $\rho^N C^{\infty}(\overline{M})$  in any boundary coordinates and frame (in the sense of Definition 5.53). Suppose also that the connection symbols of  $\nabla^{\mathcal{E}}$  are in  $\rho^N C^{\infty}(\overline{M}; \mathbb{R})$  in any boundary coordinates and frame (in the sense of Definition 2.6). Then for any given boundary data  $h \in C^{\infty}(\partial_{-}{}^{b}S^{*}\overline{M}; \pi_{b}^{*}\mathcal{E}|_{\partial_{-}{}^{b}S^{*}\overline{M}})$ , there exists a unique solution  $u \in C^{\infty}({}^{b}S^{*}\overline{M}; \pi_{b}^{*}\mathcal{E})$  to

$$Xu + \Phi u = \phi + A \quad \text{on } SM, \tag{5.55}$$

with  $u|_{\partial_{-}^{b}S^{*}\overline{M}} = h.$ 

Furthermore, for any  $l \ge 0$  there exists an  $N_0 \ge 0$  dependent only on (M, g) and l such that the following holds. If  $N \ge N_0$ , then for all solutions u as above that also satisfy  $h \equiv 0$  and  $u|_{\partial_+S^*\overline{M}} \equiv 0$ , it holds that  $u \in \mathcal{R}^l(SM; \pi^*\mathcal{E})$ .

**Proof:** Recall that  $d = \operatorname{rank} \mathcal{E}$ . From Lemma 2.1 in [14] we have that  $X = \rho \overline{X}$  for some smooth vector field  $\overline{X}$  over  ${}^{b}S^{*}\overline{M}$  that is nonvanishing and transverse to  $\partial^{b}S^{*}\overline{M}$ . Let  $(\rho, y^{\mu}) =$  $(x^{i})$  be asymptotic boundary normal coordinates of  $\overline{M}$ ,  $(b_{k})$  a frame for  $\mathcal{E}$  over their domain, and consider the coordinates  ${}^{v_{i}\partial}/\partial x^{i} \mapsto (x^{i}, v^{i})$  of TM. Let  ${}^{\mathcal{E}}\Gamma^{k}_{ij}$  denote the connection symbols of  $\nabla^{\mathcal{E}}$  with respect to  $(\partial/\partial x^i)$  and  $(b_k)$ . Then observe that the components of (5.55) with respect to  $(\pi^* b_k)$  and  $(\partial/\partial x^i)$  are given by

$$Xu^k + {}^{\mathcal{E}}\Gamma^k_{ij}v^i u^j + \Phi^k_j u^j = \phi^k + A^k_i v^i$$
(5.56)

where  $k = 1, \ldots, d$ . Dividing through by  $\rho$  gives

$$\overline{X}u^{k} + (\rho^{-1})^{\mathcal{E}}\Gamma^{k}_{ij}v^{i}u^{j} + \rho^{-1}\Phi^{k}_{j}u^{j} = \rho^{-1}\phi^{k} + \rho^{-1}A^{k}_{i}v^{i}.$$
(5.57)

Take the coordinates  $\eta_0 {}^{d\rho}/_{\rho} + \eta_{\lambda} dy^{\lambda} \mapsto (x^i, \eta_i)$  of  ${}^{b}T^*\overline{M}$  and observe that the canonical identification  $\sharp \circ (F^*)^{-1} : {}^{b}T^*\overline{M}|_M \to TM$  is given by

$$v^{i} = g^{ii'} \begin{cases} \eta_{0}/\rho & \text{if } i' = 0\\ \eta_{i'} & \text{if } i' \ge 1 \end{cases}$$
(5.58)

We denote the right-hand side by  $g^{ii'}\{\eta_{i'}\}$ . Pulling the above equation (5.57) to  ${}^{b}S^*\overline{M}$  (in the sense of Remark 4.10) gives

$$\overline{X}u^{k} + \rho^{-1\mathcal{E}}\Gamma^{k}_{ij}g^{ii'}\{\eta_{i'}\}u^{j} + \rho^{-1}\Phi^{k}_{j}u^{j} = \rho^{-1}\phi^{k} + \rho^{-1}A^{k}_{i}g^{ii'}\{\eta_{i'}\}.$$
(5.59)

We remind the reader that each  $g^{ij}$  is  $\rho^2$  times something smooth on  $\overline{M}$ . Hence it follows from our assumptions that all of the terms in this differential equation are smooth on  ${}^{b}S^*\overline{M}$ .

Since we assumed that g is nontrapping, it follows from the proof of Corollary 2.5 in [14] that any maximal integral curve<sup>8</sup>  $\sigma$  of  $\overline{X}$  is of the form  $\sigma : [a, b] \to {}^{b}S^{*}\overline{M}$  where a and b are finite with  $\sigma(a) \in \partial_{-}{}^{b}S^{*}\overline{M}$  and  $\sigma(b) \in \partial_{+}{}^{b}S^{*}\overline{M}$ . Hence, (5.59) can be viewed as ordinary differential equations (ODEs) along such curves  $\sigma$ . Hence, since  $\overline{X}$  is nonvanishing and transverse to  $\partial^{b}S^{*}\overline{M}$ , it follows from the theory of flows and the existence, uniqueness, and smooth dependence on initial condition of linear ODEs (see [5] and [23]) that indeed a unique smooth solution u exists to (5.59) and hence (5.55) satisfying the given boundary data.

Next suppose that  $h \equiv 0$  and  $u|_{\partial_+S^*\overline{M}} \equiv 0$ . If  $l \ge 1$ , suppose that we also have smooth vector fields  $V_1, ..., V_l \in C^{\infty}({}^0S^*\overline{M}; T^0S^*\overline{M})$  over the 0-cosphere bundle. Pick any point  $x_0 \in \partial M$  contained in our coordinates  $(x^i)$ . We will show that (5.2) holds for some compact neighborhood K of  $x_0$  (i.e. K has nonempty interior), from which it will quickly follow that  $u \in \mathcal{R}^l$ . Throughout the proof we will make  $N \ge 0$  is as big as we need whenever we need it. At the end of the proof, we will discuss why there exists a maximum upper bound on the size of N that we need that is dependent only on (M, g) and l, and hence demonstrate the existence of  $N_0$ .

We begin by establishing a few facts about the flow of X. Let  $\varphi : SM \times \mathbb{R} \to SM$  denote the flow of X. For any point  $\zeta = \eta_0 d^{\rho}/_{\rho} + \eta_\lambda dy^\lambda \in {}^{b}S^*M$  we write its identified point on SM as  $z \in SM$ . In [14] the authors explain that close enough to the boundary, the flow  $\varphi$ moves away from the boundary when  $\eta_0 > 0$  and towards the boundary when  $\eta_0 < 0$ . More precisely, by Lemma 2.3 in [14] and its proof there exists constants  $C, \varepsilon > 0$  such that if we take any  $z \in SM \cap \{\rho < \varepsilon\}$  and write  $\rho(t) = \rho \circ \varphi_z(t)$ ,

<sup>&</sup>lt;sup>8</sup>i.e. integral curve whose interval domain cannot be extended.

- 1. if  $\eta_0 \geq 0$  then  $\lim_{t \to -\infty} \varphi_z(t) \in \partial_{-}{}^{b}S^*\overline{M}$ ,  $\rho(t) \leq Ce^t$ , and  $\rho(t)$  is increasing,
- 2. if  $\eta_0 < 0$  then  $\lim_{t \to +\infty} \varphi_z(t) \in \partial_+{}^b S^* \overline{M}$ ,  $\rho(t) \leq C e^{-t}$ , and  $\rho(t)$  is decreasing.

By the same lemma and its proof, it follows that there exist compact neighborhoods  $K, K' \subseteq \{\rho < \varepsilon\}$  of  $x_0$  in  $\overline{M}$  satisfying  $K \subseteq K' \subseteq \text{dom}(x^i)$  such that for any  $z \in \pi^{-1}[K]$ ,  $\varphi_z$  will always be contained in  $\pi^{-1}[K']$  for  $t \ge 0$  if  $\eta_0 \ge 0$  or  $t \le 0$  if  $\eta_0 < 0$ . Our first goal is to show that each  $u^k \in \rho^{(n+1)/2} L^{\infty}[K]$  where our approach will be to

Our first goal is to show that each  $u^k \in \rho^{(n+1)/2} L^{\infty}[K]$  where our approach will be to study the growth of the solution to (5.56) by writing that equation as an ODE along integral curves of X. Fix any  $\zeta \cong z \in \pi^{-1}[K]$ . Suppose that  $\eta_0 \ge 0$  since the proof below is essentially the same for the case  $\eta_0 < 0$ . We set B and b to be the  $d \times d$  matrix and  $d \times 1$ column vector respectively given by

$$B_i^k = \left({}^{\mathcal{E}}\Gamma_{ij}^k v^i + \Phi_j^k\right) \qquad \text{and} \qquad b^k = \phi^k + A_i^k v^i, \tag{5.60}$$

and we denote  $B(t) = B \circ \varphi_z(t)$ ,  $b(t) = b \circ \varphi_z(t)$ , and  $u(t) = u \circ \varphi_z(t)$ . This way, (5.56) along  $\varphi_z$  is given by the ODE

$$\frac{du^k}{dt} + B^k_i u^i = b^k. ag{5.61}$$

Since  $u|_{\partial^b S^*\overline{M}} \equiv 0$  and  $\lim_{t\to\infty} \varphi_z(t) \in \partial_-{}^b S^*\overline{M}$ , we have that our solution u satisfies the "initial condition"  $\lim_{t\to\infty} u(t) = 0$ . Now, for any matrix M or column vector w, let |M| and |w| denote the norms  $\sum_{ik} |M_i^k|$  and  $\sum_i |w^i|$ . Fixing any time  $t_0 < 0$ , we get from (5.61), the fundamental theorem of calculus, and the triangle inequality that

$$|u(t)| \le |u(t_0)| + \int_{t_0}^t |B(s)| |u(s)| ds + \int_{t_0}^t |b(s)| ds.$$
(5.62)

We will use this to bound |u(0)| = |u(z)| in terms of a power of  $\rho$ . We employ the standard technique in ODEs of defining the function  $R : (-\infty, 0] \to \mathbb{R}$  given by the first integral on the right-hand side. Whenever  $|B(t)| \neq 0$ ,

$$\frac{1}{|B(t)|}R'(t) \le |u(t_0)| + R(t) + \int_{t_0}^t |b(s)| ds$$

and so

$$R'(t) \le |B(t)|R(t) + |B(t)| \left( |u(t_0)| + \int_{t_0}^t |b(s)| ds \right)$$

Observe that this inequality also holds when |B(t)| = 0 and hence for all  $t \leq 0$ . This is a separable equation. In particular, if we take |B(t)|R(t) to the left-hand side, multiply through by  $\exp\left[-\int_{t_0}^t |B(s)|ds\right]$ , integrate from  $t = t_0$  to t = 0 (using that  $R(t_0) = 0$ ), and finally divide through by  $\exp\left[-\int_{t_0}^0 |B(s)|ds\right]$ , we will get

$$R(0) = \int_{t_0}^0 |B(s)| |u(s)| ds$$
(5.63)

$$\leq e^{\int_{t_0}^0 |B(s)|ds} \int_{t_0}^0 e^{-\int_{t_0}^t |B(s)|ds} |B(t)| \left( |u(t_0)| + \int_{t_0}^t |b(s)|ds \right) dt.$$

Before we let  $t_0 \to -\infty$ , let us discuss integrability. Because we're working on the sphere bundle and  $g = \rho^{-2}\overline{g}$  for a smooth metric  $\overline{g}$ , there exists a constant  $C_2 > 0$  such that the magnitude of  $v^i$  in the expressions for B and b in (5.60) are less than  $C_2\rho$  on  $\pi^{-1}[K']$ . Hence by (5.60) we have that there exists constants  $C_3, C_N > 0$  such that  $|B| \leq C_3\rho$  and  $|b| \leq C_N \rho^{N+1}$  on  $\pi^{-1}[K']$  where  $N \geq 0$  is an integer whose size is to be determined later (i.e.  $C_N$  depends on N). In particular, since  $\rho(t) \leq Ce^t$  for  $t \leq 0$  we have that all of the integrals on the right-hand side of (5.63) converge. Hence letting  $t_0 \to -\infty$  in (5.63) gives that

$$\int_{-\infty}^{0} |B(s)| |u(s)| ds \le e^{\int_{-\infty}^{0} |B(s)| ds} \int_{-\infty}^{0} |B(t)| \int_{-\infty}^{t} |b(s)| ds dt$$
(5.64)

where we've used that  $\lim_{t\to\infty} u(t) = 0$  and that  $\exp\left[-\int_{t_0}^t |B(s)|ds\right] \leq 1$ . To estimate the right-hand side further, observe that since  $\rho(t)$  is increasing on  $t \leq 0$  we have that  $|b(t)| \leq C_N \rho(t) \rho^N(0) \leq C_N C e^t \rho^N(0)$  and so

$$\int_{-\infty}^{0} |b(s)| ds \le C_N C \rho^N(0).$$
(5.65)

Doing a similar sort of thing for |B(t)|, we get from (5.64) that

$$\int_{-\infty}^{0} |B(s)| |u(s)| ds \le e^{C_3 C} C_3 C C_N C \rho^N(0).$$
(5.66)

Finally, letting  $t_0 \to -\infty$  in (5.62) and plugging (5.65) and (5.66) into there finally gives us that for some constant  $C'_N > 0$ ,

$$|u(z)| \le C'_N \rho^N(z) \tag{5.67}$$

for all  $z \in \pi^{-1}[K]$ , where we've used that u(0) = u(z). If we require  $N \ge \frac{(n+1)}{2}$ , then indeed we have that each  $u^k \in \rho^{(n+1)/2} L^{\infty}[K]$ .

Next, if  $l \ge 1$  we show that  $V_1 u^k \in \rho^{(n+1)/2} L^{\infty}[K]$ , after which it should be clear how the cases of the higher order derivatives follow (i.e.  $V_2 V_1 u^k$ , ...) if  $l \ge 2$ . We do this by applying  $V_1$  to the ODE (5.61) and study the growth of the solution. Applying  $V_1$  to (5.61) and rearranging, we obtain

$$\frac{d[V_1(u^k \circ \varphi)]}{dt} + (B_i^k \circ \varphi)V_1(u^k \circ \varphi) = V_1(b^k \circ \varphi) - V_1(B_i^k \circ \varphi)(u^i \circ \varphi).$$
(5.68)

Considering that in form this is a similar ODE for  $V_1(u^k \circ \varphi)$  as (5.61) is for  $u^k$ , up to a few details we describe below, a similar proof as for (5.67) shows that for some constants  $C''_N > 0$ 

$$|V_1(u \circ \varphi)(z, 0)| = |V_1u(z)| \le C_N'' \rho^N(z)$$

for all  $z \in \pi^{-1}[K]$  and hence indeed each  $V_1 u^k \in \rho^{(n+1)/2} L^{\infty}[K]$  if we require that  $N \ge \frac{(n+1)}{2}$ . The only analogous steps that we're missing and need to show is that  $V_1(u^k \circ \varphi)$  satisfies the "initial condition"

$$\lim_{t \to -\infty} V_1(u^k \circ \varphi) = 0 \tag{5.69}$$

and that for any integer  $N' \ge 0$  whose size is to be determined later there exists a constant  $C_{N'} > 0$  such that the right-hand side of (5.68) satisfies

$$\left|V_1(b^k \circ \varphi) - V_1(B_i^k \circ \varphi)(u^i \circ \varphi)\right| \le C_{N'} \rho^{N'+1} \circ \varphi$$
(5.70)

for all  $z \in \pi^{-1}[K']$ .

By shrinking everything if needed, we assume that there is another compact subset K'' of our coordinates' domain such that  $K' \subseteq K'' \subseteq \text{dom}(x^i)$  and such that for any  $z \in \pi^{-1}[K']$ ,  $\varphi_z$  will always be contained in  $\pi^{-1}[K'']$  for  $t \ge 0$  if  $\eta_0 \ge 0$  or  $t \le 0$  if  $\eta_0 < 0$ . Throughout the rest of the proof, we again assume that we're working with points such that  $\eta_0 \ge 0$  because the proof is similar for the case  $\eta_0 < 0$ . We begin with showing the "initial condition" (5.69). Let  $(\varphi^i, \varphi_i)$  denote the components of  $\varphi$  in the coordinates  $(x^i, \eta_i)$  of  ${}^{b}T^*\overline{M}$ . Then over the interior of  ${}^{b}S^*\overline{M}$ , we have by the chain rule that

$$V_1(u^k \circ \varphi) = \left(\frac{\partial u^k}{\partial x^i} \circ \varphi\right) V_1 \varphi^i + \left(\frac{\partial u^k}{\partial \eta_i} \circ \varphi\right) V_1 \varphi_i.$$
(5.71)

Since  $u|_{\partial^b S^*\overline{M}} \equiv 0$ , and by (5.67) u vanishes like  $\rho^N$  at the boundary  $\partial^b S^*\overline{M}$  where we can require  $N \geq 2$ , all of  $u^k$ 's partials vanish at the boundary like  $\rho^{N-1}$ . In particular, for fixed  $z \in \pi^{-1}[K]$ ,

$$\left(\frac{\partial u^k}{\partial x^i} \circ \varphi\right), \left(\frac{\partial u^k}{\partial \eta_i} \circ \varphi\right) \in O(e^{(N-1)t}) \quad \text{as } t \to -\infty.$$
(5.72)

So let us take a look at size of the terms  $V_1 \varphi^i$  and  $V_1 \varphi_i$  in (5.71). Consider the coordinates  $\overline{\eta}_i^{dx^i}/\rho \mapsto (x^i, \overline{\eta}_i)$  of  ${}^0T^*\overline{M}$  (note the bars to distinguish these from our coordinates of  ${}^bT^*\overline{M}$  above) and let  $(\widetilde{\varphi}^i, \widetilde{\varphi}_i)$  denote the components of  $\varphi$  with respect to these coordinates. Since canonical identification is given by  $x^i = x^i$ ,  $\eta_0 = \overline{\eta}_0$  and  $\eta_\lambda = \rho^{-1}\overline{\eta}_\lambda$  for  $\lambda = 1, \ldots, n$ , we have that by identification  $\varphi^i = \widetilde{\varphi}^i$ ,  $\varphi_0 = \widetilde{\varphi}_0$ , and  $\varphi_\lambda = (\widetilde{\varphi}^0)^{-1}\widetilde{\varphi}_\lambda$  for  $\lambda = 1, \ldots, n$  where we point out that  $\widetilde{\varphi}^0 = \rho \circ \widetilde{\varphi}$ . Hence

$$V_1 \varphi^i = V_1 \widetilde{\varphi}^i, \qquad V_1 \varphi_0 = V_1 \widetilde{\varphi}_0, \qquad (5.73)$$

$$V_1\varphi_{\lambda} = -(\rho \circ \varphi)^{-2}V_1(\widetilde{\varphi}^0)\widetilde{\varphi}_{\lambda} + (\rho \circ \varphi)^{-1}V_1\widetilde{\varphi}_{\lambda} \quad \text{for } \lambda = 1, \dots, n.$$
 (5.74)

We point out that  $\tilde{\varphi}_{\lambda}$  are bounded over  $\pi^{-1}[K'']$  because they are smooth over the compact  $\pi_0^{-1}[K'']$ . Now, we showed in (5.11) that X extends as a smooth vector field on  ${}^0S^*\overline{M}$ . Hence, as is noted at the end of the proof of Lemma 3.13 in [14], X has a Lipschitz constant uniformly bounded on the compact  ${}^0S^*\overline{M}$ . Since  $\tilde{\varphi}$  is the flow of X over  ${}^0S^*\overline{M}$ , as the authors explain there it follows by Grönwall's inequality that there exist constants  $C_4, c_4 > 0$  such that all of the partials

$$\left|\frac{\partial \widetilde{\varphi}^{i}}{\partial x^{i}}\right|, \left|\frac{\partial \widetilde{\varphi}^{i}}{\partial \overline{\eta}_{i}}\right|, \left|\frac{\partial \widetilde{\varphi}_{i}}{\partial x^{i}}\right|, \left|\frac{\partial \widetilde{\varphi}_{i}}{\partial \overline{\eta}_{i}}\right| \le C_{4} e^{-c_{4}t}$$

$$(5.75)$$

on (z,t) for all  $z \in \pi^{-1}[K'] : \eta_0 \ge 0$  and  $t \le 0$ . Next, if we write in coordinates  $V_1 = (V_1)^{i\partial}/\partial x^i + (V_1)_i^{\partial}/\partial \overline{\eta}_i$  over  ${}^{0}S^*\overline{M}$ , we have that the components  $(V_1)^i$  and  $(V_1)_i$  are also bounded over  $\pi^{-1}[K']$  because they are smooth over the compact  $\pi_0^{-1}[K']$ . Hence there exists a constant  $C_5 \ge 0$  such that

$$\left|V_1\widetilde{\varphi}^i\right| \le C_5 e^{-c_4 t} \quad \text{and} \quad \left|V_1\widetilde{\varphi}_i\right| \le C_5 e^{-c_4 t}$$

$$(5.76)$$

on (z,t) for all  $z \in \pi^{-1}[K']$ :  $\eta_0 \ge 0$  and  $t \le 0$ . Plugging this into (5.73) and (5.74), and then into (5.71), we see that choosing  $N \ge 0$  big enough in (5.72) will make the "initial conditions" (5.69) hold (in particular, any  $N > c_4 + 1$  will work).

Finally, let's show that (5.70) holds. By (5.67), we already have that  $|(u^i \circ \varphi)| \leq C'_N \rho^N \circ \varphi$ for any  $N \geq 0$ . Next, the canonical identification  $TM \cong {}^0T^*\overline{M}$  is given by  $v^i = \rho^{-1}g^{ii'}\overline{\eta}_{i'}$ and so by (5.60) we have that over  ${}^0S^*\overline{M}$ 

$$B_i^k = \begin{pmatrix} \varepsilon \Gamma_{ij}^k \rho^{-1} g^{ii'} \overline{\eta}_{i'} + \Phi_j^k \end{pmatrix} \quad \text{and} \quad b^k = \phi^k + A_i^k \rho^{-1} g^{ii'} \overline{\eta}_{i'}.$$

In particular, we see that these are  $\rho^{N+1}$  times something smooth on  ${}^{0}S^{*}\overline{M}$ . Hence, the partials  $\partial B_{i}^{k}/\partial x^{l}$ ,  $\partial B_{i}^{k}/\partial \overline{\eta}_{l}$ ,  $\partial b^{k}/\partial x^{l}$ , and  $\partial b^{k}/\partial \overline{\eta}_{l}$  are all  $\rho^{N}$  times something smooth on  ${}^{0}S^{*}\overline{M}$ . In particular,

$$\left|\frac{\partial B_i^k}{\partial x^l} \circ \varphi\right|, \left|\frac{\partial B_i^k}{\partial \overline{\eta}_l} \circ \varphi\right|, \left|\frac{\partial b^k}{\partial x^l} \circ \varphi\right|, \left|\frac{\partial b^k}{\partial \overline{\eta}_l} \circ \varphi\right| \le C_6 \rho^N \circ \varphi \tag{5.77}$$

on (z,t) for all  $z \in \pi^{-1}[K']$ :  $\eta_0 \ge 0$  and  $t \le 0$ . Now, over  ${}^0S^*\overline{M}$  we have that

$$V_1(B_i^k \circ \varphi) = \left(\frac{\partial B_i^k}{\partial x^l} \circ \varphi\right) V_1 \widetilde{\varphi}^l + \left(\frac{\partial B_i^k}{\partial \overline{\eta}_l} \circ \varphi\right) V_1 \widetilde{\varphi}_l,$$
  

$$V_1(b^k \circ \varphi) = \left(\frac{\partial b^k}{\partial x^l} \circ \varphi\right) V_1 \widetilde{\varphi}^l + \left(\frac{\partial b^k}{\partial \overline{\eta}_l} \circ \varphi\right) V_1 \widetilde{\varphi}_l.$$
(5.78)

Hence if in (5.77) we write  $\rho^N \circ \varphi \leq Ce^{c_4t} (\rho^{N-c_4} \circ \varphi)$  it follows from (5.76) and (5.78) that requiring  $N \geq c_4 + N' + 1$  will make (5.70) holds. As discussed above, this completes the proof that  $V_1 u^k \in \rho^{(n+1)/2} L^{\infty}[K]$ .

As we mentioned above, if  $l \geq 2$  the proof that  $V_2V_1u^k \in \rho^{(n+1)/2}L^{\infty}[K]$  and similarly for the higher derivatives follows similarly, starting with applying  $V_2$  to (5.68). We end the proof with a discussion of how we can ensure that the  $N \geq 0$  that we used has an upper bound dependent only on (M, g) and l. Throughout the proof we required that N is bigger than fixed numbers (e.g. 2), numbers dependent on the dimension (e.g. (n+1)/2), the constant  $c_4$  in (5.75), and the analogs of  $c_4$  in (5.75) when carrying out the proof for higher order derivatives (i.e.  $V_2V_1u^k$ , ...). We explain why the latter two can be bounded. Due to the compactness of the boundary  $\partial \overline{M}$ , we can cover  $\partial \overline{M}$  with a finite collection of neighborhoods of the form K, K', and K'' as described above which in turn determine their own  $c_4$ 's in (5.75) and their higher derivative analogs using only the geometry of the geodesic flow  $\varphi$ . Hence, taking the maximum of all such  $c_4$ 's and their higher derivative analogs will provide us with an upper bound for N that works throughout the whole proof. In particular, this upper bound for N will be dependent only on (M, g) and l, and hence proves the existence of the claimed  $N_0$  stated in the theorem.

#### 5.5 Proof of Theorem 2.7

In our proof, we will assume that the  $N_0 \ge 0$  in the statement of the theorem is as big as we need. It will be clear that the size of  $N_0$  that we will need is dependent only on (M, g). The first step is to provide a formulation of (2.3) as a single transport equation of endomorphism fields over SM. To begin, we define a natural connection " $\nabla^{\text{En}}$ " on endomorphism fields End  $\mathcal{E}$  as follows. For any  $U \in C^{\infty}(\overline{M}; \text{End }\mathcal{E})$  and any  $v \in T_x \overline{M}$ , we define  $\nabla_v^{\text{En}} U$  to be the unique element of End  $\mathcal{E}_x$  satisfying

$$\left(\nabla_v^{\operatorname{En}}U\right)h = \left[\nabla_v^{\mathcal{E}}, U\right]h,$$

for any  $h \in C^{\infty}(\overline{M}; \operatorname{End} \mathcal{E})$ . To see what this looks like in coordinates, take coordinates  $(x^i)$  of M, a frame  $(b_i)$  for  $\mathcal{E}$  over their domain, and let  ${}^{\mathcal{E}}\Gamma^k_{ij}$  denote the connection symbols of  $\nabla^{\mathcal{E}}$  with respect to  $(\partial/\partial x^i)$  and  $(b_k)$ . A quick computation shows that

$$\nabla_v^{\rm En} U = vU + ({}^{\mathcal{E}}\Gamma)U - U({}^{\mathcal{E}}\Gamma), \qquad (5.79)$$

where on the right-hand side U is thought of as a matrix in the basis  $(b_i)$ , vU denotes applying v to every entry of U, and  ${}^{\mathcal{E}}\Gamma$  represents the matrix with entry  ${}^{\mathcal{E}}\Gamma_{ij}^k v^i$  in the  $k^{\text{th}}$ row and  $j^{\text{th}}$  column. This is considered a natural connection on End  $\mathcal{E}$  because it satisfies the product rule  $\nabla_v^{\mathcal{E}}(Uh) = (\nabla_v^{\text{En}}U)h + U\nabla_v^{\mathcal{E}}h$ .

Of more importance to us is the pullback bundle  $\pi^* \operatorname{End} \mathcal{E}$  on SM and its connection which we denote by  $\nabla^{\pi^*\operatorname{En}} := \pi^* \nabla^{\operatorname{En}}$ . We analogously define the operator  $\mathbb{X} = \nabla^{\pi^*\operatorname{En}}_X$ , using context to differentiate it from our other operators also denoted by "X." We remark that this new X satisfies all of the same properties that we proved earlier about X because we're simply considering a different vector bundle here that's also smooth over  $\overline{M}$  (i.e.  $\operatorname{End} \mathcal{E}$ ).

By looking in local coordinates, it's easy to see that  $\widetilde{\nabla}^{\mathcal{E}} = \nabla^{\mathcal{E}} + A$  where  $A \in C^{\infty}(T\overline{M}; \operatorname{End}_{\operatorname{sk}}\mathcal{E})$  is a bundle homomorphism whose entries are in  $\rho^{N_0}C^{\infty}(\overline{M})$  in any boundary coordinates and frame (in the sense of Definition 5.53). Consider solutions U and  $\widetilde{U}$  to the following transport equations on SM:

$$\begin{cases} \mathbb{X}U + \Phi U = 0, & U|_{\partial_{-}{}^{b}S^{*}\overline{M}} = \mathrm{id}, \\ \mathbb{X}\widetilde{U} + A\widetilde{U} + \widetilde{\Phi}\widetilde{U} = 0, & \widetilde{U}\Big|_{\partial_{-}{}^{b}S^{*}\overline{M}} = \mathrm{id}, \end{cases}$$
(5.80)

which exist by Proposition 5.54 (using a different connection and Higgs field) which we can apply to the second equation because " $X\tilde{U} + A\tilde{U}$ " can be thought of as " $(\pi^*\nabla)_X\tilde{U}$ " for the connection  $\nabla = \nabla^{\text{En}} + A$ .

We demonstrate the usefulness of U and  $\widetilde{U}$ . Suppose that  $\gamma : (-\infty, \infty) \to M$  is a complete unit-speed geodesic and that  $u : (-\infty, \infty) \to \mathcal{E}$  is the smooth solution along  $\gamma$  to the initial value problem

$$\nabla_{\dot{\gamma}(t)}^{\mathcal{E}} u(t) = 0, \quad \lim_{t \to -\infty} u(\gamma(t)) = e, \tag{5.81}$$

where e is an element in  $\mathcal{E}_{x_0}$  where  $x_0 \in \partial \overline{M}$  is the limit of  $\gamma(t)$  as  $t \to -\infty$ . We point out that such a u exists by Lemma 2.5 with  $\Phi = 0$ . We claim that  $(U \circ \sigma)u(t)$  and  $(\widetilde{U} \circ \sigma)u(t)$ are solutions to (2.3) and (2.3) with  $\Phi$  and  $\nabla^{\mathcal{E}}$  replaced by  $\widetilde{\Phi}$  and  $\widetilde{\nabla}^{\mathcal{E}}$  respectively.

Let  $\sigma : (-\infty, \infty) \to SM$  be the integral curve of X satisfying  $\gamma = \pi \circ \sigma$  and let  $\hat{u}$  denote u lifted to  $\sigma$  via the canonical identification of  $\mathcal{E}$  and  $\pi^* \mathcal{E}$ . It follows from (4.8) that  $\nabla_{\hat{\sigma}}^{\pi^* \mathcal{E}} \hat{u} \cong \nabla_{\hat{\gamma}}^{\mathcal{E}} u = 0$ . Thus

$$\nabla^{\mathcal{E}}_{\dot{\gamma}}((U \circ \sigma)u) + \Phi((U \circ \sigma)u) = \left[\nabla^{\mathrm{En}}_{\dot{\gamma}}(U \circ \sigma)\right]u + (U \circ \sigma)\nabla^{\mathcal{E}}_{\dot{\gamma}}u + \Phi((U \circ \sigma)u)$$

$$\cong \left[\nabla_X^{\pi^* \operatorname{En}} U\right] \hat{u} + U \nabla_{\dot{\sigma}}^{\pi^* \mathcal{E}} \hat{u} + \Phi(U \hat{u}) = 0, \quad \text{with} \quad \lim_{t \to -\infty} \left(U \circ \sigma\right) u(t) = e$$

where  $\nabla_{\dot{\gamma}}^{\text{En}}(U \circ \sigma) \cong \nabla_X^{\pi^* \text{En}} U$  follows from a local coordinate calculation using (4.9). Similarly

$$\begin{split} \widetilde{\nabla}_{\dot{\gamma}}^{\mathcal{E}} \Big( \Big( \widetilde{U} \circ \sigma \Big) u \Big) &+ \widetilde{\Phi} \Big( \Big( \widetilde{U} \circ \sigma \Big) u \Big) = \nabla_{\dot{\gamma}}^{\mathcal{E}} \Big( \Big( \widetilde{U} \circ \sigma \Big) u \Big) + (A \circ \sigma) \Big( \widetilde{U} \circ \sigma \Big) u + \widetilde{\Phi} \Big( \Big( \widetilde{U} \circ \sigma \Big) u \Big) \\ &= \Big[ \nabla_{\dot{\gamma}}^{\mathrm{En}} \Big( \widetilde{U} \circ \sigma \Big) \Big] u + \Big( \widetilde{U} \circ \sigma \Big) \nabla_{\dot{\gamma}}^{\mathcal{E}} u + (A \circ \sigma) \Big( \widetilde{U} \circ \sigma \Big) u + \widetilde{\Phi} \Big( \Big( \widetilde{U} \circ \sigma \Big) u \Big) \\ &\cong \Big( \nabla_{X}^{\pi^{*} \mathrm{En}} \widetilde{U} \Big) \hat{u} + \widetilde{U} \nabla_{\sigma}^{\pi^{*} \mathcal{E}} \hat{u} + A \Big( \widetilde{U} \hat{u} \Big) + \widetilde{\Phi} (U \hat{u}) = 0, \quad \text{with} \quad \lim_{t \to -\infty} \Big( \widetilde{U} \circ \sigma \Big) u(t) = e. \end{split}$$

Hence our claim above is indeed true. By our assumption the data (2.4) is the same for  $(\nabla^{\mathcal{E}}, \Phi)$  and  $(\widetilde{\nabla}^{\mathcal{E}}, \widetilde{\Phi})$  and so

$$\lim_{t \to \infty} (U \circ \sigma) u(t) = \lim_{t \to \infty} \left( \widetilde{U} \circ \sigma \right) u(t).$$

Since parallel transport such as (5.81) above is an isomorphism between fibers, varying the "e" in (5.81) implies here that  $U = \widetilde{U}$  on  $\partial^b S^* \overline{M}$ .

Intuitively speaking, we've demonstrated that knowing the parallel transport (5.81), the endomorphism fields U and  $\widetilde{U}$  encode the transform that takes all possible  $(\gamma, e)$  to the data (2.4) of  $\nabla^{\mathcal{E}}$ ,  $\Phi$  and  $\widetilde{\nabla}^{\mathcal{E}}$ ,  $\widetilde{\Phi}$  respectively. Furthermore, the assumption that the two transforms are equal gives us that U and  $\widetilde{U}$  are equal on the boundary. Hence we've reformulated our task to showing that  $U = \widetilde{U}$  on  $\partial^b S^* M$  implies the gauge equivalence stated in the theorem.

Guided by the observation (2.12) in the introduction, we next study the behavior of  $UU^{-1}$ over the interior SM. We note that both U and  $\tilde{U}$  are invertible because (5.80) can be seen as ordinary differential equations along integral curves of X (c.f. (1.8) in Chapter 3 of [5]). A quick computation using (5.79) and (4.9) shows that  $\mathbb{X}(W_1W_2) = \mathbb{X}(W_1)W_2 + W_1\mathbb{X}(W_2)$ and that  $\mathbb{X}(\mathrm{id}) = 0$ . Hence by (5.80),

$$\mathbb{X}\left(\widetilde{U}^{-1}\right) = -\widetilde{U}^{-1}\mathbb{X}\left(\widetilde{U}\right)\widetilde{U}^{-1} = \widetilde{U}^{-1}\left(A\widetilde{U} + \widetilde{\Phi}\widetilde{U}\right)\widetilde{U}^{-1} = \widetilde{U}^{-1}A + \widetilde{U}^{-1}\widetilde{\Phi}.$$

Next, we have that

$$\mathbb{X}\left(U\widetilde{U}^{-1}\right) = (-\Phi U)\widetilde{U}^{-1} + U\left(\widetilde{U}^{-1}A + \widetilde{U}^{-1}\widetilde{\Phi}\right),$$

and so we finally arrive at that  $Q = U\widetilde{U}^{-1}$  satisfies

$$\mathbb{X}Q + \Phi Q - QA - Q\overline{\Phi} = 0. \tag{5.82}$$

To apply our finite degree theorems from Section 5.3 above, we need our solution to vanish at "infinity." Hence we instead consider W = Q – id which satisfies

$$XW + \Phi W - WA - W\widetilde{\Phi} = -\Phi + A + \widetilde{\Phi}, \qquad (5.83)$$

and vanishes on the boundary  $\partial^b S^* \overline{M}$ . As before,  $\mathbb{X}W - WA$  can be thought of as  $(\pi^* \nabla)_X W$  for some connection " $\nabla$ " whose connection symbols are in  $\rho^{N_0} C^{\infty}(\overline{M})$  in any boundary

coordinates and frame (in the sense of Definition 2.6), and  $\Phi W - W \widetilde{\Phi}$  can be thought of an element of  $\rho^{N_0+1}C^{\infty}(\overline{M}; \operatorname{End} \operatorname{End} \mathcal{E})$  applied to W. Hence assuming that  $N_0$  is large enough, by Proposition 5.54 we have that W is in  $\mathcal{R}^3(SM; \pi^* \operatorname{End} \mathcal{E})$ . Furthermore, by Remark 5.5 we have that W extends continuously to  ${}^0S^*\overline{M}$  and vanishes on the boundary  $\partial^0S^*\overline{M}$ .

Now, if we look at the equation (5.83) in coordinates and frame, above any fixed point  $x \in M$  the entries of the right-hand side of (5.83) are restrictions of homogeneous polynomials of order zero and one in the variable v. By the theory of spherical harmonics these are elements of (Fourier) degree zero and one respectively (see Section 2.H in [12]). Hence by Proposition 5.52 we have that W and hence Q are of degree zero. Thus, we get that  $Q \in C^{\infty}(M; \operatorname{End} \mathcal{E}) \cap C^{0}(\overline{M}; \operatorname{End} \mathcal{E})$  (i.e. up to identification) such that  $Q|_{\partial \overline{M}} = \operatorname{id}$  and  $(Q - \operatorname{id}) \in \mathcal{R}^{3}(SM; \pi^{*}\mathcal{E})$ .

As the final step, let's show that this Q is the gauge that we wanted. Adopting the coordinates and frames in (5.79), we have that

$$\mathbb{X}Q = v(Q) + ({}^{\mathcal{E}}\Gamma)Q - Q({}^{\mathcal{E}}\Gamma).$$

Plugging this into (5.82) and equating the zeroth and first Fourier modes gives

$$\Phi Q - Q \widetilde{\Phi} = 0$$
 and  $\mathbb{X}Q - QA = 0.$ 

The first equation gives  $\widetilde{\Phi} = Q^{-1} \Phi Q$  over M and hence over  $\overline{M}$  by continuity. The second equation gives

$$A = Q^{-1} \mathbb{X} Q = Q^{-1} [v(Q) + ({}^{\mathcal{E}} \Gamma)Q - Q({}^{\mathcal{E}} \Gamma)],$$
  
$$\implies A + ({}^{\mathcal{E}} \Gamma) = Q^{-1} v(Q) + Q^{-1} ({}^{\mathcal{E}} \Gamma)Q,$$

Now, take any section  $u \in C^{\infty}(M; \mathcal{E}|_M)$  and write it as a column vector with respect to the basis  $(b_i)$ . We have that (here v(u) denotes applying v to every entry of u)

$$\begin{split} \widetilde{\nabla}_{v}^{\mathcal{E}} u &= v(u) + \left[A + \left({}^{\mathcal{E}} \Gamma\right)\right] u & \text{eq. for } \widetilde{\nabla}_{v}^{\mathcal{E}}, \\ &= v\left(Q^{-1}Qu\right) + Q^{-1}v(Q)u + Q^{-1}\left({}^{\mathcal{E}} \Gamma\right)Qu & QQ^{-1} = \text{id and plug in eq. above,} \\ &= v\left(Q^{-1}Qu\right) - v\left(Q^{-1}\right)Qu + Q^{-1}\left({}^{\mathcal{E}} \Gamma\right)Qu & Q^{-1}v(Q) = -v\left(Q^{-1}\right)Q \text{ (prod. rule)}, \\ &= Q^{-1}v(Qu) + Q^{-1}\left({}^{\mathcal{E}} \Gamma\right)Qu & \text{prod. rule,} \\ &= Q^{-1}\nabla_{v}^{\mathcal{E}}(Qu) & \text{eq. for } \nabla_{v}^{\mathcal{E}}. \end{split}$$

Again this relation extends to  $\partial M$  by continuity and hence the theorem is proved.

## 6 Proof of Theorem 2.14

Suppose that u is a twisted CKT of order  $m \ge 1$  that is in  $\mathcal{R}^3(SM; \pi^*\mathcal{E})$ . We will show that it is identically zero everywhere. We already proved the existence of a  $\kappa$  as in the statement of the lemma in the proof of Lemma 5.41 above. Recalling that  $\mathbb{X}_+ u = 0$  by definition, the idea here is to use to use our curvature bounds and Proposition 5.40 to conclude that  $\mathbb{X}_{-}u = 0$  as well. To do this, observe that plugging (5.44) and (5.45) into the equation in Proposition 5.40 gives

$$(2m+n)\|\mathbb{X}_{+}u\|_{L^{2}}^{2} \geq \left\|\nabla^{\pi^{*}\mathcal{E}}u\right\|_{L^{2}}^{2} + (2m+n-2)\|\mathbb{X}_{-}u\|_{L^{2}}^{2} + (\kappa\lambda_{m} - \|F^{\mathcal{E}}\|_{L^{\infty}}\lambda_{m}^{1/2})\|u\|_{L^{2}}^{2}.$$

Recalling the definition of  $\lambda_m$ , our assumption  $\|F^{\mathcal{E}}\|_{L^{\infty}} \leq \kappa \sqrt{n}$  implies that the coefficient of  $\|u\|_{L^2}^2$  on the right-hand side is nonnegative. Hence both sides of the inequality are nonnegative, and thus the left-hand side being zero implies that  $\mathbb{X}_{-}u = 0$ . Hence  $\mathbb{X}u = \mathbb{X}_{+}u + \mathbb{X}_{-}u = 0$  as well.

Now, consider any g-geodesic  $\gamma : (-\infty, \infty) \to M$  and its lift  $\sigma : (-\infty, \infty) \to SM$  to SM (i.e. latter is an integral curve of X). As we explained in Section 2.3 above, since (M, g) is nontrapping the limit  $\lim_{t\to-\infty} \gamma(t) \in \partial \overline{M}$  exists. We have that u vanishes on the compact boundary  $\partial^0 S^* \overline{M}$  by Remark 5.5 and hence it follows that that  $u \circ \sigma(t) \to 0$  as  $t \to -\infty$ . This combined with the fact that  $Xu \equiv 0$  imply that  $u \circ \sigma(t) \equiv 0$ . Thus indeed u vanishes everywhere.

We point out that the conclusion of the theorem follows even faster if one assumes  $\|F^{\mathcal{E}}\|_{L^{\infty}} < \kappa \sqrt{n}$  since then the coefficient of  $\|u\|_{L^2}^2$  in the above inequality is positive, and hence  $\|u\|_{L^2}^2 = 0$  giving  $u \equiv 0$ .

## 7 Injectivity over Higgs Fields

In this section we prove Corollary 2.10. Recall that  $d = \operatorname{rank} \mathcal{E}$ . Since the curvature of  $\nabla^{\mathcal{E}}$  is zero, by Theorem 2.14 we have that there are no nontrivial twisted CKTs in  $\mathcal{R}^3(SM; \pi^*\mathcal{E})$ . Hence, by Theorem 2.7, if  $N_0 \geq 0$  is big enough there exists a  $Q \in C^0(\overline{M}; \operatorname{End} \mathcal{E}) \cap C^\infty(M; \operatorname{End} \mathcal{E})$  such that  $Q|_{\partial \overline{M}} = \operatorname{id}$  and satisfies (2.8) with  $\widetilde{\nabla}^{\mathcal{E}} = \nabla^{\mathcal{E}}$ . Hence, we'll be done if we show that  $Q \equiv \operatorname{id}$  everywhere. Take coordinates  $(x^i)$  of M, a frame  $(b_i)$  for  $\mathcal{E}$  over these coordinates' domain, and consider the coordinates  $v^{i\partial}/\partial x^i \mapsto (x^i, v^i)$ of TM. Let  ${}^{\mathcal{E}}\Gamma^k_{ij}$  denote the connection symbols of  $\nabla^{\mathcal{E}}$  with respect to  $(\partial/\partial x^i)$  and  $(b_k)$ . Since the curvature of  $\nabla^{\mathcal{E}}$  is zero, by Proposition 1.2 of Appendix C in Volume II of [49], we may suppose that the  $(b_i)$  were chosen so that all of the  ${}^{\mathcal{E}}\Gamma^k_{ij} \equiv 0$ . Let us represent Q as a  $d \times d$ matrix in the basis  $(b_i)$ . Similarly we represent any section  $u \in C^\infty(\overline{M}; \mathcal{E})$  as a  $d \times 1$  column vector in this basis. Then for any section  $u = u^j b_j \cong [u^1, \ldots, u^d]$  whose component functions  $u^j$  are constant, we have that for any  $v \in TM$  in our coordinates

$$\label{eq:started} \begin{split} \nabla^{\mathcal{E}}_{v} u &= 0,\\ \widetilde{\nabla}^{\mathcal{E}}_{v} u &= Q^{-1} \nabla^{\mathcal{E}}_{v}(Q u) = Q^{-1} v(Q) u. \end{split}$$

where v(Q) denotes applying v to the entries of Q. Since  $\widetilde{\nabla}_v^{\mathcal{E}} u = \nabla_v^{\mathcal{E}} u$  by assumption and the above is true for all such u, we get that  $v(Q) \equiv 0$ . Hence Q is locally constant, and thus indeed equal to "id" everywhere.

## References

- [1] Yu. E. Anikonov and V. G. Romanov. On uniqueness of determination of a form of first degree by its integrals along geodesics. J. Inverse Ill-Posed Probl., 5(6):487-490, 1997.
- [2] Carlos A. Berenstein and Enrico Casadio Tarabusi. Inversion formulas for the kdimensional Radon transform in real hyperbolic spaces. *Duke Math. J.*, 62(3):613-631, 1991.
- [3] Jan Bohr. Stability of the non-abelian X-ray transform in dimension  $\geq 3$ . J. Geom. Anal., 31(11):11226-11269, 2021.
- [4] Jan Bohr and Gabriel P. Paternain. The transport Oka-Grauert principle for simple surfaces. J. Ec. polytech. Math., 10:727-769, 2023.
- [5] Earl A. Coddington and Norman Levinson. Theory of ordinary differential equations. McGraw-Hill Book Co., Inc., New York-Toronto-London, 1955.
- [6] Bartlomiej Czech, Lampros Lamprou, Samuel McCandlish, and James Sully. Integral geometry and holography. J. High Energy Phys., (10):175, front matter+40, 2015.
- [7] N. S. Dairbekov and V. A. Sharafutdinov. On conformal Killing symmetric tensor fields on Riemannian manifolds. *Mat. Tr.*, 13(1):85-145, 2010.
- [8] Naeem M. Desai, William R. B. Lionheart, Morten Sales, Markus Strobl, and Søren Schmidt. Polarimetric neutron tomography of magnetic fields: uniqueness of solution and reconstruction. *Inverse Problems*, 36(4):045001, 17, 2020.
- [9] Nikolas Eptaminitakis. Stability estimates for the X-ray transform on simple asymptotically hyperbolic manifolds. *Pure Appl. Anal.*, 4(3):487-516, 2022.
- [10] Nikolas Eptaminitakis and C. Robin Graham. Local X-ray transform on asymptotically hyperbolic manifolds via projective compactification. *New Zealand J. Math.*, 52:733-763, 2021 [2021-2022].
- [11] David Finch and Gunther Uhlmann. The X-ray transform for a non-abelian connection in two dimensions. volume 17, pages 695-701. 2001. Special issue to celebrate Pierre Sabatier's 65th birthday (Montpellier, 2000).
- [12] Gerald B. Folland. Introduction to partial differential equations. Princeton University Press, Princeton, NJ, second edition, 1995.
- [13] Gerald B. Folland. *Real analysis.* Pure and Applied Mathematics (New York). John Wiley & Sons, Inc., New York, second edition, 1999. Modern techniques and their applications, A Wiley-Interscience Publication.
- [14] C. Robin Graham, Colin Guillarmou, Plamen Stefanov, and Gunther Uhlmann. X-ray transform and boundary rigidity for asymptotically hyperbolic manifolds. Ann. Inst. Fourier (Grenoble), 69(7):2857-2919, 2019.

- [15] Colin Guillarmou. Lens rigidity for manifolds with hyperbolic trapped sets. J. Amer. Math. Soc., 30(2):561-599, 2017.
- [16] Colin Guillarmou, Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. The X-ray transform for connections in negative curvature. *Comm. Math. Phys.*, 343(1):83-127, 2016.
- [17] Victor Guillemin and David Kazhdan. Some inverse spectral results for negatively curved n-manifolds. In Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), volume XXXVI of Proc. Sympos. Pure Math., pages 153-180. Amer. Math. Soc., Providence, RI, 1980.
- [18] Sigurdur Helgason. The totally-geodesic Radon transform on constant curvature spaces. In Integral geometry and tomography (Arcata, CA, 1989), volume 113 of Contemp. Math., pages 141-149. Amer. Math. Soc., Providence, RI, 1990.
- [19] Sigurdur Helgason. Geometric analysis on symmetric spaces, volume 39 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1994.
- [20] A. Hilger, I. Manke, N. Kardjilov, M. Osenberg, H. Markötter, and J. Banhart. Tensorial neutron tomography of three-dimensional magnetic vector fields in bulk materials. *Nat Commun*, 9(4023), 2018.
- [21] Sean Holman and Gunther Uhlmann. On the microlocal analysis of the geodesic X-ray transform with conjugate points. J. Differential Geom., 108(3):459-494, 2018.
- [22] Joonas Ilmavirta. Coherent quantum tomography. SIAM J. Math. Anal., 48(5):3039-3064, 2016.
- [23] John M. Lee. Introduction to smooth manifolds, volume 218 of Graduate Texts in Mathematics. Springer, New York, second edition, 2013.
- [24] John M. Lee. Introduction to Riemannian manifolds, volume 176 of Graduate Texts in Mathematics. Springer, Cham, second edition, 2018.
- [25] Jere Lehtonen. The geodesic ray transform on two-dimensional CartanHadamard manifolds. PhD thesis, University of Jyväskylä, 2016.
- [26] Jere Lehtonen, Jesse Railo, and Mikko Salo. Tensor tomography on Cartan-Hadamard manifolds. *Inverse Problems*, 34(4):044004, 27, 2018.
- [27] Juan Maldacena. The large N limit of superconformal field theories and supergravity [MR1633016 (99e:81204a)]. In Trends in theoretical physics, II (Buenos Aires, 1998), volume 484 of AIP Conf. Proc., pages 51-63. Amer. Inst. Phys., Woodbury, NY, 1999.
- [28] S. V. Manakov and V. E. Zakharov. Three-dimensional model of relativistic-invariant field theory, integrable by the inverse scattering transform. *Lett. Math. Phys.*, 5(3):247-253, 1981.

- [29] Rafe Mazzeo. Hodge Cohomology of Negatively Curved Manifolds. PhD thesis, Massachusetts Institute of Technology, 1982.
- [30] Richard B. Melrose. The Atiyah-Patodi-Singer index theorem, volume 4 of Research Notes in Mathematics. A K Peters, Ltd., Wellesley, MA, 1993.
- [31] Francois Monard, Richard Nickl, and Gabriel P. Paternain. Efficient nonparametric Bayesian inference for X-ray transforms. *The Annals of Statistics*, 47(2):1113 - 1147, 2019.
- [32] Francois Monard, Plamen Stefanov, and Gunther Uhlmann. The geodesic ray transform on Riemannian surfaces with conjugate points. *Comm. Math. Phys.*, 337(3):1491-1513, 2015.
- [33] R. G. Muhometov. The problem of recovery of a two-dimensional Riemannian metric and integral geometry. Doklady Akademii Nauk, 232(1):32-35, 1977.
- [34] R. G. Novikov. Non-Abelian Radon transform and its applications. In *The Radon trans-form—the first 100 years and beyond*, volume 22 of *Radon Ser. Comput. Appl. Math.*, pages 115-127. Walter de Gruyter, Berlin, Copyright 2019.
- [35] R. G. Novikov. On determination of a gauge field on  $\mathbb{R}^d$  from its non-abelian Radon transform along oriented straight lines. J. Inst. Math. Jussieu, 1(4):559-629, 2002.
- [36] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. The attenuated ray transform for connections and Higgs fields. *Geom. Funct. Anal.*, 22(5):1460-1489, 2012.
- [37] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. Tensor tomography on surfaces. *Invent. Math.*, 193(1):229-247, 2013.
- [38] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. Tensor tomography: progress and challenges. *Chinese Ann. Math. Ser. B*, 35(3):399-428, 2014.
- [39] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. Invariant distributions, Beurling transforms and tensor tomography in higher dimensions. *Math. Ann.*, 363(1-2):305-362, 2015.
- [40] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. Geometric inverse problems with emphasis on two dimensions, volume 204 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2023. With a foreword by András Vasy.
- [41] L. N. Pestov and V. A. Sharafutdinov. Integral geometry of tensor fields on a manifold of negative curvature. *Sibirsk. Mat. Zh.*, 29(3):114-130, 221, 1988.
- [42] V. A. Sharafutdinov. Integral geometry of tensor fields. Inverse and Ill-posed Problems Series. VSP, Utrecht, 1994.
- [43] V. A. Sharafutdinov. On the inverse problem of determining a connection on a vector bundle. J. Inverse Ill-Posed Probl., 8(1):51-88, 2000.

- [44] V. A. Sharafutdinov. Variations of Dirichlet-to-Neumann map and deformation boundary rigidity of simple 2-manifolds. J. Geom. Anal., 17(1):147-187, 2007.
- [45] Plamen Stefanov and Gunther Uhlmann. Boundary rigidity and stability for generic simple metrics. J. Amer. Math. Soc., 18(4):975-1003, 2005.
- [46] Plamen Stefanov and Gunther Uhlmann. Local lens rigidity with incomplete data for a class of non-simple Riemannian manifolds. J. Differential Geom., 82(2):383-409, 2009.
- [47] Plamen Stefanov and Gunther Uhlmann. The geodesic X-ray transform with fold caustics. Anal. PDE, 5(2):219-260, 2012.
- [48] Plamen Stefanov, Gunther Uhlmann, and András Vasy. Inverting the local geodesic X-ray transform on tensors. J. Anal. Math., 136(1):151-208, 2018.
- [49] Michael E. Taylor. Partial differential equations, volume 115 of Applied Mathematical Sciences. Springer, New York, second edition, 2011.
- [50] Gunther Uhlmann and András Vasy. The inverse problem for the local geodesic ray transform. *Invent. Math.*, 205(1):83-120, 2016.
- [51] R. S. Ward. Soliton solutions in an integrable chiral model in 2 + 1 dimensions. J. Math. Phys., 29(2):386-389, 1988.