



Rational, Log Canonical, Du Bois Singularities: On the Conjectures of Kollár and Steenbrink^{*}

SÁNDOR J. KOVÁCS

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, U.S.A.
e-mail: kovacs@math.uchicago.edu skovacs@member.ams.org
Current address: Department of Mathematics, University of Chicago, Chicago, IL 60637, U.S.A.

(Received: 12 September 1997; accepted in final form: 10 March 1998)

Abstract. Let X be a proper complex variety with Du Bois singularities. Then $H^i(X, \mathbb{C}) \rightarrow H^i(X, \mathcal{O}_X)$ is surjective for all i . This property makes this class of singularities behave well with regard to Kodaira type vanishing theorems. Steenbrink conjectured that rational singularities are Du Bois and Kollár conjectured that log canonical singularities are Du Bois. Kollár also conjectured that under some reasonable extra conditions Du Bois singularities are log canonical. In this article Steenbrink's conjecture is proved in its full generality, Kollár's first conjecture is proved under some extra conditions and Kollár's second conjecture is proved under a set of reasonable conditions, and shown that these conditions cannot be relaxed.

Mathematics Subject Classifications (1991): Primary: 14F17; Secondary: 14B05.

Key words: singularities, rational, log canonical, Du Bois.

0. Introduction

Following P. Deligne's work on mixed Hodge structures, it was shown by Ph. Du Bois that for every complex scheme X one can define a filtered complex $\underline{\Omega}_X$ that 'resolves' \mathbb{C}_X , its associated graded quotients, $\mathrm{Gr}_{\mathrm{filt}}^p \underline{\Omega}_X$, have \mathcal{O}_X -linear differentials and coherent cohomology sheaves and the analogue of the Hodge–de Rham spectral sequence exists and degenerates at the E_1 term [DB81].

Du Bois singularities were introduced by J. H. M. Steenbrink. They have the property that the zeroth graded piece of Du Bois' complex, $\mathrm{Gr}_{\mathrm{filt}}^0 \underline{\Omega}_X$, 'resolves' \mathcal{O}_X . In particular, let (X, x) be a normal isolated singularity, $f: Y \rightarrow X$ a resolution of singularities such that $E = f^{-1}(0)$ is a divisor with only simple normal crossings and $Y \setminus E \simeq X \setminus \{x\}$. Then (X, x) is Du Bois if and only if $R^i f_* \mathcal{O}_Y \simeq R^i f_* \mathcal{O}_E$ for all $i > 0$.

Cohomological properties of Du Bois singularities are often similar to those of smooth points. For instance, every proper, flat degeneration f over the unit disk Δ is cohomologically insignificant provided $f^{-1}(0)$ has Du Bois singularities [S81].

^{*} Supported in part by NSF Grant DMS-9600089.

As a simple consequence of the degeneration of the Hodge–de Rham spectral sequence, one easily sees that if X is proper with Du Bois singularities, then $H^i(X, \mathbb{C}) \rightarrow H^i(X, \mathcal{O}_X)$ is surjective for all i . According to J. Kollár’s Principle [K87, 23; CKM88, 8.1; K95, 9.12] this surjectivity property makes these singularities the natural class for Kodaira type vanishing theorems [K95, 9.12, 12.3, 12.10].

Therefore it is interesting to know what kind of singularities have the Du Bois property. After classifying normal Gorenstein two-dimensional Du Bois singularities Steenbrink made the following conjecture:

CONJECTURE S. *Rational singularities are Du Bois.*

At the same time he proved the conjecture for isolated singularities (cf. [S83]).

Conjecture S was recently confirmed for projective varieties by Kollár [K95, Ch. 12]. This would have completely solved the problem if one were able to prove that a variety with rational singularities can be embedded into a projective variety with only rational singularities. Unfortunately, that is still unknown. The first result of this article is that in fact Steenbrink’s conjecture is true in general.

THEOREM S. *Conjecture S holds.*

The proof is somewhat similar to the one in [ibid.]. There are two main additional ideas: First, one observes that the degeneration of the Hodge–de Rham spectral sequence implies another surjectivity (in fact a weaker one) than the one used by Kollár. The second idea is that this surjectivity allows one to pass to local cohomology, where the other necessary ingredient of the proof holds. For details, see the argument in Section 2.

Kollár has also made a conjecture regarding Du Bois singularities [K92,1.13]:

CONJECTURE K. *Log canonical singularities are Du Bois.*

This has only been confirmed for isolated singularities. S. Ishii showed that a normal isolated Gorenstein singularity is Du Bois if and only if it is log canonical [I85] (see definition at the end of the Introduction). More generally it was shown by K. Watanabe and Ishii that an isolated log canonical singularity is Du Bois [W86], [I86b]. The second result of this paper is another step toward Kollár’s conjecture.

THEOREM K. *Let X be a complex variety with log canonical Cohen–Macaulay singularities. Let $\Sigma = \text{Sing } X$ be the set of singular points of X and let Σ_r denote the smallest closed subset of X such that $X \setminus \Sigma_r$ has rational singularities. Assume that either $\dim \Sigma + \dim \Sigma_r + 1 < \dim X$ or Σ has Du Bois singularities. Then X has Du Bois singularities.*

Remark. Unfortunately, this still does not solve even the three-dimensional case completely.* However, it gives some new results already in that case: It implies that a log canonical Cohen–Macaulay threefold with only finitely many nonrational points has Du Bois singularities. Similarly a log canonical Cohen–Macaulay threefold whose singular set is a curve with only ordinary singularities is Du Bois. In higher dimensions there are many more new cases: for instance a d -dimensional log canonical Cohen–Macaulay variety whose singular set is a curve is Du Bois as soon as $d \geq 4$, a log canonical Cohen–Macaulay variety whose singular set is smooth is always Du Bois.

On the other hand, one would also like to know how far Du Bois singularities are from being log canonical. Simple examples show that there are Du Bois singularities whose canonical divisor is not \mathbb{Q} -Cartier and there are Du Bois singularities whose canonical divisor is \mathbb{Q} -Cartier, but the singularity fails to be log canonical cf. [W86, 4.13; I86b, 2.5; I86a, 3.3]. One fact to keep in mind is that rational singularities are not necessarily log canonical, in particular there are two-dimensional rational singularities (and therefore Du Bois) that are not log canonical, but have a \mathbb{Q} -Cartier canonical divisor.

In light of these facts the following seems to be the best one can hope for in this direction.

THEOREM K'. *Let U be a normal variety and assume that K_U is Cartier and U has Du Bois singularities. Then U is log canonical.*

This was also conjectured by Kollár.

DEFINITIONS AND NOTATION. Throughout the article the groundfield will always be \mathbb{C} , the field of complex numbers. A *complex scheme* (resp. *complex variety*) will mean a separated scheme (resp. variety) of finite type over \mathbb{C} .

A divisor D is called \mathbb{Q} -Cartier if mD is Cartier for some $m > 0$. A normal variety X is said to have *log canonical* (resp. *log terminal*, *canonical*) singularities if K_X is \mathbb{Q} -Cartier and for any resolution of singularities $f: Y \rightarrow X$, with the collection of exceptional prime divisors $\{E_i\}$, there exist $a_i \in \mathbb{Q}$, $a_i \geq -1$ (resp. $a_i > -1$, $a_i \geq 0$) such that $K_Y \equiv f^*K_X + \sum a_i E_i$ (cf. [CKM88]). The *index* of a normal variety X with K_X \mathbb{Q} -Cartier is the smallest positive integer m such that mK_X is Cartier. Note that for a normal variety X with K_X \mathbb{Q} -Cartier, there exists locally an *index 1 cover*, i.e., a finite surjective morphism $\sigma: X' \rightarrow X$ such that X' has index 1 [R87, 3.6].

A singularity is called *Gorenstein* (resp. *Cohen–Macaulay*) if its local ring is a Gorenstein (resp. Cohen–Macaulay) ring. A variety is *Gorenstein* (resp. *Cohen–Macaulay*) if it admits only Gorenstein (resp. Cohen–Macaulay) singularities. Let

* Recently, I found that the results in this article give basis to a rather easy proof of the three-dimensional Cohen–Macaulay case. This is included in the next article.

X be a normal variety and $f: Y \rightarrow X$ a resolution of singularities. X is said to have *rational singularities* if $R^i f_* \mathcal{O}_Y = 0$ for all $i > 0$.

Let X be a complex scheme of dimension n . $D_{\text{filt}}(X)$ denotes the derived category of filtered complexes of \mathcal{O}_X -modules with differentials of order ≤ 1 and $D_{\text{filt, coh}}(X)$ the subcategory of $D_{\text{filt}}(X)$ of complexes K^\cdot , such that for all i , the cohomology sheaves of $\text{Gr}_{\text{filt}}^i K^\cdot$ are coherent (cf. DB81, [GNPP88]). $D(X)$ and $D_{\text{coh}}(X)$ denotes the derived categories with the same definition except that the complexes are assumed to have the trivial filtration. The superscripts $+$, $-$, b carry the usual meaning (bounded below, bounded above, bounded). $C(X)$ is the category of complexes of \mathcal{O}_X -modules with differentials of order ≤ 1 and for $u \in \text{Mor}(C(X))$, $M(u) \in \text{Ob}(C(X))$ denotes the mapping cone of u (cf. [H66]). The isomorphism in these categories is denoted by \simeq_{qis} . If K^\cdot is a complex in any of the above defined categories, then $h^i(K^\cdot)$ denotes the i th cohomology sheaf of K^\cdot . In particular every sheaf is naturally a complex with $h^i = 0$ for $i \neq 0$.

The right derived functor of an additive functor F , if exists, is denoted by RF and $R^i F$ stands for $h^i \circ RF$. In particular, \mathbb{H}^i denotes $R^i \Gamma$ and \mathbb{H}_Z^i denotes $R^i \Gamma_Z$ where Γ is the functor of global sections and Γ_Z is the functor of global sections with support in the closed subset Z . Finally $\omega_X = f^! \mathbb{C}$ is the dualizing complex of X where $f: X \rightarrow \text{Spec } \mathbb{C}$ is the natural morphism (cf. [H66]). Note that if X has Gorenstein singularities, then $\omega_X \simeq_{\text{qis}} \omega_X[n]$ and ω_X is a line bundle.

The dimension of the empty set is $-\infty$.

1. Du Bois Singularities

The actual construction of Du Bois' complex will not be used in this article. Therefore it is not repeated here. Instead the interested reader is referred to the original article. Note also that a simplified construction was later obtained by [GNPP88] via the general theory of cubic resolutions. An easily accessible introduction can be found in [S85].

The basic results regarding $\underline{\Omega}_X$ that are essential in the sequel are summarized in the following theorem.

THEOREM 1.1 [DB81, 3.2, 3.10, 4.5, 4.11], [GNPP88, III.1.12, III.1.17, V.3.5].
For every complex scheme X there exists an $\underline{\Omega}_X \in \text{Ob}(D_{\text{filt}}(X))$ with the following properties.

- (1.1.1) It is functorial, i.e., if $\phi: Y \rightarrow X$ is a morphism of complex schemes, then there exists a natural map ϕ^* of filtered complexes $\phi^*: \underline{\Omega}_X \rightarrow R\phi_* \underline{\Omega}_Y$. Furthermore, $\underline{\Omega}_X \in \text{Ob}(D_{\text{filt, coh}}^b(X))$ and if ϕ is proper, then ϕ^* is a morphism in $D_{\text{filt, coh}}^b(X)$.
- (1.1.2) Let Ω_X be the usual De Rham complex of Kähler differentials considered with the 'filtration bête'. Then there exists a natural map of filtered complexes $\Omega_X \rightarrow \underline{\Omega}_X$ and if X is smooth, it is a quasi-isomorphism.

(1.1.3) Let $U \subseteq X$ be an open subscheme of X . Then $\underline{\Omega}_X|_U \simeq_{\text{qis}} \underline{\Omega}_U$.

(1.1.4) If X is proper, then there exists a spectral sequence degenerating at E_1 and abutting to the singular cohomology of X : $E_1^{pq} = \mathbb{H}^q(X, \text{Gr}_{\text{filt}}^p \underline{\Omega}_X[p]) \Rightarrow H^{p+q}(X, \mathbb{C})$.

(1.1.5) $\dim \text{Supp } h^i(\text{Gr}_{\text{filt}}^p \underline{\Omega}_X) \leq \dim X - i + p$.

(1.1.6) Let $\underline{\Omega}_X^0 = \text{Gr}_{\text{filt}}^0 \underline{\Omega}_X$. Let $\Sigma = \text{Sing } X$ be the singular set of X and $f: Y \rightarrow X$ a resolution of singularities such that it is an isomorphism outside Σ and $E = f^{-1}(\Sigma)$ is a divisor with normal crossings. Then there exists a distinguished triangle, $\underline{\Omega}_X^0 \rightarrow \underline{\Omega}_\Sigma^0 \oplus Rf_* \mathcal{O}_Y \rightarrow Rf_* \mathcal{O}_E \xrightarrow{+1}$, where the morphisms are those of (1.1.1) and (1.1.2).

Remark 1.1.7. Let $f: Y \rightarrow X$ be a resolution of singularities of X . Then by (1.1.1) and (1.1.2) the natural morphism $\mathcal{O}_X \rightarrow Rf_* \mathcal{O}_Y$ factors through $\underline{\Omega}_X^0$ and if X is proper, then by (1.1.4) $H^i(X, \mathbb{C}) \rightarrow \mathbb{H}^i(X, \underline{\Omega}_X^0)$ is surjective for all i .

DEFINITION 1.2 [S83]. X is said to have *Du Bois singularities* if $\mathcal{O}_X \rightarrow \underline{\Omega}_X^0$ is a quasi-isomorphism. (I.e., $h^0(\underline{\Omega}_X^0) \simeq \mathcal{O}_X$ and $h^i(\underline{\Omega}_X^0) = 0$ for all $i \neq 0$.) In particular, if X is proper and has Du Bois singularities, then $H^i(X, \mathbb{C}) \rightarrow H^i(X, \mathcal{O}_X)$ is surjective for all i .

2. Rational Singularities – The Key Lemma

2.1. Let X be a complex scheme that one would like to prove to have Du Bois singularities. Let F^\cdot be a complex such that $\mathcal{O}_X \rightarrow \underline{\Omega}_X^0 \rightarrow F^\cdot \xrightarrow{+1}$ forms a distinguished triangle and let $\Sigma_{DB} = \cup \text{Supp } h^i(F^\cdot)$ the union of the supports of the cohomology sheaves of F^\cdot . Then Σ_{DB} is the *non-Du Bois locus* of U . By taking general hyperplane sections, as in [K95, 12.8], one may assume that $\dim \Sigma_{DB} \leq 0$. Therefore as long as the assumptions on X are invariant under taking hyperplane sections, one can restrict to the case when the possibly non-Du Bois locus is at most a set of finite points.

The following is the key step in the proof of both Theorem S and Theorem K.

LEMMA 2.2. *Let U be a complex scheme with a finite set of points, P , such that $U \setminus P$ has only Du Bois singularities and assume that $H_P^i(U, \mathcal{O}_U) \rightarrow \mathbb{H}_P^i(U, \underline{\Omega}_U^0)$ is injective for all $i = 0, \dots, \dim U$. Then U has Du Bois singularities.*

Proof. Since the statement is local, one may assume that U is affine. Let F^\cdot be the complex defined in (2.1). By assumption P contains $\Sigma_{DB} = \cup \text{Supp } h^i(F^\cdot)$, the non-Du Bois locus of U .

Next let X be a projective closure of U , and let $Q = X \setminus U$ and $Z = P \cup Q$. Then $X \setminus Z \simeq U \setminus P$ has only Du Bois singularities, i.e., $\mathcal{O}_{X \setminus Z} \simeq_{\text{qis}} \underline{\Omega}_{X \setminus Z}^0$.

Now by (1.1.4) the composition $H^i(X, \mathbb{C}) \rightarrow H^i(X, \mathcal{O}_X) \rightarrow \mathbb{H}^i(X, \underline{\Omega}_X^0)$ is surjective for all i . Then in the commutative diagram

$$\begin{array}{ccccccc} H^{i-1}(X \setminus Z, \mathcal{O}_{X \setminus Z}) & \longrightarrow & H_Z^i(X, \mathcal{O}_X) & \longrightarrow & H^i(X, \mathcal{O}_X) & \longrightarrow & H^i(X \setminus Z, \mathcal{O}_{X \setminus Z}) \\ \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta \\ \mathbb{H}^{i-1}(X \setminus Z, \underline{\Omega}_{X \setminus Z}^0) & \longrightarrow & \mathbb{H}_Z^i(X, \underline{\Omega}_X^0) & \longrightarrow & \mathbb{H}^i(X, \underline{\Omega}_X^0) & \longrightarrow & \mathbb{H}^i(X \setminus Z, \underline{\Omega}_{X \setminus Z}^0) \end{array}$$

the rows are exact, α and δ are isomorphisms, and γ is surjective. Hence β is surjective by the 5-lemma.

Observe that since $\dim P \leq 0$, $P \cap Q = \emptyset$, and then

$$H_Z^i(X, \mathcal{O}_X) \simeq H_P^i(X, \mathcal{O}_X) \oplus H_Q^i(X, \mathcal{O}_X),$$

$$\mathbb{H}_Z^i(X, \underline{\Omega}_X^0) \simeq \mathbb{H}_P^i(X, \underline{\Omega}_X^0) \oplus \mathbb{H}_Q^i(X, \underline{\Omega}_X^0)$$

and by excision $H_P^i(X, \mathcal{O}_X) \simeq H_P^i(U, \mathcal{O}_U)$ and $\mathbb{H}_P^i(X, \underline{\Omega}_X^0) \simeq \mathbb{H}_P^i(U, \underline{\Omega}_U^0)$.

Therefore $H_P^i(U, \mathcal{O}_U) \rightarrow \mathbb{H}_P^i(U, \underline{\Omega}_U^0)$ is surjective. By assumption it is also injective, hence an isomorphism.

The cohomology sheaves of F^\cdot are supported on P , so $\mathbb{H}^i(U \setminus P, F^\cdot) = 0$ for all i . Hence $\mathbb{H}^i(U, F^\cdot) = \mathbb{H}_P^i(U, F^\cdot) = 0$. Using again that $\dim P \leq 0$, one finds that $\mathbb{H}^i(U, F^\cdot) = H^0(U, h^i(F^\cdot))$, so in fact $h^i(F^\cdot) = 0$ for all i , thus $\mathcal{O}_U \simeq \underline{\Omega}_U^0$. \square

THEOREM 2.3. *Let U be a complex scheme such that $\mathcal{O}_U \rightarrow \underline{\Omega}_U^0$ has a left inverse, then U has Du Bois singularities.*

Proof. The statement is local, so one may assume that U is affine. Since $\mathcal{O}_H \simeq \mathcal{O}_U \otimes_{\mathbb{L}} \mathcal{O}_H$ and $\underline{\Omega}_H^0 \simeq_{\text{qis}} \underline{\Omega}_U^0 \otimes_{\mathbb{L}} \mathcal{O}_H$ for a general hyperplane H (cf. [K95, 12.6.2]), one can use (2.1), and then (2.2) can be applied. \square

COROLLARY 2.4 (cf. [K95, 12.8]). *Let V be a complex scheme with Du Bois singularities and $f: V \rightarrow U$ a morphism to a complex scheme such that $\mathcal{O}_U \rightarrow Rf_*\mathcal{O}_V$ has a left inverse. Then U has Du Bois singularities as well.*

Proof. $\mathcal{O}_U \rightarrow \underline{\Omega}_U^0 \rightarrow Rf_*\mathcal{O}_V$ has a left inverse, so the statement follows by (2.3). \square

COROLLARY 2.5 (cf. [K95, 12.8.2]). *Let V be a complex scheme with Du Bois singularities and $f: V \rightarrow U$ a finite and dominant morphism to a normal variety. Then U also has Du Bois singularities.*

Proof. $R^i f_*\mathcal{O}_V = 0$ for $i > 0$ and the normalized trace map splits $\mathcal{O}_U \rightarrow f_*\mathcal{O}_V$. \square

COROLLARY 2.6 (cf. [S83, 3.7], [K95, 12.9]). *Let U be a complex variety with rational singularities. Then U has Du Bois singularities.*

Proof. Let $f: V \rightarrow U$ be a resolution of singularities. Then $\mathcal{O}_U \simeq_{\text{qis}} Rf_*\mathcal{O}_V$, so the statement again follows by (2.4). \square

COROLLARY 2.7. *Let U be a complex variety with log terminal singularities. Then U has Du Bois singularities.*

Proof. First assume that U has canonical singularities of index 1. Let $f: V \rightarrow U$ be a resolution of singularities. Now $f^*\omega_U$ is a subsheaf of ω_V , so by the Grauert–Riemenschneider vanishing theorem $Rf_*\omega_V \simeq_{\text{qis}} \omega_U$. Then $\omega_U \rightarrow Rf_*f^*\omega_U \rightarrow Rf_*\omega_V \simeq_{\text{qis}} \omega_U$ shows that $\omega_U \rightarrow Rf_*f^*\omega_U$ has a left inverse, so tensoring by ω_U^{-1} one obtains that $\mathcal{O}_U \rightarrow Rf_*\mathcal{O}_V$ has a left inverse. Then (2.4) implies the result.

The statement is local, so in the general case one can take the index 1 cover, U' , of U which has canonical singularities of index 1 (cf. [R87, 3.6]). Then U' has Du Bois singularities by the first part, so U has Du Bois singularities by (2.5). \square

Remark 2.7.1. The last result certainly follows also from (2.6), but then one would have to appeal to the nontrivial fact that log terminal singularities are rational (cf. [E81], [F81], [KMM87]), whereas the above proof is considerably short and simple.

3. Log Canonical Singularities

The following notation will be used through the rest of the article.

NOTATION 3.1. Let U be a complex scheme and $\Sigma = \text{Sing } U$ the set of singular points of U . Further let Σ_r denote the smallest closed subset of U , such that $U \setminus \Sigma_r$ has rational singularities. Σ and Σ_r will be considered with the reduced induced subscheme structure. Let $f: V \rightarrow U$ be a resolution of singularities such that it is an isomorphism outside Σ and $E = f^{-1}(\Sigma)$ is a divisor with normal crossings. Finally let $\mathcal{Q} = \omega_U/f_*\omega_V$. Note that $\text{Supp } \mathcal{Q} \subseteq \Sigma_r$.

Grothendieck duality will be used in the following form (cf. [H66, III.11.1]):

For $f: V \rightarrow U$ as above and for all F^\bullet bounded complexes of \mathcal{O}_V -modules $Rf_*R\mathcal{H}om_V(F^\bullet, \omega_V) \simeq R\mathcal{H}om_U(Rf_*F^\bullet, \omega_U)$.

LEMMA 3.2. *Let U be a complex variety of dimension n with log canonical Gorenstein singularities. Then the natural maps*

$$R^i f_*\mathcal{O}_V \rightarrow R^i f_*\mathcal{O}_E, \quad h^i(\underline{\Omega}_U^0) \rightarrow h^i(\underline{\Omega}_\Sigma^0)$$

are injective for all $i > 0$.

Proof. U has log canonical singularities, so $f^*\omega_U \subseteq \omega_V(E)$, and thus $\omega_U \simeq f_*\omega_V(E)$ by [KMM87, 1-3-2]. Consider the following commutative diagram

$$\begin{array}{ccc} f_*\omega_V & \longrightarrow & f_*\omega_V(E) \simeq \omega_U \\ \downarrow & & \downarrow \\ Rf_*\omega_V & \longrightarrow & Rf_*\omega_V(E). \end{array}$$

By the Grauert–Riemenschneider vanishing theorem, the first vertical arrow is a quasi-isomorphism, so the natural morphism on the bottom factors through ω_U

$$Rf_*\omega_V \rightarrow \omega_U \rightarrow Rf_*\omega_V(E).$$

Hence the same holds for the dualizing complexes

$$Rf_*\omega_V \rightarrow \omega_U \rightarrow Rf_*\omega_V(E)[n]. \quad (3.2.1)$$

Next apply $R\mathcal{H}om_U(-, \omega_U)$ to (3.2.1). By Grothendieck duality

$$Rf_*\mathcal{O}_V(-E) \simeq_{\text{qis}} R\mathcal{H}om_U(Rf_*\omega_V(E)[n], \omega_U),$$

$$\mathcal{O}_U \simeq_{\text{qis}} R\mathcal{H}om_U(\omega_U, \omega_U),$$

$$Rf_*\mathcal{O}_V \simeq_{\text{qis}} R\mathcal{H}om_U(Rf_*\omega_V, \omega_U),$$

so (3.2.1) implies that the natural morphism $Rf_*\mathcal{O}_V(-E) \rightarrow Rf_*\mathcal{O}_V$ factors through \mathcal{O}_U

$$Rf_*\mathcal{O}_V(-E) \rightarrow \mathcal{O}_U \rightarrow Rf_*\mathcal{O}_V. \quad (3.2.2)$$

Observe that the natural morphism $\mathcal{O}_U \rightarrow Rf_*\mathcal{O}_V$ factors through $\underline{\Omega}_U^0$, so (3.2.2) gives a natural morphism $Rf_*\mathcal{O}_V(-E) \rightarrow \underline{\Omega}_U^0$ that factors through \mathcal{O}_U

$$Rf_*\mathcal{O}_V(-E) \rightarrow \mathcal{O}_U \rightarrow \underline{\Omega}_U^0. \quad (3.2.3)$$

By (1.1.6) there exists a distinguished triangle $\underline{\Omega}_U^0 \rightarrow \underline{\Omega}_\Sigma^0 \oplus Rf_*\mathcal{O}_V \rightarrow Rf_*\mathcal{O}_E \xrightarrow{+1}$.

Then it is easy to see, that one has the following commutative diagram of distinguished triangles (cf. [DB90, 7.7])

$$\begin{array}{ccccccc} Rf_*\mathcal{O}_V(-E) & \longrightarrow & \underline{\Omega}_U^0 & \longrightarrow & \underline{\Omega}_\Sigma^0 & \xrightarrow{+1} & \\ \downarrow & & \downarrow & & \downarrow & & \\ Rf_*\mathcal{O}_V(-E) & \longrightarrow & Rf_*\mathcal{O}_V & \longrightarrow & Rf_*\mathcal{O}_E & \xrightarrow{+1} & . \end{array} \quad (3.2.4)$$

Now (3.2.2) and (3.2.3) implies that the natural morphisms on the higher cohomology sheaves must be zero

$$R^i f_* \mathcal{O}_V(-E) \rightarrow 0 \rightarrow R^i f_* \mathcal{O}_V, \quad R^i f_* \mathcal{O}_V(-E) \rightarrow 0 \rightarrow h^i(\underline{\Omega}_U^0).$$

Hence the statement follows by (3.2.4). \square

The following is probably known to experts, but I could not find a reference, so a proof is included here

LEMMA 3.3. *Let U be a complex Cohen–Macaulay scheme of dimension n . Then $R^i f_* \mathcal{O}_V = 0$ for $0 < i < n - \dim \Sigma_r - 1$.*

Proof. Apply $R\mathcal{H}om_U(-, \omega_U)$ to the short exact sequence $0 \rightarrow f_* \omega_V \rightarrow \omega_U \rightarrow \mathcal{Q} \rightarrow 0$.

By the Grauert–Riemenschneider vanishing theorem, $f_* \omega_V \simeq_{\text{qis}} Rf_* \omega_V$, so by Grothendieck duality $R\mathcal{H}om_U(\mathcal{Q}, \omega_U) \rightarrow \mathcal{O}_U \rightarrow Rf_* \mathcal{O}_V \xrightarrow{+1}$ forms a distinguished triangle, hence for $i > 0$, $R^i f_* \mathcal{O}_V \simeq \mathcal{E}xt_U^{i+1}(\mathcal{Q}, \omega_U)$.

Let $x \in U$ be a closed point. The local ring of x , \mathcal{O}_x , is a Cohen–Macaulay ring of dimension n , so $(R^i f_* \mathcal{O}_V)_x \simeq \mathcal{E}xt_U^{i+1}(\mathcal{Q}, \omega_U)_x \simeq \mathcal{E}xt_{\mathcal{O}_x}^{i+1}(\mathcal{Q}_x, \omega_{\mathcal{O}_x}) = 0$ for $1 < i + 1 < \dim \mathcal{O}_x - \dim \mathcal{Q}_x$ (cf. BH93, 3.5.11). Now the statement follows since $\dim \mathcal{Q}_x \leq \dim \Sigma_r$. \square

Remark 3.3.1. In this lemma f may be an arbitrary resolution of singularities (as opposed to the assumption in (3.1)).

THEOREM 3.4. *Let U be a complex variety of dimension n with log canonical Gorenstein singularities. Assume that $h^i(\underline{\Omega}_\Sigma^0) = 0$ for $i \geq n - \dim \Sigma_r - 1$. Then U has Du Bois singularities.*

Proof. Consider the following commutative diagram where the rows are distinguished triangles

$$\begin{array}{ccccccc} \mathcal{O}_U & \longrightarrow & \underline{\Omega}_U^0 & \longrightarrow & F^\cdot & \xrightarrow{+1} & \\ \downarrow \simeq & & \downarrow & & \downarrow & & \\ \mathcal{O}_U & \longrightarrow & Rf_* \mathcal{O}_V & \longrightarrow & R^\cdot & \xrightarrow{+1} & \end{array}$$

For $i > 0$, $h^i(F^\cdot) \simeq h^i(\underline{\Omega}_U^0) \subset h^i(\underline{\Omega}_\Sigma^0)$ by (3.2), so $h^i(F^\cdot) = 0$ for $i \geq n - \dim \Sigma_r - 1$.

This remains true after taking general hyperplane sections, so as in (2.1), one may assume that $\dim \Sigma_{DB} \leq 0$, i.e., for a finite set of points, P , $U \setminus P$ has Du Bois singularities.

The cohomology sheaves of F^\cdot are supported on P , so $\mathbb{H}^i(U \setminus P, F^\cdot) = 0$ for all i and since $\dim P \leq 0$, one finds that $\mathbb{H}_P^i(U, F^\cdot) = H_P^0(U, h^i(F^\cdot))$.

$H_p^0(U, h^i(F^\cdot)) = 0$ for $i \geq n - \dim \Sigma_r - 1$, so $\mathbb{H}_p^q(U, F^\cdot) = 0$ for $q \geq n - \dim \Sigma_r - 1$.

On the other hand, $h^i(R^\cdot) = 0$ for $i \leq 0$, since $\mathcal{O}_U \simeq f_*\mathcal{O}_V$, and $h^i(R^\cdot) \simeq R^i f_*\mathcal{O}_V = 0$ for $0 < i < n - \dim \Sigma_r - 1$ by (3.3). Then $H_p^j(U, h^i(R^\cdot)) = 0$ for $i < n - \dim \Sigma_r - 1$ and all j trivially, and then $\mathbb{H}_p^q(U, R^\cdot) = 0$ for $q < n - \dim \Sigma_r - 1$.

Therefore $\mathbb{H}_p^q(U, F^\cdot) \rightarrow \mathbb{H}_p^q(U, R^\cdot)$ is zero for all q . Now the following commutative diagram

$$\begin{array}{ccccc} \mathbb{H}_p^{i-1}(U, F^\cdot) & \longrightarrow & H_p^i(U, \mathcal{O}_U) & \longrightarrow & \mathbb{H}_p^i(U, \underline{\Omega}_U^0) \\ \downarrow 0 & & \downarrow \simeq & & \downarrow \\ \mathbb{H}_p^{i-1}(U, R^\cdot) & \longrightarrow & H_p^i(U, \mathcal{O}_U) & \longrightarrow & \mathbb{H}_p^i(U, Rf_*\mathcal{O}_V) \end{array}$$

implies, that $\mathbb{H}_p^{i-1}(U, F^\cdot) \rightarrow H_p^i(U, \mathcal{O}_U)$ is zero, thus $H_p^i(U, \mathcal{O}_U) \rightarrow \mathbb{H}_p^i(U, \underline{\Omega}_U^0)$ is injective. The result now follows by (2.2). \square

COROLLARY 3.5. *Let U be a complex variety with log canonical Cohen–Macaulay singularities. Assume that either $\dim \Sigma + \dim \Sigma_r + 1 < \dim U$ or Σ has Du Bois singularities. Then U has Du Bois singularities.*

Proof. Passing to the index 1 cover, as in Corollary 2.7, one may assume that U has Gorenstein singularities. By (1.1.5) $\dim \text{Supp } h^i(\underline{\Omega}_\Sigma^0) \leq \dim \Sigma - i$, so the assumptions imply that $\dim \text{Supp } h^i(\underline{\Omega}_\Sigma^0) < \dim U - \dim \Sigma_r - 1 - i$ for all $i > 0$. Then the statement follows by (3.4). \square

Finally let us regard the opposite direction, namely that Du Bois singularities are not far from being log canonical.

THEOREM 3.6. *Let U be a normal variety and assume that K_U is Cartier and U has Du Bois singularities. Then U is log canonical.*

Proof. The distinguished triangle of (1.1.6) $\underline{\Omega}_U^0 \rightarrow \underline{\Omega}_\Sigma^0 \oplus Rf_*\mathcal{O}_V \rightarrow Rf_*\mathcal{O}_E \xrightarrow{+1}$ implies that the natural morphism $Rf_*\mathcal{O}_V(-E) \rightarrow Rf_*\mathcal{O}_V$ factors through $\underline{\Omega}_U^0 \simeq_{\text{qis}} \mathcal{O}_U$. Hence there exists a morphism $Rf_*\mathcal{O}_V(-E) \rightarrow \mathcal{O}_U$ that is a quasi-isomorphism on $U \setminus \Sigma$. Applying $R\mathcal{H}om_U(-, \omega_U)$ to this morphism one obtains $\omega_U \rightarrow Rf_*\omega_V(E)[n]$ and taking the $-n$ th cohomology gives a morphism $\omega_U \rightarrow f_*\omega_V(E)$ that is an isomorphism on $U \setminus \Sigma$ (in particular it is not the zero morphism). By adjointness this gives a nonzero morphism $f^*\omega_U \rightarrow \omega_V(E)$. $f^*\omega_U$ is a line bundle, so this implies that $f^*\omega_U \subseteq \omega_V(E)$, and therefore U is log canonical. \square

References

- [BH93] Bruno, W. and Herzog, J.: *Cohen-Macaulay Rings*, Cambridge University Press, 1993.
 [CKM88] Clemens, H., Kollár, J. and Mori, S.: Higher dimensional complex geometry, *Astérisque* **166** (1988).

- [DB81] Du Bois, Ph.: Complexe de De Rham filtré d'une variété singulière, *Bull. Soc. Math. France* **109** (1981), 41–81.
- [DB90] Du Bois, Ph.: Dualité dans la catégorie des complexes filtrés d'opérateurs différentiels d'ordre ≤ 1 , *Collect. Math.* **41** (1990), 89–121.
- [E81] Elkik, R.: Rationalité des singularités canoniques, *Invent. Math.* **64** (1981), 1–6.
- [F81] Flenner, H.: Rationale quasihomogene Singularitäten, *Arch. Math.* **36** (1981), 35–44.
- [GR70] Grauert, H. and Riemenschneider, O.: Verschwindungssätze für analytische Kohomologiegruppen auf komplexen Räumen, *Invent. Math.* **11** (1970), 263–292.
- [GNPP88] Guillén, F., Navarro Aznar, V., Pascual-Gainza, P. and Puerta, F.: *Hyperrésolutions cubiques et descente cohomologique*, Lecture Notes in Math. 1335, Springer-Verlag, 1988.
- [H66] Hartshorne, R.: *Residues and Duality*, Lecture Notes in Math. 20, Springer-Verlag, 1966.
- [I85] Ishii, S.: On Isolated Gorenstein Singularities, *Math. Ann.* **270** (1985), 541–554.
- [I86a] Ishii, S.: Du Bois singularities on a normal surface, in: *Complex Analytic Singularities*, Adv. Stud. Pure Math. 8, Kinokuniya–North-Holland, 1986, 153–163.
- [I86b] Ishii, S.: Isolated \mathbb{Q} -Gorenstein singularities of dimension three, in: *Complex Analytic Singularities*, Adv. Stud. Pure Math. 8, Kinokuniya–North-Holland, 1986, 165–198.
- [KMM87] Kawamata, Y., Matsuda, K. and Matsuki, K.: Introduction to the minimal model problem, in: T. Oda (ed.), *Algebraic Geometry*, Sendai, Adv. Stud. Pure Math. 10, Kinokuniya–North-Holland, 1987, 283–360.
- [K87] Kollár, J.: Vanishing theorems for cohomology groups, *Algebraic Geometry (Bowdoin 1985)*, Proc. Sympos. Pure Math. 46, Amer. Math. Soc., Providence, 1987, pp. 233–243.
- [K95] Kollár, J.: *Shafarevich Maps and Automorphic Forms*, Princeton University Press, 1995.
- [K92] Kollár, J.: (with 14 coauthors): Flips and abundance for algebraic threefolds, *Astérisque* **211** (1992).
- [R87] Reid, M.: Young person's guide to canonical singularities, *Algebraic Geometry (Bowdoin 1985)*, Proc. Sympos. Pure Math. 46, Amer. Math. Soc., Providence, 1987, pp. 345–416.
- [S81] Steenbrink, J. H. M.: Cohomologically insignificant degenerations, *Compositio Math.* **42** (1981), 315–320.
- [S83] Steenbrink, J. H. M.: Mixed Hodge structures associated with isolated singularities, *Singularities – Part 2*, Proc. Sympos. Pure Math. 40, Amer. Math. Soc., Providence, 1983, pp. 513–536.
- [S85] Steenbrink, J. H. M.: Vanishing theorems on singular spaces, *Astérisque* **130** (1985), 330–341.
- [W86] Watanabe, K.: *On Plurigenera of Normal Isolated Singularities II*, in: *Complex Analytic Singularities*, Adv. Stud. Pure Math. 8, Kinokuniya–North-Holland, 1986, 671–685.