

Your Name

Your Signature

Student ID #

--	--	--	--	--	--	--

Quiz Section

--	--

Professor's Name

TA's Name

- Turn off and put away all electronic devices except your non-graphing calculator.
- This exam is closed book. You may use one $8\frac{1}{2} \times 11$ sheet of handwritten notes (both sides may be used).
- Graphing calculators are not allowed. Do not share notes.
- In order to receive full credit, you must show all of your work on the exam paper (**even** if you could do the work in your head!). Remember to read each problem carefully and answer the questions being asked.
- Place a box around **YOUR FINAL ANSWER** to each question.
- If you need more room, use the **back of the previous page** and indicate to the reader that you have done so.
- Raise your hand if you have a question.

Problem	Total Points	Score
1	12	
2	10	
3	10	
4	10	
5	20	

Problem	Total Points	Score
6	10	
7	8	
8	10	
9	10	
Total	100	

1. Consider the function $f(x) = \sin(x - 4) + \cos(x - 4) + 4\sqrt{x}$.

(a) [6 points] Find the second Taylor polynomial T_2 of $f(x)$ based at $b = 4$.

(b) [2 points] Use the second Taylor polynomial T_2 to approximate $f(4.1)$.

(c) [4 points] Use Taylor's inequality to find an upper bound for the error in your approximation above.

2. Consider the function $f(x) = \frac{1 - \cos(x^2)}{x^3}$ for $x \neq 0$, and $f(0) = 0$.

(a) [5 points] Find the Taylor series for the function $f(x)$ about $b = 0$.

Write your answer in summation notation.

(b) [2 points] Find the first three nonzero terms of the Taylor series.

(c) [3 points] Find the interval on which the series in (a) converges.

3. Given points $P(1, 0, 2)$, $Q(3, -1, 5)$, and $R(0, 1, 1)$.

(a) [5 points] Find the equation of the plane containing the three points P , Q , and R .

(b) [5 points] Write the vector \overrightarrow{PQ} as a sum of two vectors, one parallel to \overrightarrow{PR} , and the other perpendicular to \overrightarrow{PR} .

4. [10 points] Decide if the following statements are TRUE or FALSE. You need not explain your answer.

(a) _____ The planes $2x - 2y + z = 4$ and $x - y + z = 2$ are parallel.

(b) _____ The vectors $\langle 1, -2, 5 \rangle$ and $\langle 2, 1, 0 \rangle$ are perpendicular.

(c) _____ If $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$, then $\vec{b} = \vec{c}$.

(d) _____ The lines

$$x = 2 - 3t, \quad y = 5 + t, \quad z = 4t$$

and

$$x = -3t, \quad y = 1 + t, \quad z = 2 + 4t$$

are parallel.

(e) _____ The line

$$x = 1 - t, \quad y = t, \quad z = 4 + 7t$$

intersects the plane $x - y - 7z = 3$ at some point.

(f) _____ The line

$$x = 2 + 2t \quad y = 5 - 8t, \quad z = -4 + 3t$$

is parallel to the plane $-x - y - 2z = 3$.

(g) _____ For any three vectors $\vec{a}, \vec{b}, \vec{c}$, we have $|\vec{a} \cdot (\vec{b} \times \vec{c})| = |(\vec{a} \times \vec{b}) \cdot \vec{c}|$.

(h) _____ For any three vectors $\vec{a}, \vec{b}, \vec{c}$, we have $\vec{a} \cdot (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{b}) \times (\vec{a} \cdot \vec{c})$.


(i) _____ Two distinct lines parallel to a third line in 3D-space are parallel to each other.

(j) _____ Two distinct lines perpendicular to a third line in 3D-space are parallel to each other.


5. A curve in the xy -plane, called cardioid, is determined by the polar equation

$$r = 1 + \cos \theta.$$


- (a) [5 points] Find all values of θ at which the tangent line is vertical. You should find them using calculation, not the picture.



- (b) [5 points] Find the area of the region bounded by the x -axis and the cardioid $r = 1 + \cos \theta$ from $\theta = 0$ to $\theta = \pi$.




- (c) [10 points] Let R be the region in the first quadrant of the xy -plane that lies inside the cardioid $r = 1 + \cos \theta$ and outside the circle $r = 1$. Find the volume of the solid that lies above R and below the plane $z = y$.



6. [10 points] Find the radius of curvature of the Archimedean spiral

$$x = t \cos t, \quad y = t \sin t$$

at the point $(-\pi, 0)$. (You are finding the radius of the circle shown in the figure below.)



7. [8 points] Evaluate the integral $\int_0^2 \int_0^{4-x^2} \frac{xe^{2y}}{4-y} dy dx$.

8. [10 points] A lamina occupies the region in the xy -plane bounded by the lines $x = 1$, $x = 2$, $y = ax$, and $y = 2ax$ for some positive number a . The lamina has density function $\rho(x, y) = \frac{1}{x} + \frac{1}{y^2}$. Find the value of a that minimizes the mass of the lamina.

9. [10 points] Find three positive numbers x , y , and z whose sum is 100 and for which the product

$$xy^2z^3$$

is maximum.