

Lecture 10-28: The rational canonical and Jordan forms

October 28, 2024

Having classified finitely generated modules over a PID, I now turn to some linear algebraic applications, following section 12.2 in the text.

Let k be a field and V a finite-dimensional vector space over k . Let $T : V \rightarrow V$ be a linear transformation. One can make V into a module over the polynomial ring $k[x]$ by decreeing that k act as usual by scalar multiplication, while x acts by T , so that any polynomial in x acts by the same polynomial in T (see the Example on p. 340). Since V is finitely generated as a module (being finite-dimensional), the classification theorem proved last time applies to it. Given two such transformations T, T' on the same V , we see that the two module structures on V (one with T , the other with T') are isomorphic if and only if the matrices of T and T' with respect to any fixed basis of V are similar (or conjugate),

By the elementary divisor version of this classification we can write V as a direct sum $\bigoplus_{i=1}^m k[x]/(q_i)$ of quotients of $k[x]$ by principal ideals (q_i) with each q_i of the form $p_i^{n_i}$, a power of a monic irreducible polynomial p_i over k (there is no room for any summands $k[x]$). Now given *any* monic polynomial $q = x^n + \sum_{i=0}^{n-1} q_i x^i \in k[x]$, there is an obvious basis for the quotient $W = k[x]/(q)$ consisting of the first n powers $1, x, \dots, x^{n-1}$ of x . Multiplication by x sends any power x^i to x^{i+1} if $i < n-1$, while $xx^{n-1} = x^n = -\sum_{i=0}^{n-1} q_i x^i$.

Accordingly the matrix of multiplication by x on W with respect to this basis is

$$\begin{pmatrix} 0 & 0 & 0 & \dots & -q_0 \\ 1 & 0 & 0 & \dots & -q_1 \\ 0 & 1 & 0 & \dots & -q_2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & -q_{n-1} \end{pmatrix}$$

We denote this last matrix by $C(q)$ and call it the **companion matrix of q** (p. 475). Clearly it has minimal polynomial q as an $n \times n$ matrix over k ; that is, q is the unique monic polynomial of least degree with $q(C(q)) = 0$. In particular, given any monic polynomial q of degree n , there is an explicit $n \times n$ matrix with minimal polynomial q . One checks directly that the characteristic polynomial of $C(q)$ is also $\pm q$. Combining the classification with the above choice of basis on each quotient $k[x]/(p_i^{n_i})$ above we get

Rational canonical form for matrices: Theorem 16, p. 477

Any square matrix M over k is similar to a block diagonal one B for which the blocks are companion matrices $C(p_i^{n_i})$ of powers of irreducible polynomials p_i over k . Two such matrices B are similar if and only if one can be obtained from the other by reordering the blocks.

The minimal polynomial of B is clearly the least common multiple of the powers $p_i^{n_i}$. As a consequence, given an irreducible polynomial p over k of degree d , the companion matrix $C(p^m)$ is up to similarity the only $md \times md$ matrix with minimal polynomial p^m ; all other matrices of this size will have minimal polynomials either involving a factor other than p or equal to a lower power of p .

Theorem 16 on p. 477 uses the invariant factor decomposition rather than the elementary divisor decomposition, so that the block diagonal factors take the form $C(q_1), \dots, C(q_m)$, where the q_i are monic polynomials with $q_1 | q_2 | \dots | q_m$. For most purposes the elementary divisor version is more convenient.

A further remarkable consequence is that **any square matrix M over k is similar to its transpose M^t** . Indeed, by putting M into rational canonical form we see that it is enough to prove this for companion matrices $C = C(p^m)$ of powers of irreducible polynomials p . But now a polynomial $q(C) = 0$ if and only if $q(C^t) = 0$, so C^t has the same minimal polynomial as C and must be similar to it. Note that any Math 208 student could understand the statement of this last result, but its proof is well beyond the scope of that course.

Now assume that k is algebraically closed, or more generally that all eigenvalues (roots of the characteristic polynomial) of a linear transformation T from k^n to itself lie in k . Then the only possible powers occurring in the rational canonical form of any matrix of T are powers $(x - a_i)^{n_i}$ of linear polynomials. In this case it is customary to use a different basis for $W = k[x]/((x - a_i)^{n_i})$, namely the one consisting of the powers $(x - a_i)^{n_i-1}, \dots, (x - a_i)^0$ in reverse order. The matrix of multiplication by x on W now takes the form

$$\begin{pmatrix} a_i & 1 & 0 & \dots & 0 \\ 0 & a_i & 1 & \dots & 0 \\ 0 & 0 & a_i & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a_i \end{pmatrix}$$

We call this matrix a **Jordan block** (with eigenvalue a_i).

Then we have

Jordan canonical form for matrices: Theorem 23, p. 493

Any square matrix M over a field k containing all of its eigenvalues is similar to a block diagonal matrix with Jordan blocks corresponding to eigenvalues of M . Two such block diagonal matrices are similar if and only if one can be obtained from the other by reordering the blocks.

In particular, for any element $a \in k$ and a fixed size n , there are only finitely many similarity classes of $n \times n$ matrices with a as their only eigenvalue. The similarity class of any such matrix is characterized by the sizes of its Jordan blocks. These form an unordered set of positive integers whose sum is n (a partition of n). Partitions are well-studied objects in enumerative combinatorics and a great deal is known about them. Another consequence of the Jordan form is the following result: any $n \times n$ complex invertible matrix M has a logarithm, so that

$$M = e^L = \sum_{i=0}^{\infty} \frac{L^i}{i!} \text{ for some complex matrix } L.$$

I will close by returning to the rational canonical form in a very special case. Let $q = p^n$ be a power of a prime p . I showed in class that some nonzero x in the finite field F_q of order q has multiplicative order $q - 1$ (assuming that such a field F_q exists). Hence any such x generates all of F_q over its prime subfield F_p , so that its minimal polynomial f over F_p necessarily has degree n . Then the matrix of multiplication by x , regarded as a linear transformation of F_q , also has multiplicative order $q - 1$ and minimal polynomial f , whence its rational canonical form must be $C(f)$. Thus $M = C(f)$ and its transpose $M^t C(f)^t$ are invertible $n \times n$ matrices over F_p having order $q - 1$.

In an upcoming homework assignment, you will use $C(f)^t$ to construct a **deBruijn sequence** of length q ; that is, a sequence a_0, \dots, a_{q-1} of elements of F_p such that its consecutive subsequences a_i, \dots, a_{i+n-1} of length n run through all sequences of n elements of F_p , each such sequence occurring exactly once. Here the index i runs from 0 to $q - 1$ and addition of subscripts takes place modulo q (so that the sequence “wraps around”). The construction uses the transpose M^t of the companion matrix M of the last slide. A crucial property is that **no power M^i or $(M^t)^i$ of M or M^t with $1 \leq i < q - 1$ has any eigenvalue equal to 1**, since M_i is the matrix of multiplication by $x^i \neq 1$ in F_q . Likewise for $(M^t)^i$, since its eigenvalues are the same as those of M^i .