Math 334 Sample Problems

One side of one notebook sized page of notes will be allowed on the test. You may work together on the sample problems – I encourage you to do that. The test will cover up to §3.3.

- 1. Find the maximum of xy^2z^3 on the set $\{(x, y, z) : x \ge 0, y \ge 0, z \ge 0, x + y + z = 12\}$.
- 2. Suppose that F(u) is a differentiable function with $F'(u) \neq 0$ and z = f(x, y) is a differentiable function that satisfies and

$$z = F(ax + by + cz),$$

where a, b, c are constants with $c \neq 0$. Prove

$$bf_x = af_y$$
.

- 3. Prove that the level set $y^2 = (x-1)(x-2)(x-3)$ is the disjoint union of two smooth connected curves. Prove that one of the curves is compact and the other is not.
- 4. Suppose that (x_0, y_0, z_0, u_0) satisfies the equations

$$x + y + z = F(u)$$

 $x^{2} + y^{2} + z^{2} = G(u)$
 $x^{3} + y^{3} + z^{3} = H(u)$,

where, F, G, H are C^1 in a neighborhood of u_0 . State a sufficient condition for being able to solve these equations for x, y, z as C^1 functions of u in a neighborhood of (x_0, y_0, z_0, u_0) .

- 5. Is the set $\{(x,y): y^2+x^2e^y=0\}$ a smooth curve? Is the set $\{(a\cos t,b\sin t): t\in (0,\pi)\}$, where a>0,b>0 a smooth curve?
- 6. Expand $(1 x + 2y)^3$ in powers of x 1 and y 2 in two different ways. The first way is by using algebra and the second way is by computing the Taylor polynomial of degree three centered at (1, 2).
- 7. Using the method of Lagrange multipliers, find the highest and lowest points of the circle

$$x^{2} + y^{2} + z^{2} = 16$$
, $(x+1)^{2} + (y+1)^{2} + (z+1)^{2} = 27$

8. Show that the surface $z = 3x^2 - 2xy + 2y^2$ lies entirely above every one of its tangent planes. Hint: Look at the Taylor expansion at every point.

1

Sample Problems 2

9. Let a > 0, b > 0 and a + b = 1. Also let x > 0, y > 0. prove that

$$x^a y^b \le ax + by,$$

by using the method of Lagrange multipliers applied to maximize $x^a y^b$ subject to ax + by = c, where c > 0 is some constant.

- 10. Let $f(x,y) = x^2(1+y)^3 + 7y^2$ define a function on \mathbb{R}^2 . Find and classify its critical points. What is $\sup\{f(x,y):(x,y)\in\mathbb{R}^2\}$? What is $\inf\{f(x,y):(x,y)\in\mathbb{R}^2\}$?
- 11. Define $f(x) = (\log x)^{\log x}$, for x > 1. Using the chain rule, compute f'(x).
- 12. Folland, §2.9, problem 16.
- 13. Suppose F(x,y) is a C^2 function that satisfies the equations F(x,y) = F(y,x), F(x,x) = x. Prove that the quadratic term in the Taylor polynomial of F based at the point (a,a) is $\frac{1}{2}F_{xx}(a,a)(x-y)^2$.
- 14. There may be homework problems or example problems from the text or lectures on the midterm.
- 15. The following topics have been covered since the first midterm:
 - (a) Chain rule.
 - (b) Mean value theorem.
 - (c) Higher order partials and equality of mixed partials.
 - (d) Taylor's theorem in one and several variables with Lagrange's form of the remainder.
 - (e) Behavior near critical points second derivative test for extrema in the case of two variables.
 - (f) Max-min problems with constraints. The method of Lagrange multipliers.
 - (g) Implicit Function Theorem.
 - (h) Smooth curves and surfaces.