Math 336 Sample Problems

One notebook sized page of notes (both sides) will be allowed on the test. The test will be comprehensive. The final exam is at 8:30 am, Monday, June 6, in the regular classroom.

1. Let p(z) be a polynomial of degree n. Let $M(r) = \max\{|p(z)| : |z| = r\}$. Let r > s > 0. Prove that

$$\frac{M(s)}{s^n} \ge \frac{M(r)}{r^n}.$$

- 2. Is there an analytic function f that maps |z| < 1 into |z| < 1 such that $f(\frac{1}{2}) = \frac{2}{3}, f(\frac{1}{4}) = \frac{1}{3}$?
- 3. Let $D = \{|z| < 1\}$. Suppose g is a real valued function on D and $0 \le g(z) \le |z|$. Suppose there is an $f \in \mathcal{O}(D)$ so that $|f(z)| = e^{g(z)}$. Prove that g is identically 0.
- 4. Suppose u_n is a sequence of harmonic functions on a domain W and suppose the sequence converges uniformly on compact sets to a function u. Prove that u is harmonic.

5. Let $f(z) = \frac{z-a}{1-\bar{a}z}$, where |a| < 1. Let $D = \{z : |z| < 1\}$. Prove that

(a)

$$\frac{1}{\pi} \int_D |f'(z)|^2 dx dy = 1.$$

(b)

$$\frac{1}{\pi} \int_D |f'(z)| dx dy = \frac{1 - |a|^2}{|a|^2} \log\left(\frac{1}{1 - |a|^2}\right).$$

Hint: Use the Poisson integral formula.

6. Let u(x, y), v(x, y) be continuously differentiable as functions of (x, y) in a domain Ω . Let f(z) = u(z) + iv(z). Suppose that for every $z_0 \in \Omega$ there is an r_0 (depending on z_0) such that

$$\int_{|z-z_0|=r} f(z)dz = 0,$$

for all r with $r < r_0$. Prove that f is analytic in Ω . Hint: Show that f satisfies the Cauchy-Riemann equations in Ω .

7. Compute

$$\int_{|z|=4} \frac{\sin z}{z^2} dz$$

Sample Problems

- 8. Let $D_2 = \{z : |z| < 2\}$ and $I = \{x \in \mathbf{R} : -1 \le x \le 1\}$. Find a bounded harmonic function u, defined in $D_2 I$ such that u does not extend to a harmonic function defined in all of D_2 .
- 9. Suppose f is analytic on $D = \{|z| < 1\}$ and f(0) = 0. Prove that

$$\sum f(z^n)$$

converges uniformly on compact subsets of D.

10. Let a_k be a sequence of distinct complex numbers such that $\sum_{k=1}^{\infty} \frac{1}{|a_k|}$ converges. Let $A = \{a_k : k = 1, ..., \infty\}$. Prove that

$$\sum_{k=1}^{\infty} \frac{1}{z - a_k}$$

converges to an analytic function on $\mathbb{C} - A$.

- 11. Let f and g be entire functions so that satisfy $f^2 + g^2 = 1$. Prove that there is an entire function h so that $f = \cos(h), g = \sin(h)$.
- 12. Find a function, h(x, y), harmonic in $\{x > 0, y > 0\}$, such that

$$h(x,y) = \begin{cases} 0 & \text{if } 0 < x < 2, y = 0, \\ 1 & \text{if } x > 2, y = 0, \\ 2 & \text{if } x = 0, y > 0 \end{cases}$$

- 13. Suppose that u is harmonic on all of \mathbb{C} and $u \ge 0$. Prove that u is constant.
- 14. Suppose f is analytic on $H = \{z = x + iy : y > 0\}$ and suppose $|f(z)| \le 1$ on H and f(i) = 0. Prove

$$|f(z)| \le \left|\frac{z-i}{z+i}\right|.$$

15. Compute

$$\int_0^\infty \frac{1 - \cos x}{x^2} dx.$$

16. Let f be a non-constant analytic function on the connected open set W. Let $Z = \{z : f(z) = 0\}$. Prove that W - Z is connected.

Sample Problems

- 17. (a) Prove that 1/z does not have an analytic antiderivative on $\mathbb{C} \{0\}$.
 - (b) Find all integers $0, \pm 1, \pm 2, \ldots$ such that the function $z^n e^{1/z}$ has an analytic antiderivative on $\mathbb{C} \{0\}$.
- 18. Find the radius of convergence of

$$\sum \frac{n^n}{n!} z^{2n}.$$

- 19. Suppose $f \in \mathcal{O}(0 < |z a| < \epsilon)$ and that $\operatorname{Re}(f)$ is bounded. Prove that a is a removable singularity.
- 20. Let f be a non-constant analytic function defined on $\{|z| < 1\}$ such that $\operatorname{Re}(f(z)) \ge 0$.
 - (a) Prove that $\operatorname{Re}(f(z)) > 0$.
 - (b) Suppose f(0) = 1. Prove that

$$\frac{1-|z|}{1+|z|} \le |f(z)| \le \frac{1+|z|}{1-|z|}.$$

21. Suppose f and g are analytic on a connected open set Ω .

- (a) If |f(z)| + |g(z)| is constant, then both f and g are constant.
- (b) If |f(z)| + |g(z)| assumes a local maximum in Ω , then f and g are constant.

22. Prove that
$$\sum_{1}^{\infty} \frac{\sin nz}{2^n}$$
 represents an analytic function on $|\operatorname{Im}(z)| < \log 2$

- 23. Find all real valued harmonic functions on \mathbb{C} that are constant on vertical lines (the constant may depend on the line).
- 24. Let f and g be two analytic functions on an open connected set W. Suppose that $f(z)\overline{g(z)}$ is real for all $z \in W$. Prove that either f = cg or g is identically 0.
- 25. (a) Prove that the series

$$\sum_{1}^{\infty} 2^{-n^2} z^{2^n}$$

converges uniformly on $|z| \leq 1$.

- (b) Prove that the radius of convergence of the series is 1.
- 26. There may be homework problems or example problems from the text on the final. Don't forget previous sample problems.