

Highly neighborly centrally symmetric spheres

Isabella Novik*

Department of Mathematics
University of Washington
Seattle, WA 98195-4350, USA
novik@uw.edu

Hailun Zheng

Department of Mathematics
University of Michigan
Ann Arbor, Michigan 48109-1043, USA
hailunz@umich.edu

January 29, 2020

Abstract

In 1995, Jockusch constructed an infinite family of centrally symmetric 3-dimensional simplicial spheres that are *cs-2-neighborly*. Here we generalize his construction and show that for all $d \geq 3$ and $n \geq d + 1$, there exists a centrally symmetric d -dimensional simplicial sphere with $2n$ vertices that is *cs- $\lfloor d/2 \rfloor$ -neighborly*. This result combined with work of Adin and Stanley completely resolves the upper bound problem for centrally symmetric simplicial spheres.

1 Introduction

In this paper we construct highly neighborly centrally symmetric d -dimensional spheres with an arbitrarily large even number of vertices. A simplicial complex is *centrally symmetric* (*cs*, for short) if it possesses a free simplicial involution. We refer to a pair of vertices that form an orbit under this involution as antipodes or antipodal vertices. One large class of examples is given by the boundary complexes of *cs* simplicial polytopes: a polytope P is *cs* if $P = -P$; the involution, in this case, is induced by the map $v \mapsto -v$ on the vertices.

A (non-*cs*) simplicial complex is called ℓ -neighborly if every ℓ of its vertices form a face. For instance, the boundary complex of the $(d + 1)$ -dimensional simplex is $(d + 1)$ -neighborly, while the boundary complex of the $(d + 1)$ -dimensional cyclic polytope with $n \geq d + 3$ vertices, $C(d + 1, n)$, is $\lfloor d/2 \rfloor$ -neighborly. The interest in neighborly polytopes arises from the celebrated upper bound theorem [12, 22] asserting that among all d -dimensional simplicial spheres with n vertices, the boundary complex of $C(d+1, n)$ simultaneously maximizes all the face numbers. The cyclic polytope in this statement can be replaced with any $\lfloor d/2 \rfloor$ -neighborly d -dimensional simplicial sphere — the objects that abound in nature, see [18].

This notion of ℓ -neighborliness can be easily modified for the class of *cs* complexes: a *cs* simplicial complex Δ is *cs- ℓ -neighborly* if every set of ℓ of its vertices, no two of which are antipodes, is a face of Δ . Furthermore, the same notion applies to any (not necessarily *cs*) full-vertex subcomplex Γ of Δ . For instance, the boundary complex of the $(d+1)$ -dimensional cross-polytope is *cs- $(d+1)$ -neighborly*, while the boundary complex of the same cross-polytope with one facet removed is *cs- d -neighborly*. Adin [1] and Stanley (unpublished) proved that in a complete analogy with Stanley's upper bound

*Research is partially supported by NSF grant DMS-1664865, and by Robert R. & Elaine F. Phelps Professorship in Mathematics.

theorem, among all cs simplicial spheres of dimension d and with $2n$ vertices, a cs- $\lceil d/2 \rceil$ -neighborly sphere simultaneously maximizes all the face numbers, *assuming such a sphere exists*. (See [16, 17] for an extension of this result to cs simplicial manifolds.)

Thus, two natural questions to consider are: do there exist cs simplicial polytopes of dimension $d + 1 \geq 4$ with arbitrarily many vertices whose boundary complexes are cs- $\lceil d/2 \rceil$ -neighborly? Do there exist cs simplicial spheres of dimension $d \geq 3$ with arbitrarily many vertices that are cs- $\lceil d/2 \rceil$ -neighborly?

The answer to the first question was given by McMullen and Shephard [13] more than fifty years ago: extending the 4-dimensional case worked out by Grünbaum [4, p. 116], they proved that while there do exist cs $(d + 1)$ -dimensional polytopes with $2(d + 2)$ vertices that are cs- $\lceil d/2 \rceil$ -neighborly, a cs $(d + 1)$ -dimensional polytope with $2(d + 3)$ vertices cannot be more than cs- $\lfloor (d + 2)/3 \rfloor$ -neighborly. Moreover, according to [10], a cs $(d + 1)$ -dimensional polytope with more than 2^{d+1} vertices cannot be even cs-2-neighborly.

The second question remained a total mystery until in 1995 Jockusch [6] showed that, in a sharp contrast with the situation for cs 4-dimensional polytopes, for *every* value of $n \geq 4$, there exists a cs 3-dimensional sphere with $2n$ vertices that is cs-2-neighborly. In addition, for $d \leq 6$, Lutz [11] found (by a computer search) several cs d -dimensional spheres with $2(d + 3)$ vertices that are cs- $\lceil d/2 \rceil$ -neighborly.

Here, we build on work of Jockusch to provide a complete answer to the second question: for *all* values of $d \geq 3$ and $n \geq d + 1$, there exists a cs d -dimensional combinatorial sphere with $2n$ vertices, Δ_n^d , that is cs- $\lceil d/2 \rceil$ -neighborly. Thus, our result combined with work of Adin and Stanley completely resolves the upper bound problem for cs simplicial spheres. (At the same time, there is not even a plausible upper bound conjecture for cs polytopes.)

Our construction is by induction on both d and n . The key idea in constructing Δ_{n+1}^d from Δ_n^d is to define for each $i \leq \lceil d/2 \rceil - 1$, an auxiliary d -dimensional ball with $2n$ vertices, $B_n^{d,i} \subset \Delta_n^d$, that is both i -stacked and cs- i -neighborly, see Sections 2 and 3 for definitions. For $d = 3$, our construction reduces to Jockusch's construction. It is worth mentioning that using the same balls in fact allows us to construct for any $\ell \leq \lceil d/2 \rceil$, a family of cs d -dimensional combinatorial spheres that are cs- ℓ -neighborly but not cs- $(\ell + 1)$ -neighborly.

The structure of the paper is as follows. In Section 2 we discuss several basic facts and definitions pertaining to simplicial complexes and PL topology. Section 3 is a hard duty section that contains our inductive construction and the proof that this construction works. In Section 4, we study some other properties of the spheres Δ_n^d . We close in Section 5 with several remarks and open questions.

2 Preliminaries

In this section we review some background related to simplicial complexes and PL topology. For all undefined terminology we refer the readers to [2].

A *simplicial complex* Δ with vertex set $V = V(\Delta)$ is a collection of subsets of V that is closed under inclusion and contains all singletons: $\{v\} \in \Delta$ for all $v \in V$. The elements of Δ are called *faces*. The *dimension of a face* $\tau \in \Delta$ is $\dim \tau := |\tau| - 1$. The *dimension of Δ* , $\dim \Delta$, is the maximum dimension of its faces. A face of a simplicial complex Δ is a *facet* if it is maximal w.r.t. inclusion. We say that Δ is *pure* if all facets of Δ have the same dimension.

Let V be a set of size $d + 1$. Two fundamental examples of pure simplicial complexes are the d -dimensional simplex on V , $\bar{V} := \{\tau : \tau \subseteq V\}$, and its boundary complex, $\partial \bar{V} := \{\tau : \tau \subsetneq V\}$.

To simplify the notation, for a face that is a vertex, an edge, or a triangle, we write v , uv , and uvw instead of $\{v\}$, $\{u, v\}$, and $\{u, v, w\}$, respectively. We denote by (v_1, v_2, \dots, v_n) a path with edges $v_1v_2, v_2v_3, \dots, v_{n-1}v_n$ if $v_n \neq v_1$, or a cycle if $v_n = v_1$. In particular, a path of length one (v_1, v_2) is a 1-dimensional simplex, so it can also be written as $\overline{v_1v_2}$.

Let Δ be a simplicial complex. The k -skeleton of Δ , $\text{Skel}_k(\Delta)$, is the subcomplex of Δ consisting of all faces of dimension $\leq k$. If τ is a face of Δ , then the *link of τ in Δ* is the following subcomplex of Δ :

$$\text{lk}(\tau, \Delta) := \{\sigma \in \Delta : \sigma \cap \tau = \emptyset \text{ and } \sigma \cup \tau \in \Delta\}.$$

Finally, if Δ is pure and Γ is a full-dimensional pure subcomplex of Δ , then $\Delta \setminus \Gamma$ is the subcomplex of Δ generated by those facets of Δ that are not in Γ .

If Δ and Γ are simplicial complexes on disjoint vertex sets, then the *join* of Δ and Γ is the simplicial complex $\Delta * \Gamma = \{\sigma \cup \tau : \sigma \in \Delta \text{ and } \tau \in \Gamma\}$. Two important special cases are the *cone* over Δ with apex v defined as the join $\Delta * \bar{v}$ and the *suspension* of Δ , $\Sigma\Delta$, defined as the join of Δ with a 0-dimensional sphere. In the rest of the paper, we write $\Delta * v$ in place of $\Delta * \bar{v}$.

Let V be a set of size $d + 1$ and let \bar{V} be the d -dimensional simplex on V . A *combinatorial d -ball* is a simplicial complex PL homeomorphic to \bar{V} . Similarly, a *combinatorial $(d - 1)$ -sphere* is a simplicial complex PL homeomorphic to $\partial\bar{V}$.

One advantage of working with combinatorial balls and spheres is that they satisfy several natural properties that fail in the class of simplicial balls and spheres. For instance, if Δ is a combinatorial d -sphere and $\Gamma \subset \Delta$ is a combinatorial d -ball, then so is $\Delta \setminus \Gamma$, see [5]. Furthermore, the link of any face in a combinatorial sphere is a combinatorial sphere. On the other hand, the link of a face τ in a combinatorial d -ball B is either a combinatorial ball or a combinatorial sphere; in the former case we say that τ is a *boundary face* of B , and in the latter case that τ is an *interior face* of B . The *boundary complex* of B , ∂B , is the subcomplex of B that consists of all boundary faces of B ; in particular, ∂B is a combinatorial $(d - 1)$ -sphere.

The following lemma summarizes a few basic but very useful properties of combinatorial balls. We only prove the last part; the proofs of the first two parts along with additional information on PL topology can be found in [5], see also [9].

Lemma 2.1.

1. *Let B and B' be combinatorial balls. Then $B * B'$ is a combinatorial ball; its interior faces are sets of the form $F \cup F'$, where F is an interior face of B and F' is an interior face of B' . Furthermore, the cone over ∂B , $\partial B * v$, is a combinatorial ball; its boundary complex is ∂B .*
2. *Let B and B' be combinatorial d -balls such that $B \cap B' \subseteq \partial B \cap \partial B'$ is a combinatorial $(d - 1)$ -ball. Then $B \cup B'$ is a combinatorial d -ball. The set of interior faces of $B \cup B'$ consists of the interior faces of B , the interior faces of B' , and the interior faces of $B \cap B'$.*
3. *Assume that combinatorial balls B and B' are full-dimensional subcomplexes of a combinatorial sphere Γ . If B and B' share a common interior face, then they share a facet.*

Proof: For part 3, let τ be a common interior face of B and B' . Assume that $\dim \Gamma = d$ and $\dim \tau = k$. Then the link of τ in B is a combinatorial $(d - k - 1)$ -sphere, and so is the link of τ in Γ . Furthermore, the link of τ in B is contained in the link of τ in Γ . Thus, these two links must be equal. In particular, every facet F of Γ containing τ must be a facet of B . By the same argument, such an F must also be a facet of B' . The result follows. \square

A simplicial complex Δ is *centrally symmetric* or *cs* if its vertex set is endowed with a *free involution* $\alpha : V(\Delta) \rightarrow V(\Delta)$ that induces a free involution on the set of all nonempty faces of Δ . In more detail, for all nonempty faces $\tau \in \Delta$, the following holds: $\alpha(\tau) \in \Delta$, $\alpha(\tau) \neq \tau$, and $\alpha(\alpha(\tau)) = \tau$. To simplify notation, we write $\alpha(\tau) = -\tau$ and refer to τ and $-\tau$ as *antipodal faces* of Δ . Similarly, if Γ is a subcomplex of Δ we write $-\Gamma$ in place of $\alpha(\Gamma)$.

One example of a cs combinatorial d -sphere is the boundary complex of the $(d+1)$ -dimensional *cross-polytope*, $\partial\mathcal{C}_{d+1}^*$. The polytope \mathcal{C}_{d+1}^* is the convex hull of $\{\pm e_1, \pm e_2, \dots, \pm e_{d+1}\}$, where e_1, e_2, \dots, e_{d+1} are the endpoints of the standard basis in \mathbb{R}^{d+1} . As an abstract simplicial complex, $\partial\mathcal{C}_{d+1}^*$ is the $(d+1)$ -fold suspension of $\{\emptyset\}$. Equivalently, it is the collection of all subsets of $V_{d+1} := \{\pm v_1, \dots, \pm v_{d+1}\}$ that contain at most one vertex from each pair $\{\pm v_i\}$. In particular, every cs simplicial complex with vertex set V_n is a subcomplex of $\partial\mathcal{C}_n^*$.

We close this section with a discussion of neighborliness and stackedness. Let $\Delta \subseteq \partial\mathcal{C}_n^*$ be a simplicial complex, possibly non-cs, and let $1 \leq i \leq n$. We say that Δ is *cs- i -neighborly* (w.r.t. V_n), if $\text{Skel}_{i-1}(\Delta) = \text{Skel}_{i-1}(\partial\mathcal{C}_n^*)$. For $i = 1$, this simply means that $V(\Delta) = V_n$. For convenience, we also refer to simplices (i.e., faces of $\partial\mathcal{C}_n^*$) as cs-0-neighborly complexes.

A combinatorial d -ball B is called *i -stacked* (for some $0 \leq i \leq d$), if all interior faces of B are of dimension $\geq d - i$, that is, $\text{Skel}_{d-i-1}(B) = \text{Skel}_{d-i-1}(\partial B)$. In particular, 0-stacked balls are simplices, and 1-stacked balls are also known as stacked balls. The notion of stackedness takes its origins in the generalized lower bound theorem [14, 15, 23].

The following lemmas will be handy.

Lemma 2.2. *Let B_1 and B_2 be combinatorial balls of dimension d_1 and d_2 , respectively. If B_1 is i_1 -stacked and B_2 is i_2 -stacked, then*

1. *The complex $B_1 * B_2$ is an $(i_1 + i_2)$ -stacked combinatorial $(d_1 + d_2 + 1)$ -ball.*
2. *Furthermore, if $d_1 = d_2 = d$, $i_1 \leq i_2$, and $B_1 \cap B_2 \subseteq \partial B_1 \cap \partial B_2$ is a combinatorial $(d-1)$ -ball that is i_3 -stacked for some $i_3 < i_2$, then $B_1 \cup B_2$ is an i_2 -stacked combinatorial d -ball.*

Proof: For part 1, observe that by definition of stackedness, all interior faces of B_j have dimension $\geq d_j - i_j$ for $j = 1, 2$. Hence by Lemma 2.1, the interior faces of $B_1 * B_2$, which is a combinatorial $(d_1 + d_2 + 1)$ -ball, have dimension $\geq (d_1 + d_2) - (i_1 + i_2) + 1$. Thus, $B_1 * B_2$ is $(i_1 + i_2)$ -stacked.

Part 2 similarly follows from Lemma 2.1 and the definition of stackedness. Indeed, the interior faces of both B_1 and B_2 have dimension $\geq d - i_2$. All other interior faces of $B_1 \cup B_2$ are interior faces of $B_1 \cap B_2$, and so have dimension $\geq (d-1) - i_3 \geq d - i_2$. Hence $B_1 \cup B_2$ is i_2 -stacked. \square

Lemma 2.3. *Let $k \geq 1$ be an integer. Let $\Delta \subseteq \partial\mathcal{C}_n^*$ be a combinatorial $(2k-1)$ -sphere that is cs- k -neighborly w.r.t. V_n , and let $B \subseteq \Delta$ be a combinatorial $(2k-1)$ -ball that is both cs- $(k-1)$ -neighborly w.r.t. V_n and $(k-1)$ -stacked. Then $\Delta \setminus B$ is a combinatorial $(2k-1)$ -ball that is cs- k -neighborly and k -stacked.*

Proof: Let $F \in \partial\mathcal{C}_n^*$ be any set of size $\leq k$. Since Δ is cs- k -neighborly, F is a face of Δ . Thus, if $F \notin B$, then $F \in \Delta \setminus B$. On the other hand, if $F \in B$, then since $\dim F \leq k-1 < (2k-1) - (k-1)$ and since B is $(k-1)$ -stacked, F must be a boundary face of B and thus also a face of $\Delta \setminus B$. We infer that $\Delta \setminus B$ is cs- k -neighborly.

Next let F be a face of $\Delta \setminus B$ of dimension $< (2k-1) - k = k-1$, i.e., $|F| \leq k-1$. Since B is cs- $(k-1)$ -neighborly, it follows that $F \in B$. Thus F must lie on the boundary of B and hence also on the boundary of $\Delta \setminus B$. We conclude that $\Delta \setminus B$ is k -stacked. \square

3 The construction

Our goal is to construct Δ_n^d — a cs combinatorial d -sphere with $2n \geq 2d + 2$ vertices that is cs- $\lceil d/2 \rceil$ -neighborly. Our method is to build a certain ball on those $2n$ vertices that is both cs- $(\lceil d/2 \rceil - 1)$ -neighborly and $(\lceil d/2 \rceil - 1)$ -stacked. The following lemma explains the significance of these balls and outlines the inductive procedure on n we will use once such balls are constructed. For all values of $d + 1 \leq n$, the vertex set of Δ_n^d will be $V_n = V(\partial\mathcal{C}_n^*)$. In particular, our initial complex Δ_{d+1}^d is $\partial\mathcal{C}_{d+1}^*$.

Lemma 3.1. *Let $d \geq 1$ and $1 \leq i \leq \lceil d/2 \rceil$ be integers. Assume that $\Delta_n^{d,i}$ is a cs combinatorial d -sphere with $V(\Delta_n^{d,i}) = V_n$ that is cs- i -neighborly. Assume further that $B_n^{d,i-1} \subseteq \Delta_n^{d,i}$ is a combinatorial d -ball that satisfies the following properties:*

- the ball $B_n^{d,i-1}$ is both cs- $(i-1)$ -neighborly w.r.t. V_n and $(i-1)$ -stacked, and
- the balls $B_n^{d,i-1}$ and $-B_n^{d,i-1}$ share no common facets.

Then the complex $\Delta_{n+1}^{d,i}$ obtained from $\Delta_n^{d,i}$ by replacing $B_n^{d,i-1}$ with $\partial B_n^{d,i-1} * v_{n+1}$ and $-B_n^{d,i-1}$ with $\partial(-B_n^{d,i-1}) * (-v_{n+1})$ is a cs combinatorial d -sphere with $V(\Delta_{n+1}^{d,i}) = V_{n+1}$ that is cs- i -neighborly.

Proof: Observe that $B_n^{d,i-1}$ and $\partial B_n^{d,i-1} * v_{n+1}$ are two combinatorial balls with the same boundary. The fact that $B_n^{d,i-1}$ and $-B_n^{d,i-1}$ share no common facets combined with Lemma 2.1 then implies that $\Delta_{n+1}^{d,i}$ is a combinatorial sphere. Moreover, the definition of $\Delta_{n+1}^{d,i}$ along with the fact that $\Delta_n^{d,i}$ is a cs complex guarantees that the complex $\Delta_{n+1}^{d,i}$ is also cs.

To show that $\Delta_{n+1}^{d,i}$ is cs- i -neighborly, consider a set $F \in \partial\mathcal{C}_{n+1}^*$ with $|F| \leq i \leq \lceil d/2 \rceil$. If v_{n+1} is in F , then $F \setminus v_{n+1}$ is a face of $\partial\mathcal{C}_n^*$ of size at most $i-1$. Since $B_n^{d,i-1}$ is cs- $(i-1)$ -neighborly and $(i-1)$ -stacked, it follows that

$$F \setminus v_{n+1} \in \text{Skel}_{i-2}(B_n^{d,i-1}) \subseteq \text{Skel}_{d-i}(B_n^{d,i-1}) \subseteq \partial B_n^{d,i-1}.$$

Hence $F \in \partial B_n^{d,i-1} * v_{n+1} \subseteq \Delta_{n+1}^{d,i}$. If $-v_{n+1} \in F$, then by the above argument, $-F \in \Delta_{n+1}^{d,i}$, and so by symmetry, $F \in \Delta_{n+1}^{d,i}$. Finally, if $\pm v_{n+1} \notin F$, then since $\Delta_n^{d,i}$ is cs- i -neighborly, $F \in \Delta_n^{d,i}$. As any face of $\pm B_n^{d,i-1}$ of dimension $\leq i-1$ is on the boundary of $\pm B_n^{d,i-1}$, we conclude that $F \in \Delta_n^{d,i} \setminus \pm B_n^{d,i-1} \subseteq \Delta_{n+1}^{d,i}$. \square

The idea is to build $B_n^{d,\lceil d/2 \rceil - 1}$ from balls that are less cs-neighborly using intertwined induction.

Definition 3.2. Let $d \geq 1$, $i \leq \lceil d/2 \rceil$, and $n \geq d + 1$ be integers. Define Δ_n^d and $B_n^{d,i}$ inductively as follows:

- For the initial cases, define $\Delta_n^1 := (v_1, v_2, \dots, v_n, -v_1, -v_2, \dots, -v_n, v_1)$, $\Delta_{d+1}^d := \partial\mathcal{C}_{d+1}^*$, $B_n^{d,j} := \emptyset$ if $j < 0$, and $B_n^{1,0} := \overline{(-v_1)v_n}$. (In particular, $B_n^{1,j} \subseteq \Delta_n^{1,1}$ for all $j \leq 0$.)
- Assume that Δ_m^{d-1} and $B_m^{d-1,i} \subseteq \Delta_m^{d-1}$ are already defined for all $i \leq \lfloor (d-1)/2 \rfloor$ and $m \geq d$. If $d = 2k$, then define $B_m^{2k-1,k} := \Delta_m^{2k-1} \setminus B_m^{2k-1,k-1}$ (for $m \geq 2k$). Then, for all $n \geq d + 1$ and $i \leq \lceil d/2 \rceil$, define

$$B_n^{d,i} := \left(B_{n-1}^{d-1,i} * v_n \right) \cup \left((-B_{n-1}^{d-1,i-1}) * (-v_n) \right).$$

- If Δ_n^d is already defined (and assuming that $\Delta_n^d \supseteq B_n^{d, \lceil d/2 \rceil - 1}$ and also that $B_n^{d, \lceil d/2 \rceil - 1}$ is a combinatorial ball that shares no common facets with $-B_n^{d, \lceil d/2 \rceil - 1}$), define Δ_{n+1}^d as the complex obtained from Δ_n^d by replacing $B_n^{d, \lceil d/2 \rceil - 1}$ with $\partial B_n^{d, \lceil d/2 \rceil - 1} * v_{n+1}$ and $-B_n^{d, \lceil d/2 \rceil - 1}$ with $\partial(-B_n^{d, \lceil d/2 \rceil - 1}) * (-v_{n+1})$.

To get the feel for this construction, we start by computing several explicit examples of complexes produced by Definition 3.2. Note that the join of any simplicial complex with the void complex \emptyset is the void complex. Hence by induction, for all $d \geq 1$, $B_n^{d,0}$ is the simplex on the vertex set $\{-v_1, v_{n-d+1}, v_{n-d+2}, \dots, v_n\}$. In particular, the link of v_{i+1} in Δ_{i+1}^2 is $\partial B_i^{2,0}$, the boundary of the triangle $\{-v_1, v_{i-1}, v_i\}$; similarly, the link of $-v_{i+1}$ is the boundary of $\{v_1, -v_{i-1}, -v_i\}$. It follows that Δ_n^2 is obtained from $\partial \mathcal{C}_3^*$ by symmetric stacking — an operation defined in [7].

To get a handle on $B_n^{3,1}$, it is worth noting that for $i \leq \lceil d/2 \rceil - 1$, Definition 3.2 implies that

$$\begin{aligned} B_n^{d,i} &= \left(B_{n-2}^{d-2,i} * (v_{n-1}, v_n) \cup (-B_{n-2}^{d-2,i-1}) * (-v_{n-1}, v_n) \right) \\ &\quad \cup \left((-B_{n-2}^{d-2,i-1}) * (-v_{n-1}, -v_n) \cup B_{n-2}^{d-2,i-2} * (v_{n-1}, -v_n) \right) \\ &= \left(B_{n-2}^{d-2,i} * (v_{n-1}, v_n) \right) \cup \left((-B_{n-2}^{d-2,i-1}) * (v_n, -v_{n-1}, -v_n) \right) \cup \left(B_{n-2}^{d-2,i-2} * (v_{n-1}, -v_n) \right). \end{aligned}$$

If we let $d = 3$ and $i = 1$, then

$$B_n^{3,1} = \left((v_{n-2}, v_{n-3}, \dots, v_1, -v_{n-2}, -v_{n-3}, \dots, -v_1) * (v_{n-1}, v_n) \right) \cup \left((v_1, -v_{n-2}) * (v_n, -v_{n-1}, -v_n) \right).$$

In particular, $B_n^{3,1}$ is a combinatorial 3-ball that has $2n - 3$ facets; it is cs-1-neighborly w.r.t. V_n and 1-stacked. This complex plays a key role in Jockusch's construction [6] of cs-2-neighborly 3-spheres: in fact, Jockusch's 3-spheres are exactly the complexes Δ_n^3 , for $n \geq 4$.

Several other easy consequences of Definition 3.2 are: if $i \leq \lfloor d/2 \rfloor$ and if $B_n^{d,i}$ is well-defined, then every facet of $B_n^{d,i}$ contains either v_n or $-v_n$, and no face of $B_n^{d,i}$ contains two antipodal vertices. This latter property and the fact that $\Delta_{d+1}^d = \partial \mathcal{C}_{d+1}^*$ is cs- $(d+1)$ -neighborly implies that Δ_{d+1}^d does contain all balls $B_{d+1}^{d,i}$ for $i \leq \lfloor d/2 \rfloor$, so we can at least start executing the algorithm in Definition 3.2 for all d (once we have taken care of smaller dimensions).

Our remaining task is to show that this algorithm never gets stuck and that its output, Δ_n^d , is a cs combinatorial d -sphere with $2n$ vertices that is cs- $\lfloor d/2 \rfloor$ -neighborly. To start, we verify in Lemma 3.3 that if Definition 3.2 allowed us to reach a point where the complex Δ_n^d and its subcomplex $B_n^{d, \lceil d/2 \rceil - 1}$ were constructed, then all complexes produced by the definition up to that point satisfy all the assumptions of Lemma 3.1. This allows us to advance one more step and construct Δ_{n+1}^d . We then need to show that Δ_{n+1}^d produced in this way contains $B_{n+1}^{d, \lceil d/2 \rceil - 1}$.

Lemma 3.3. *If the algorithm reached the (d, n) -th step and produced a pair $\Delta_n^d \supseteq B_n^{d, \lceil d/2 \rceil - 1}$, then Δ_n^d is a cs combinatorial d -sphere with vertex set V_n that is cs- $\lfloor d/2 \rfloor$ -neighborly while the complexes $B_n^{d,i}$ (for $0 \leq i \leq \lfloor d/2 \rfloor$) are combinatorial d -balls that are cs- i -neighborly w.r.t. V_n and i -stacked. Furthermore, they satisfy the following “nesting property”: $-B_n^{d,i-1} \subseteq B_n^{d,i}$ (for all $i \leq \lfloor d/2 \rfloor$). Finally, for all $i \leq \lfloor d/2 \rfloor$, $B_n^{d,i}$ and $-B_n^{d,i}$ share no common facets.*

Proof: We verify the properties of Δ_n^d and $B_n^{d,i}$ by induction on the dimension and the number of vertices. Since $\Delta_{d+1}^d = \partial \mathcal{C}_{d+1}^*$, the complex Δ_{d+1}^d is indeed a cs combinatorial d -sphere with

$V(\Delta_{d+1}^d) = V_{d+1}$ that is $\text{cs-}\lceil d/2 \rceil$ -neighborly. Furthermore, for any $m \geq 2$, $B_m^{1,0}$ and $B_m^{1,1}$ satisfy all the conditions: $B_m^{1,0}$ is an edge, and hence it is a $\text{cs-}0$ -neighborly and 0 -stacked 1 -ball, while $B_m^{1,1}$ is a path on V_m , and hence a $\text{cs-}1$ -neighborly (w.r.t. V_m) and 1 -stacked 1 -ball. Finally, $-B_m^{1,0} = \overline{v_1(-v_m)}$ and

$$B_m^{1,1} = \Delta_m^1 \setminus B_m^{1,0} = (-v_1, -v_2, \dots, -v_m, v_1, v_2, \dots, v_m) \supseteq -B_m^{1,0}.$$

For the inductive step, since the algorithm reached the (d, n) -th step, we can assume that the complexes $B_{n-1}^{d-1,j}$ (for $j \leq \lceil (d-1)/2 \rceil$) satisfy all the conditions of the lemma and that if $n > d+2$, then Δ_{n-1}^d is a cs combinatorial d -sphere (with vertex set V_{n-1}) that is $\text{cs-}\lceil d/2 \rceil$ -neighborly. We now show that then the same holds for $B_n^{d,i}$ (for all $0 \leq i \leq \lceil d/2 \rceil$) and Δ_n^d .

We start with the nesting property. By definition, for all $i \leq \lfloor d/2 \rfloor = \lceil (d-1)/2 \rceil$,

$$\begin{aligned} -B_n^{d,i-1} &= \left((-B_{n-1}^{d-1,i-1}) * (-v_n) \right) \cup \left(B_{n-1}^{d-1,i-2} * v_n \right), \quad \text{and} \\ B_n^{d,i} &= \left((-B_{n-1}^{d-1,i-1}) * (-v_n) \right) \cup \left(B_{n-1}^{d-1,i} * v_n \right). \end{aligned}$$

By the inductive hypothesis, $B_{n-1}^{d-1,i-2}$ is a subcomplex of $B_{n-1}^{d-1,i}$. Hence $-B_n^{d,i-1} \subseteq B_n^{d,i}$ for $i \leq \lfloor d/2 \rfloor$. We will treat the case of $d = 2k - 1$, $i = k$ a bit later.

Next, it follows from the nesting property and the definition of $B_{n-1}^{d-1,j}$ that $B_{n-1}^{d-1,i}$ and $B_{n-1}^{d-1,i-1}$ share no common facets for all $1 \leq i \leq \lfloor d/2 \rfloor \leq \lceil (d-1)/2 \rceil$, and hence neither do $B_n^{d,i}$ and $-B_n^{d,i}$. This result also implies the nesting property for $d = 2k - 1$ and $i = k$: since $B_n^{2k-1,k-1}$ and $-B_n^{2k-1,k-1}$ have no common facets, $-B_n^{2k-1,k-1} \subseteq \Delta_n^{2k-1} \setminus B_n^{2k-1,k-1} = B_n^{2k-1,k}$.

We now show that $B_n^{d,i}$ is an i -stacked combinatorial d -ball. For the case of $i \leq \lfloor d/2 \rfloor$, recall that $B_n^{d,i}$ is the union of $D_1 = B_{n-1}^{d-1,i} * v_n$ and $D_2 = (-B_{n-1}^{d-1,i-1}) * (-v_n)$. By the inductive hypothesis and Lemma 2.1, D_1 and D_2 are combinatorial d -balls. The nesting property implies that $D_1 \cap D_2 = -B_{n-1}^{d-1,i-1}$ and by the inductive hypothesis it is also a combinatorial $(d-1)$ -ball. Since it is contained in $\partial D_1 \cap \partial D_2$, Lemma 2.1 guarantees that $B_n^{d,i} = D_1 \cup D_2$ is a combinatorial d -ball. Furthermore, by the inductive hypothesis and Lemma 2.2, D_1 is i -stacked, D_2 is $(i-1)$ -stacked and the intersection is $(i-1)$ -stacked. Thus the union $B_n^{d,i}$ is i -stacked.

Next we turn to treating cs-neighborliness : we show that for all $i \leq \lfloor d/2 \rfloor$, $B_n^{d,i}$ is $\text{cs-}i$ -neighborly w.r.t. V_n while Δ_n^d is a cs combinatorial d -sphere that is $\text{cs-}\lceil d/2 \rceil$ -neighborly w.r.t. V_n . The statement about $B_n^{d,i}$ follows easily from the definition of $B_n^{d,i}$ and the inductive hypothesis asserting that $B_{n-1}^{d-1,j}$ is $\text{cs-}j$ -neighborly w.r.t. V_{n-1} for $j = i-1, i$. Now, if $n = d+1$, then $\Delta_n^d = \partial \mathcal{C}_{d+1}^*$, so it is a cs combinatorial d -sphere on V_n that is $\text{cs-}\lceil d/2 \rceil$ -neighborly, and if $n > d+1$, then the inductive hypothesis on Δ_{n-1}^d along with Lemma 3.1 and the established properties of $B_{n-1}^{d,\lceil d/2 \rceil-1}$ imply that Δ_n^d is a cs combinatorial d -sphere on V_n that is $\text{cs-}\lceil d/2 \rceil$ -neighborly.

Finally, we discuss the case of $d = 2k - 1$, $i = k$. Since $B_n^{2k-1,k} = \Delta_n^{2k-1} \setminus B_n^{2k-1,k-1}$, we obtain from the previous paragraph and Lemma 2.3 that $B_n^{2k-1,k}$ is a combinatorial $(2k-1)$ -ball that is $\text{cs-}k$ -neighborly and k -stacked. This completes the proof. \square

It now remains to show that $B_{n+1}^{d,\lceil d/2 \rceil-1}$ is a subcomplex of Δ_{n+1}^d . (Recall that our assumptions include that $B_m^{d-1,\lceil (d-1)/2 \rceil-1}$ is a subcomplex of Δ_m^{d-1} for all $m \geq d$ and that $B_n^{d,\lceil d/2 \rceil-1}$ is a subcomplex of Δ_n^d .) To facilitate the proof of this result, we rely on the following lemma; its proof is an immediate consequence of the definition of $B_n^{d,i}$, Lemma 2.1, and the nesting property.

Lemma 3.4. *Under the assumptions of Lemma 3.3, for all $d \geq 2$ and $0 \leq i \leq \lfloor d/2 \rfloor$,*

$$\partial B_n^{d,i} = \left(\partial B_{n-1}^{d-1,i} * v_n \right) \cup \left(\partial(-B_{n-1}^{d-1,i-1}) * (-v_n) \right) \cup \left(B_{n-1}^{d-1,i} \setminus (-B_{n-1}^{d-1,i-1}) \right).$$

A combinatorial sphere Δ is called k -stacked if it is the boundary complex of a combinatorial k -stacked ball.

Corollary 3.5. *Under the assumptions of Lemma 3.3, for all $k \geq 1$ and $n \geq 2k + 1$, $\partial B_n^{2k,k} = \Delta_n^{2k-1}$. In particular, Δ_n^{2k-1} is k -stacked.*

Proof: By definition, the complex Δ_n^{2k-1} is obtained from $\Delta_{n-1}^{2k-1} = B_{n-1}^{2k-1,k-1} \cup B_{n-1}^{2k-1,k}$ by replacing $\pm B_{n-1}^{2k-1,k-1}$ with $\pm(\partial B_{n-1}^{2k-1,k-1} * v_n)$. On the other hand, Lemma 3.4 and the fact that $\partial B_n^{2k-1,k} = \partial B_n^{2k-1,k-1}$ yield that

$$\partial B_n^{2k,k} = \left(\partial B_{n-1}^{2k-1,k-1} * v_n \right) \cup \left(\partial(-B_{n-1}^{2k-1,k-1}) * (-v_n) \right) \cup \left(B_{n-1}^{2k-1,k} \setminus (-B_{n-1}^{2k-1,k-1}) \right).$$

Hence $\partial B_n^{2k,k} = \Delta_n^{2k-1}$. Since by Lemma 3.3, the ball $B_n^{2k,k}$ is k -stacked, the sphere Δ_n^{2k-1} is also k -stacked. \square

We are now in a position to show that the ball $B_n^{d, \lfloor d/2 \rfloor - 1}$ is indeed a subcomplex of Δ_n^d . This will require one additional lemma.

Lemma 3.6. *Under the assumptions of Lemma 3.3, for all $d \geq 2$, $n \geq d+1$, and $0 \leq i \leq j \leq \lfloor d/2 \rfloor$, the following inclusion holds: $B_n^{d-1,i} \subseteq \partial B_n^{d,j}$.*

Proof: The proof is by induction on d . We start by checking the statement in the following base cases.

1. If $j = 0$ (and hence $i = 0$), then for any $d \geq 2$,

$$\partial B_n^{d,0} = \partial \overline{\{-v_1, v_{n-d+1}, \dots, v_n\}} \supseteq \overline{\{-v_1, v_{n-d+2}, \dots, v_n\}} = B_n^{d-1,0}.$$

2. If $i = 0$, then by Lemmas 3.4 and 3.5, for any $j > 0$ and $d \geq 2j$,

$$\partial B_n^{d,j} \supseteq \partial B_{n-1}^{d-1,j} * v_n \supseteq \dots \supseteq \partial B_{n-d+2j}^{2j,j} * \overline{\{v_{n-d+2j+1}, \dots, v_n\}} = \Delta_{n-d+2j}^{2j-1} * \overline{\{v_{n-d+2j+1}, \dots, v_n\}}.$$

Since $B_n^{d-1,0} = B_{n-d+2j}^{2j-1,0} * \overline{\{v_{n-d+2j+1}, \dots, v_n\}}$ and $\Delta_{n-d+2j}^{2j-1} \supseteq B_{n-d+2j}^{2j-1,0}$, we conclude that $B_n^{d-1,0} \subseteq \partial B_n^{d,j}$.

3. If $j \geq 1$ and $d = 2j$, then Lemma 3.5 implies that for any $i \leq j$, $\partial B_n^{2j,j} = \Delta_n^{2j-1} \supseteq B_n^{2j-1,i}$.

For the inductive step we can thus assume that $1 \leq j < d/2$ (equivalently, that $1 \leq j \leq \lfloor (d-1)/2 \rfloor$), and that the statement holds for $d-1$ and all $0 \leq i \leq j \leq \lfloor (d-1)/2 \rfloor$. Then

$$\begin{aligned} B_n^{d-1,i} &= \left(B_{n-1}^{d-2,i} * v_n \right) \cup \left((-B_{n-1}^{d-2,i-1}) * (-v_n) \right) \\ &\stackrel{(*)}{\subseteq} \left(\partial B_{n-1}^{d-1,j} * v_n \right) \cup \left((-\partial B_{n-1}^{d-1,j-1}) * (-v_n) \right) \\ &\stackrel{(**)}{\subseteq} \partial B_n^{d,j}. \end{aligned}$$

Here $(*)$ follows from the inductive hypothesis and $(**)$ follows from Lemma 3.4. This completes the proof. \square

An immediate corollary of Lemma 3.6 is that $B_{n+1}^{d, \lfloor d/2 \rfloor - 1} \subseteq \Delta_{n+1}^d$:

Corollary 3.7. For all $d \geq 2$ and $n \geq d + 1$,

$$B_{n+1}^{d, \lceil d/2 \rceil - 1} \subseteq \left(\partial B_n^{d, \lceil d/2 \rceil - 1} * v_{n+1} \right) \cup \left(\partial(-B_n^{d, \lceil d/2 \rceil - 1}) * (-v_{n+1}) \right) \subseteq \Delta_{n+1}^d.$$

Proof: The first inclusion is by definition of $B_n^{d, \lceil d/2 \rceil - 1}$ and Lemma 3.6, and the second inclusion is by definition of Δ_{n+1}^d . \square

Corollary 3.7 together with Lemma 3.3 completes the proof that the construction described in Definition 3.2 never gets stuck and that for every $d \geq 2$ and $n \geq d + 1$ it outputs a cs combinatorial d -sphere with vertex set V_n that is cs- $\lceil d/2 \rceil$ -neighborly. We summarize this in the theorem below:

Theorem 3.8. For all $d \geq 2$ and $n \geq d + 1$, the complex Δ_n^d is a cs combinatorial d -sphere with vertex set V_n that is cs- $\lceil d/2 \rceil$ -neighborly.

Remark 3.9. Let $1 \leq i \leq \lceil d/2 \rceil$. Since by nesting property,

$$B_n^{d, i-1} \subseteq B_n^{d, \lceil d/2 \rceil - 1} \cup \left(-B_n^{d, \lceil d/2 \rceil - 1} \right),$$

it follows from Corollary 3.7 that $B_n^{d, i-1} \subseteq \Delta_n^d$. This allows us to construct cs- i -neighborly spheres that are not cs- $(i + 1)$ -neighborly for all $1 \leq i \leq \lceil d/2 \rceil$. Indeed, Lemmas 3.1 and 3.3 along with Theorem 3.8 imply that the complex $\Delta_{n+1}^{d, i}$ obtained from Δ_n^d by replacing the subcomplexes $\pm B_n^{d, i-1}$ with $\pm(\partial B_n^{d, i-1} * v_{n+1})$ is a cs combinatorial d -sphere that is cs- i -neighborly. To see that $\Delta_{n+1}^{d, i}$ is not cs- $(i + 1)$ -neighborly, note that the inductive proof of Lemma 3.3 in fact shows that while $B_n^{d, i-1}$ is cs- $(i - 1)$ -neighborly, it is not cs- i -neighborly. In particular, $\partial B_n^{d, i-1}$ is not cs- i -neighborly, and hence Δ_{n+1}^d is not cs- $(i + 1)$ -neighborly.

4 Properties

It is our hope that the spheres Δ_n^d constructed in this paper will find many other applications. With this in mind, in this section we discuss several additional properties that these spheres possess. One important property is that certain face links in Δ_n^d are also highly cs-neighborly.

Proposition 4.1. For all $k \geq 2$ and $n \geq 2k - 1$,

$$\text{lk}(v_n v_{n+1}, \Delta_{n+1}^{2k-1}) = \Delta_{n-1}^{2k-3} = \text{lk}(v_n v_{n+1} v_{n+2}, \Delta_{n+2}^{2k}).$$

Proof: The assertion holds if $n = 2k - 1$ since in this case all complexes appearing in the statement are the boundary complexes of cross-polytopes. Thus assume $n \geq 2k$. By definition of Δ_{n+1}^{2k-1} , Lemma 3.4, and Lemma 3.5,

$$\text{lk}(v_n v_{n+1}, \Delta_{n+1}^{2k-1}) = \text{lk}(v_n, \partial B_n^{2k-1, k-1}) = \partial B_{n-1}^{2k-2, k-1} = \Delta_{n-1}^{2k-3}.$$

Similarly,

$$\text{lk}(v_n v_{n+1} v_{n+2}, \Delta_{n+2}^{2k}) = \text{lk}(v_n v_{n+1}, \partial B_{n+1}^{2k, k-1}) = \text{lk}(v_n, \partial B_n^{2k-1, k-1}) = \partial B_n^{2k-2, k-1} = \Delta_{n-1}^{2k-3}.$$

This completes the proof. \square

In view of Proposition 4.1, it is natural to ask whether the link of v_{n+1} in Δ_{n+1}^{2k} is Δ_n^{2k-1} , and, more generally, whether Δ_{n+1}^{2k} is a suspension. This is especially relevant to our discussion since suspensions of cs- k -neighborly spheres are also cs- k -neighborly; in particular, both complexes $\Sigma\Delta_n^{2k-1}$ and Δ_{n+1}^{2k} are cs combinatorial $2k$ -spheres that are cs- k -neighborly w.r.t V_{n+1} . Are they isomorphic? The following proposition shows that the answer is no.

Proposition 4.2. *If $n \geq 2k + 1$, then the complex Δ_{n+1}^{2k} is not a suspension. In particular, Δ_{n+1}^{2k} is not isomorphic to $\Sigma\Delta_n^{2k-1}$.*

Proof: We will rely on the following three observations.

1. If $\Delta = \Sigma\Gamma$ and $F \in \Gamma$, then $\text{lk}(F, \Delta) = \Sigma\text{lk}(F, \Gamma)$.
2. For $k \geq 2$ and $n \geq 2k + 1$, Δ_{n-2}^{2k-3} is not a suspension.

To see this, recall that Δ_{n-2}^{2k-3} is cs- $(k-1)$ -neighborly. Now, a cs $(2k-4)$ -sphere that is not the boundary of the cross-polytope can be at most cs- $(k-2)$ -neighborly. Hence, if Δ_{n-2}^{2k-3} were a suspension, it would also be at most cs- $(k-2)$ -neighborly.

3. For $\ell \geq 1$ and $m \geq 2\ell + 1$, the complex $\partial B_m^{2\ell, \ell-1}$ is not a cs complex.

If $\partial B_m^{2\ell, \ell-1}$ were a cs complex, then the link $\text{lk}(v_m, \partial B_m^{2\ell, \ell-1}) = \partial B_{m-1}^{2\ell-1, \ell-1}$ would coincide with $-\text{lk}(-v_m, \partial B_m^{2\ell, \ell-1}) = -\partial(-B_{m-1}^{2\ell-1, \ell-2}) = \partial B_{m-1}^{2\ell-1, \ell-2}$. Since the balls $B_{m-1}^{2\ell-1, \ell-1}$ and $B_{m-1}^{2\ell-1, \ell-2}$ share no common facets, their union would then be a $(2\ell-1)$ -sphere strictly contained in the $(2\ell-1)$ -sphere $\Delta_m^{2\ell-1}$, which is impossible.

We are now ready to prove the proposition. If $k = 1$, then Δ_{n+1}^2 is obtained from $\partial\mathcal{C}_3^*$ by symmetric stacking, and so Δ_{n+1}^2 is not a suspension for $n \geq 3$. Thus let $k \geq 2$, and assume by contradiction that Δ_{n+1}^{2k} is a suspension with suspension vertices $\pm v_i$ for some $i \leq n + 1$.

If $i < n - 1$, then by Observation 1, the link of $v_{n-1}v_nv_{n+1}$ in Δ_{n+1}^{2k} must be a suspension. However, by Proposition 4.1, $\text{lk}(v_{n-1}v_nv_{n+1}, \Delta_{n+1}^{2k}) = \Delta_{n-2}^{2k-3}$, which according to Observation 2 is not a suspension.

If $i = n + 1$, then $\text{lk}(v_{n+1}, \Delta_{n+1}^{2k})$ must equal $\text{lk}(-v_{n+1}, \Delta_{n+1}^{2k})$. In particular, $\text{lk}(v_{n+1}, \Delta_{n+1}^{2k}) = \partial B_n^{2k, k-1}$ must be a cs complex, which contradicts Observation 3.

If $i = n$, then by Observation 1, $\text{lk}(v_{n+1}, \Delta_{n+1}^{2k}) = \partial B_n^{2k, k-1}$ must be a suspension with suspension vertices $\pm v_n$. However, since $-B_{n-1}^{2k-1, k-2}$ is strictly contained in $B_{n-1}^{2k-1, k-1}$, the complexes $\text{lk}(v_n, \partial B_n^{2k, k-1}) = \partial B_{n-1}^{2k-1, k-1}$ and $\text{lk}(-v_n, \partial B_n^{2k, k-1}) = \partial(-B_{n-1}^{2k-1, k-2})$ are not equal.

Finally, if $i = n - 1$, then by Observation 1, we must have

$$\text{lk}(v_{n-1}, \text{lk}(v_nv_{n+1}, \Delta_{n+1}^{2k})) = \text{lk}(-v_{n-1}, \text{lk}(v_nv_{n+1}, \Delta_{n+1}^{2k})).$$

The proof of Proposition 4.1 implies that the former complex is Δ_{n-2}^{2k-3} while the latter complex is $\partial(-B_{n-2}^{2k-2, k-2})$. These complexes cannot be equal since one is cs and the other one is non-cs. \square

Remark 4.3. The ball $B_{n+1}^{2k, k}$ is “special” in the following sense: while $B_{n+1}^{d, i} \subseteq \Delta_{n+1}^d$ for all $d \geq 1$, $i \leq \lceil d/2 \rceil - 1$, and $n \geq d$ (see Remark 3.9), the ball $B_{n+1}^{2k, k}$ is **not** a subcomplex of Δ_{n+1}^{2k} for $n > 2k$. Indeed, a straightforward computation using that $\Delta_n^{2k-1} = B_n^{2k-1, k} \cup B_n^{2k-1, k-1}$ along with the definition of $B_{n+1}^{2k, k}$ implies that $B_{n+1}^{2k, k} \cup (-B_{n+1}^{2k, k}) = \Sigma\Delta_n^{2k-1}$. Our claim that $B_{n+1}^{2k, k} \not\subseteq \Delta_{n+1}^{2k}$ (for $n \geq 2k + 1$) then follows from Proposition 4.2. However, in the case of $n = 2k$, the same computation shows that $B_{2k+1}^{2k, k} \cup (-B_{2k+1}^{2k, k}) = \Sigma\Delta_{2k}^{2k-1} = \Sigma(\partial\mathcal{C}_{2k}^*) = \partial\mathcal{C}_{2k+1}^* = \Delta_{2k+1}^{2k}$.

To close this section, we show that while Δ_{n+1}^{2k} is not the suspension of Δ_n^{2k-1} , the complexes Δ_n^{2k} and Δ_n^{2k-1} are closely related:

Proposition 4.4. *Let $k \geq 2$ and $n \geq 2k + 1$. Then $\Delta_n^{2k} \supseteq \Delta_n^{2k-1}$.*

Proof: To prove the statement, it suffices to construct a combinatorial $2k$ -ball D_n that satisfies the following properties:

- $D_n \supseteq -B_n^{2k,k-1}$.
- D_n and $-D_n$ share no common facets and their union is Δ_n^{2k} ; in particular, $\pm D_n \subseteq \Delta_n^{2k}$.
- $\partial D_n = \partial(-D_n) = \Delta_n^{2k-1}$, and so $\Delta_n^{2k-1} \subseteq \Delta_n^{2k}$.

We construct D_n by induction on n . For the base case, define $D_{2k+1} := B_{2k+1}^{2k,k}$. This ball has all the desired properties: this follows from Lemma 3.3, Corollary 3.5, and Remark 4.3.

For the inductive step, assume that there exists a combinatorial $2k$ -ball D_n that satisfies all of the above properties. In particular, $-D_n$ contains $B_n^{2k,k-1}$ as a subcomplex. Recall also that $\partial B_n^{2k,k-1} \supseteq B_n^{2k-1,k-1}$ (see Lemma 3.6). Thus, the following complex is well-defined:

$$D_{n+1} := \left((-D_n) \setminus B_n^{2k,k-1} \right) \cup \left((-B_n^{2k-1,k-1}) * (-v_{n+1}) \right) \cup \left((\partial B_n^{2k,k-1}) \setminus B_n^{2k-1,k-1} * v_{n+1} \right). \quad (4.1)$$

This definition along with the inductive assumption that D_n and $-D_n$ share no common facets implies that D_{n+1} and $-D_{n+1}$ share no common facets either. Further, since $\partial B_n^{2k,k-1}$ contains $B_n^{2k-1,k-2}$ (see Lemma 3.6) and since $B_n^{2k-1,k-2}$ and $B_n^{2k-1,k-1}$ share no common facets (this follows from the nesting property), equation (4.1) guarantees that

$$D_{n+1} \supseteq \left((-B_n^{2k-1,k-1}) * (-v_{n+1}) \right) \cup \left(B_n^{2k-1,k-2} * v_{n+1} \right) = -B_{n+1}^{2k,k-1}.$$

Recall that Δ_{n+1}^{2k} is obtained from Δ_n^{2k} by replacing $\pm B_n^{2k,k-1}$ with $\pm(\partial B_n^{2k,k-1} * v_{n+1})$. This together with the definition of D_{n+1} and the inductive hypothesis asserting that $D_n \cup (-D_n) = \Delta_n^{2k}$ implies that $D_{n+1} \cup (-D_{n+1}) = \Delta_{n+1}^{2k}$.

It only remains to show that D_{n+1} is a combinatorial $2k$ -ball with $\partial D_{n+1} = \partial(-D_{n+1}) = \Delta_{n+1}^{2k-1}$. We use eq. (4.1) and the inductive hypothesis that $\partial(-D_n) = \Delta_n^{2k-1}$. First, the boundary of $\partial B_n^{2k,k-1} * v_{n+1}$ coincides with the boundary of $B_n^{2k,k-1}$. Thus, replacing the subcomplex $B_n^{2k,k-1}$ of the combinatorial ball $-D_n$ with $\partial B_n^{2k,k-1} * v_{n+1}$ results in a combinatorial ball D'_{n+1} that has the same boundary as $-D_n$. The ball D'_{n+1} has v_{n+1} as an interior vertex whose link in D'_{n+1} is $\partial B_n^{2k,k-1}$. Now, $B_n^{2k-1,k-1}$ is contained in both $\partial B_n^{2k,k-1}$ and $\partial D'_{n+1} = \partial(-D_n) = \Delta_n^{2k-1}$. Since v_{n+1} is an interior vertex of D'_{n+1} , removing $B_n^{2k-1,k-1} * v_{n+1}$ from D'_{n+1} results in a combinatorial ball D''_{n+1} whose boundary is obtained from that of $-D_n$ by replacing $B_n^{2k-1,k-1}$ with $\partial B_n^{2k-1,k-1} * v_{n+1}$. Finally, the balls $(-B_n^{2k-1,k-1}) * (-v_{n+1})$ and D''_{n+1} intersect along $-B_n^{2k-1,k-1}$, which is a subcomplex of their boundaries. This yields that $D_{n+1} = D''_{n+1} \cup \left((-B_n^{2k-1,k-1}) * (-v_{n+1}) \right)$ is a combinatorial ball. Further, the boundary of D_{n+1} is obtained from the boundary of D''_{n+1} by replacing $-B_n^{2k-1,k-1}$ with $\partial(-B_n^{2k-1,k-1}) * (-v_{n+1})$. We conclude that $\partial D_{n+1} = \Delta_{n+1}^{2k-1}$. \square

Observe that the $2k$ -ball D_n constructed in the proof of Proposition 4.4 is k -stacked. (Indeed, since the boundary of D_n is a cs - k -neighborly complex w.r.t. V_n , no face of D_n of dimension $\leq k-1$ is an interior face of D_n , and so all interior faces of D_n have dimension $\geq k = 2k - k$.) Thus one curious consequence of Proposition 4.4 and Corollary 3.5 is that for $n > 2k + 1$, Δ_n^{2k-1} is the boundary complex of at least four distinct k -stacked $2k$ -balls, namely, D_n , $-D_n$, $B_n^{2k,k}$, and $-B_n^{2k,k}$.

5 Closing remarks and questions

We close with a few open questions.

The spheres Δ_n^d we constructed here are cs combinatorial spheres that are cs- $\lceil d/2 \rceil$ -neighborly. Hence, according to [13], for $d \geq 3$ and $n \geq d + 3$, they are not the boundary complexes of cs polytopes. In fact, it follows from results of Pfeifle [19, Chapter 10] that they are not even *cs fans* (at least for d big enough). However, these spheres might still possess some additional “liked by all” properties:

Problem 5.1. *Let $d \geq 3$ and $n \geq d + 3$. Are the spheres Δ_n^d vertex decomposable or at least shellable? Are they realizable as (non-cs) fans? Are they even realizable as boundary complexes of (non-cs) polytopes?*

It is well-known that there are many (non-cs) neighborly polytopes and neighborly spheres. For example, the number of combinatorial types of $\lceil d/2 \rceil$ -neighborly $(d + 1)$ -polytopes with n vertices is at least $n^{\frac{d-1}{2}n(1+o(1))}$ for $d > 1$ and $n \rightarrow \infty$, see [18, Section 6]. Are there many cs d -spheres that are cs- $\lceil d/2 \rceil$ -neighborly?

Problem 5.2. *Find many new constructions of cs combinatorial d -spheres that are cs- $\lceil d/2 \rceil$ -neighborly.*

Finally, it is worth mentioning that in a contrast with cs combinatorial spheres, there exist cs combinatorial $2k$ -manifolds with $n > 2(2k + 1)$ vertices that are cs- $(k + 1)$ -neighborly. (The interest in such complexes arises in part from Sparla’s conjecture [20, 21] that posits an upper bound on the Euler characteristic of cs combinatorial $2k$ -manifolds with $2n$ vertices; see [17] for some results on this conjecture.) One construction of such an infinite family is given in [8]: for each $k \geq 1$, it produces a cs triangulation of the product of two k -dimensional spheres with $4k + 4$ vertices that is cs- $(k + 1)$ -neighborly. For additional constructions in low dimensions, see [3, 11, 20].

Problem 5.3. *Find new constructions of (infinite families of) cs combinatorial $2k$ -manifolds that are cs- $(k + 1)$ -neighborly.*

Acknowledgments

We are grateful to the referee for suggesting an approach that greatly simplified and streamlined our main definitions and presentation.

References

- [1] R. M. Adin. *Combinatorial structure of simplicial complexes with symmetry*. PhD thesis, Hebrew University, Jerusalem, 1991.
- [2] A. Björner. Topological methods. In *Handbook of combinatorics, Vol. 1, 2*, pages 1819–1872. Elsevier Sci. B. V., Amsterdam, 1995.
- [3] F. Effenberger. *Hamiltonian submanifolds of regular polytopes*. Dissertation, University of Stuttgart, Stuttgart, 2010.
- [4] B. Grünbaum. *Convex polytopes*, volume 221 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2003. Prepared and with a preface by Volker Kaibel, Victor Klee and Günter M. Ziegler.

- [5] J. F. P. Hudson. *Piecewise linear topology*. University of Chicago Lecture Notes. W. A. Benjamin Inc., New York-Amsterdam, 1969.
- [6] W. Jockusch. An infinite family of nearly neighborly centrally symmetric 3-spheres. *J. Combin. Theory Ser. A*, 72(2):318–321, 1995.
- [7] S. Klee, E. Nevo, I. Novik, and H. Zheng. A lower bound theorem for centrally symmetric simplicial polytopes. *Discrete Comput. Geom.*, 61:541–561, 2019.
- [8] S. Klee and I. Novik. Centrally symmetric manifolds with few vertices. *Adv. Math.*, 229:487–500, 2012.
- [9] W. B. R. Lickorish. Simplicial moves on complexes and manifolds. In *Proceedings of the Kirbyfest (Berkeley, CA, 1998)*, volume 2 of *Geom. Topol. Monogr.*, pages 299–320. Geom. Topol. Publ., Coventry, 1990.
- [10] N. Linial and I. Novik. How neighborly can a centrally symmetric polytope be? *Discrete Comput. Geom.*, 36:273–281, 2006.
- [11] F. Lutz. *Triangulated manifolds with few vertices and vertex-transitive group actions*. Dissertation, Technischen Universität Berlin, Berlin, 1999.
- [12] P. McMullen. The maximum numbers of faces of a convex polytope. *Mathematika*, 17:179–184, 1970.
- [13] P. McMullen and G. C. Shephard. Diagrams for centrally symmetric polytopes. *Mathematika*, 15:123–138, 1968.
- [14] P. McMullen and D. W. Walkup. A generalized lower-bound conjecture for simplicial polytopes. *Mathematika*, 18:264–273, 1971.
- [15] S. Murai and E. Nevo. On the generalized lower bound conjecture for polytopes and spheres. *Acta Math.*, 210(1):185–202, 2013.
- [16] I. Novik. Remarks on the upper bound theorem. *J. Combin. Theory Ser. A*, 104(1):201–206, 2003.
- [17] I. Novik. On face numbers of manifolds with symmetry. *Adv. Math.*, 192:183–208, 2005.
- [18] A. Padrol. Many neighborly polytopes and oriented matroids. *Discrete Comput. Geom.*, 50(4):865–902, 2013.
- [19] J. Pfeifle. *Extremal Constructions for Polytopes and Spheres*. Dissertation, Technischen Universität Berlin, Berlin, 2003.
- [20] E. Sparla. *Geometrische und kombinatorische Eigenschaften triangulierter Mannigfaltigkeiten*. Berichte aus der Mathematik. [Reports from Mathematics]. Verlag Shaker, Aachen, 1997. Dissertation, Universität Stuttgart, Stuttgart, 1997.
- [21] E. Sparla. An upper and a lower bound theorem for combinatorial 4-manifolds. *Discrete Comput. Geom.*, 19:575–593, 1998.
- [22] R. P. Stanley. The upper bound conjecture and Cohen-Macaulay rings. *Studies in Applied Math.*, 54:135–142, 1975.
- [23] R. P. Stanley. The number of faces of a simplicial convex polytope. *Adv. Math.*, 35:236–238, 1980.