

## Math 135: Homework 6

Due Thursday, February 10

1. Fix a real constant  $\alpha$ . Show that  $\sum_{k=0}^{\infty} \frac{\sin \alpha k}{2^k}$  converges. Evaluate the sum exactly (in terms of  $\alpha$ ) by using the fact that  $\sin \alpha k = \text{Im}(e^{i\alpha k})$ . **Hints:** You may assume that all of the series we've been discussing work equally well with complex numbers as with real numbers. Also, note that for any real numbers  $a$  and  $b$ , we have  $\frac{1}{a+ib} = \frac{a}{a^2+b^2} - i\frac{b}{a^2+b^2}$ .

2. Fix an interval  $I = (a, b)$ . Let  $y_1(x)$  and  $y_2(x)$  be solutions of  $y'' + p(x)y' + q(x)y = 0$  on  $I$ , where  $p(x), q(x)$  are continuous functions on  $I$ . Show that if there is a point in  $I$  where both  $y_1$  and  $y_2$  vanish or where both have maxima or minima, then one of  $y_1$  and  $y_2$  is a multiple of the other.

3. In the following, you are given a differential equation and one solution of it. Use *reduction of order* to find the general solution. We will discuss this method in class on Monday or Tuesday: try a solution of the form  $y_2(x) = u(x)y_1(x)$  and solve for the unknown function  $u(x)$ . When you plug  $y_2(x)$  into the differential equation, the  $u$  terms will cancel, leaving an equation of the form  $f(x)u'' + g(x)u' = 0$ . This is a first order linear equation (and also separable) in the function  $u'$ , so solve for  $u'$ . Integrate to get  $u$ , and then you can get the second solution  $y_2 = uy_1$ .

(This technique is also discussed in the book: in the constant coefficient case, see Lesson 20C, and with nonconstant coefficients but using slightly different notation, see Lesson 23B.)

(a)  $x^2y'' - x(x+2)y' + (x+2)y = 0$ ,  $y_1(x) = x$ .

(b)  $xy'' - (x+2)y' + 2y = 0$ ,  $y_1(x) = e^x$ .