

Math 308H
Spring 2016
Midterm 2
May 13, 2016

Name (Print): _____

Time Limit: 50 Minutes

Instructor _____

This exam contains 5 pages (including this cover page) and 4 problems. Check to see if any pages are missing. Enter all requested information on the top of this page, and put your initials on the top of every page, in case the pages become separated.

You may use a sheet of notes, 8.5×11 and handwritten by you, but no other devices, books, or notes are permitted.

Unless otherwise stated, you are required to show your work on each problem on this exam. The following rules apply:

- **Organize your work**, in a reasonably neat and coherent way, in the space provided. Work scattered all over the page without a clear ordering will receive very little credit.
- **Mysterious or unsupported answers will not receive full credit.** A correct answer, unsupported by calculations, explanation, or algebraic work will receive no credit; an incorrect answer supported by substantially correct calculations and explanations might still receive partial credit.
- If you need more space, use the back of the pages; clearly indicate when you have done this.

Problem	Points	Score
1	9	
2	10	
3	11	
4	10	
Total:	40	

Do not write in the table to the right.

1. Suppose V is a subset of \mathbb{R}^n . For V to be a subspace, we require the three conditions.
 1. $\mathbf{0} \in V$.
 2. If $\mathbf{u}, \mathbf{v} \in V$, then $\mathbf{u} + \mathbf{v} \in V$. (closed under addition.)
 3. If $\mathbf{u} \in V$, $r \in \mathbb{R}$, then $r\mathbf{u} \in V$. (closed under scalar multiplication.)

For each of the following subsets V , decide whether each condition is satisfied. For example, you might say that conditions 1 and 3 hold, but condition 2 does not.

- (a) (3 points) Let $V \subseteq \mathbb{R}^3$ be the set of all vectors of the form $\begin{bmatrix} x \\ x^2 \\ x^3 \end{bmatrix}$ for $x \in \mathbb{R}$.

- (b) (3 points) Let $V \subseteq \mathbb{R}^2$ be the set of all vectors of the form $\begin{bmatrix} x \\ 2x \end{bmatrix}$ for $x \geq 0$.

- (c) (3 points) Let $V \subseteq \mathbb{R}^3$ be the set of all vectors of the form $\begin{bmatrix} x \\ y \\ z \end{bmatrix}$ where y and z are any real numbers and x is an integer.

2. Read each of the following statements carefully, and decide whether it is true or false. You are not required to justify your answers, but I recommend justifying them to yourself.

(a) (2 points) If A and B are equivalent matrices, then $\text{col}(A) = \text{col}(B)$.

(b) (2 points) If A and B are $n \times n$ diagonal matrices, then $AB = BA$.

(c) (2 points) Suppose $m > n$ and $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is a linear transformation with $T(\mathbf{x}) = A\mathbf{x}$. Then for any $\mathbf{b} \in \mathbb{R}^n$, the system $A\mathbf{x} = \mathbf{b}$ is consistent.

(d) (2 points) If A and B are $n \times n$ matrices with AB invertible, then A and B are also invertible.

(e) (2 points) Suppose $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is a linear transformation, and $V \subseteq \mathbb{R}^n$ is a subspace. Then the set of all $\mathbf{x} \in \mathbb{R}^m$ with $T(\mathbf{x}) \in V$ is a subspace of \mathbb{R}^m .

3. (a) (3 points) Suppose A and B are $n \times n$ matrices and that A , B , and $A + B$ are invertible. Simplify the following expression as much as possible.

$$(AB)^{-1}BA + (A + B)^T(A^T + B^T)^{-1}$$

- (b) (3 points) Solve the following equation for the matrix X , where A, B and C are all $n \times n$ matrices and A and B are invertible.

$$AXB = C.$$

- (c) (3 points) Give an example of an onto linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ satisfying
- $$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ 2 \end{bmatrix}.$$

- (d) (2 points) Give an example of two 2×2 invertible matrices A and B whose sum $A + B$ is not invertible.

4. Consider the matrix

$$A = \begin{bmatrix} 0 & 2 & 0 & -2 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 2 & 3 \end{bmatrix}.$$

(a) (5 points) Find a basis for $\text{null}(A)$.

(b) (5 points) Find a basis for $\text{col}(A)$.