

## DERIVATIVE OF $\Lambda$

JUNE 24, 2010

Let  $\Lambda_\gamma$  denote the Dirichlet-to-Neumann map for an electrical network with conductivity  $\gamma$ . By way of the Kirchhoff matrix  $K = (\kappa_{ij})$ , consider the space of conductivities to be a subset of  $\mathbb{R}^{N \times N}$ , where  $N$  is the number of vertices. We denote the map from  $\gamma$  to  $\Lambda_\gamma$  by  $L$ . In this note we compute the directional derivative  $D_\epsilon L$  of  $L$ .

A direction in this context is represented by a matrix  $\epsilon$ , with arbitrary real entries, which is symmetric and has row sum 0.

**Lemma 0.1.** *Let*

$$K = \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}, \quad \epsilon = \begin{bmatrix} \epsilon_A & \epsilon_B \\ \epsilon_B^T & \epsilon_C \end{bmatrix},$$

where  $\epsilon$  is symmetric and  $K$  is a Kirchhoff matrix. Let

$$K(t) = \begin{bmatrix} A + t\epsilon_A & B + t\epsilon_B \\ B^T + t\epsilon_B^T & C + t\epsilon_C \end{bmatrix},$$

and

$$\Lambda(t) = [A + t\epsilon_A - (B + t\epsilon_B)(C + t\epsilon_C)^{-1}(B^T + t\epsilon_B^T)].$$

Then

$$D_\epsilon L = \Lambda'(0) = \epsilon_A - \epsilon_B C^{-1} B^T - B C^{-1} \epsilon_B^T + B C^{-1} \epsilon_C C^{-1} B^T.$$

*Proof.* Let  $C(t) = C + t\epsilon_C$ .  $C(t)(C(t))^{-1} - I$ . By the product rule  $C'(t)C(t) + C(t)(C(t)^{-1})' = 0$ , hence  $(C^{-1})'(0) = -C^{-1}\epsilon_C C^{-1}$ . Using the product rule again

$$D_\epsilon L = \Lambda'(0) = \epsilon_A - \epsilon_B C^{-1} B^T - B C^{-1} \epsilon_B^T + B C^{-1} \epsilon_C C^{-1} B^T. \quad \square$$

**Corollary 0.1.** *Let  $\phi, \psi$  be boundary functions. Let  $u, v$  be  $\gamma$  harmonic functions with boundary values  $\phi, \psi$ . Then*

$$\phi^T D_\epsilon L \psi = \sum_{i \neq j} \epsilon_{ij} (u_i - u_j)(v_i - v_j).$$

*Proof.* The interior values of the  $\gamma$ -harmonic function with boundary values  $\phi$  is  $-C^{-1}B^T\phi$  and with boundary values  $\psi$  is  $-C^{-1}B^T\psi$ . Hence by computation

$$\phi^T D_\epsilon L \psi = [\phi^T \quad -\phi^T B C^{-1}] \begin{bmatrix} \epsilon_A & \epsilon_B \\ \epsilon_B^T & \epsilon_C \end{bmatrix} \begin{bmatrix} \psi \\ -C^{-1}B^T\psi \end{bmatrix} = \sum_{i \neq j} \epsilon_{ij} (u_i - u_j)(v_i - v_j)$$

